

NOTICE OF PESTICIDE REGISTRATION
 REREISTRATION

NAME OF PESTICIDE PRODUCT

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NO. (10)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Monteclay Chemical Company
P.O. Box 5224
Fresno, CA 93755

161 157430
18 1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

 ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Henry M. Sandy

DATE

8/26/55

AUG 26 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

481421

NORDOX

FUNGICIDE

FUNGICIDE

ACTIVE INGREDIENT:

COPPER (IN CUPROUS OXIDE (Cu₂O)) 50%

INERT INGREDIENT: 50%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED.

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water or milk and induce vomiting by pushing back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes: Flush eyes with water for at least 15 minutes, holding eyelids apart. Seek medical attention immediately.

See left side panel for additional precautionary statements.

FUNGICIDE



FUNGICIDE

ACTIVE INGREDIENT:

COPPER (IN CUPROUS OXIDE (Cu₂O) 50%

INERT INGREDIENT: 50%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED.

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water or milk and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes: Flush eyes with water for at least 15 minutes, holding eye-lids apart. Seek medical attention immediately.

See left side panel for additional precautionary statements.

NORDOX A/S
Monterey Chemical Co.
P.O. Box 5317
Fresno, CA 93755

EPA Reg. No. _____ Net Contents _____
EPA Est. No. _____

Manufactured by Nordox A/S, Oslo 6, Norway

MONTEREY CHEMICAL

**HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid breathing vapors or dust. Avoid contact with skin, eyes and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Do not store near feed or food stuffs.

ENVIRONMENTAL HAZARDS

Do not apply directly to any open bodies of water. Do not contaminate water by the cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinseate that cannot be used according to label directions must be disposed of according to Federal, State, or local procedures.

3. CONTAINER DISPOSAL Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

This is to certify that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph applicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there shall be no warranty, expressed or implied, by this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product.

Harmful if swallowed. Avoid breathing vapors or dust. Avoid contact with skin, eyes and clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Do not store near feed or food stuffs.

ENVIRONMENTAL HAZARDS

Do not apply directly to any open bodies of water. Do not contaminate water by the cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food, or feed by storage or disposal.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinsate that cannot be used according to label directions must be disposed of according to Federal, State, or local procedures.

3. CONTAINER DISPOSAL Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

It is critical that this product be used and mixed only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages: (i) arising from any use of this product in a manner or for a purpose not recommended in its label directions, or from mixing this product before use with any substance except as recommended by the product's label, (ii) arising from handling or storage in violation of label instructions, (iii) for all indirect special or consequential damages, (iv) when not reported to this company within one year of discovery, and (v) arising from product not used within the label-designated shelf life or four years from date of purchase, whichever first occurs. THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.

GENERAL INSTRUCTIONS

Use **NORDOX** as noted. **NORDOX** is adaptable to spraying from all types of spray equipment. Depending on the equipment used and the specific crop, the volume applied per acre will differ. For dilute, high volume sprays: use from 25 to 100 gallons of water per acre (GPA) for most vegetable crops, 400 to 800 GPA for fruit orchards and up to 1500 GPA as may be required for large citrus groves. For concentrate ground sprays apply from 5 to 20 GPA for most vegetable crops and 25 to 100 GPA for fruit and nut crops. For aerial spraying, 3 to 15 GPA are commonly used. Add **NORDOX** slowly to a spray tank partially filled with water. Spraying fungicides, insecticides, nutrients, etc. should be added last. Observe all cautions and limitations on label of all products used in mixtures.

The following specific instructions are based on general applications. The recommendations of State Agricultural Extension Offices should be closely followed as to timing, frequency and number of sprays per season.

TREE AND VINE CROPS

CROP	DISEASE	LATIN NAME	RATE	REMARKS
Almonds	Brown rot blossom blight	<i>Monilinia laxa</i>	Dilute: 2-5 lbs/100 gal	Dormant application: Nov 15-Jan 15
	Coryneum blight (shot hole)	<i>Coryneum carpophium</i>	Concentrate: 8-20 lbs/A Aerial: 8-20 lbs/A	Apply as bud spray at pink bud throughout popcorn
	Bacterial blast	<i>Pseudomonas syringae</i>	Dilute: 3-4 lbs/100 gal	Apply at dormant to early pink stage. Where disease is severe apply 1/4 lb at 2-week intervals post-bloom. Use 2-4 applications just prior to sprinkling. Slight leaf injury may occur.
Apples	Bacterial canker	<i>Pseudomonas</i> spp.	Dilute: 2-4 lbs/100 gal	Apply as a dormant spray
	Apple scab (black spot)	<i>Venturia inaequalis</i>	Concentrate: 3-5 lbs/A	Apply only at bud break to control primary infection.
Apricots	Brown rot blossom blight	<i>Monilinia</i> spp.	Dilute: 2-5 lbs/100 gal	Dormant application: Nov 15-Jan 15
	Coryneum blight (shot hole)	<i>Coryneum beijerinckii</i>	Concentrate: 8-20 lbs/A Aerial: 8-20 lbs/A	Apply as a bud spray through bud swell and popcorn stage. Do not apply after bloom.
Avocados	Anthraxnose	<i>Glomerella singulata</i>	Dilute: 2-3 lbs/100 gal	Apply when blossom buds open
	Scab	<i>Sphaceloma persea</i>	Concentrate: 8-10 lbs/A	Repeat at monthly intervals for 5-6 applications.
Bananas	Sigatoka	<i>Cercospora musae</i>	2.5 to 7 lbs/A	Apply every 3-4 weeks
Caneberries	Anthraxnose	<i>Gloeosporium arianasporum</i>	2 lbs/100 gal	Apply when leaf buds open. Repeat when flower buds show white and continue at 10-14 day intervals.
	Leaf and cane spot	<i>Septoria rub.</i>	2-4 lbs/100 gal	Apply delayed dormant spray in the spring and make a fall application.
	Yellow rust	<i>Kuchincola uredinis</i> <i>Phragmidium rubroideae</i>		
Cherries	Brown rot blossom blight	<i>Monilinia</i> spp.	2-5 lbs/100 gal	Apply at popcorn to full bloom stage
	Coryneum blight	<i>Coryneum beijerinckii</i>	Dilute: 4-6 lbs/100 gal	Apply before heavy rains in the fall and again in January. If disease is severe, apply a late summer spray after harvest.
	Dead bud	<i>Pseudomonas syringae</i>	Concentrate: 8-20 lbs/A	
Citrus	Melanose	<i>Phomopsis</i> spp.	Dilute: 3/4-3 lbs/100 gal	Florida and Texas
	Fruit rot	<i>Diplodia natalensis</i> <i>Diaplothe C. fr.</i>	Concentrate: 4-20 lbs/A Aerial: 4-20 lbs/A	Apply as pre-bloom and post bloom sprays.
	Spot anthracnose (scab)	<i>Elsinoe fawcettii</i>		
	Greasy spot (pink pitting)	<i>Mycosphaerella citri</i>		
	Brown rot	<i>Phytophthora citrophthora</i>	Dilute: 1/2-2 lbs/100 gal	Apply in the fall just before the rain season. Apply as an overall or trunk spray. Apply to the soil beneath the trees. Use with care in areas where copper injury is known to occur.
Coconuts	Black pod rot	<i>Phytophthora palmivora</i>	2-4 lbs/A	Apply on a 14-21 day schedule in high rainfall areas
Coffee	Iron spot	<i>Cercospora</i> spp.	2-8 lbs/A	Apply 3 applications at monthly intervals at the beginning of the wet season
	Pink Disease	<i>Corticium salmonicolor</i>		
	Leaf rust	<i>Hemileia vastatrix</i>	3.5-8 lbs/A	Apply as locally recommended usually at 3-4 week intervals depending upon disease severity and rainfall condition
	Leaf spot Berry spot	<i>Mycena citricola</i> <i>Lecanospora coffeicola</i>		
Cranberry	Fruit rot	<i>Ceratospora lunata</i>	8 lbs/A	Single application in late bloom. 1-2 additional applications at 10-14 day intervals may be necessary.
Currants and Gooseberries	Leaf spot (cane blight)	<i>Batryosphaeria ribis</i>	5-10 lbs/A	Make 3 applications starting after harvest, before bloom and after petal fall
Filberts	Bacterial blight	<i>Pseudomonas</i> spp. <i>Xanthomonas</i> spp.	4-5 lbs/100 gal	Apply in late August or early September. Under severe conditions apply another application when 1/4 of the leaves have dropped.
Grapes	Black rot	<i>Gulignardia bidwellii</i>	2 lbs/A	Apply the first application when clusters are 1-8 inches long. A second application should be applied just after bloom and a third before fruit maturity.
	Downy mildew	<i>Plasmopara viticola</i>		

Apples	Bacterial canker Apple scab (black spot)	<i>Pseudomonas</i> spp. <i>Venturia</i> <i>sp.</i>	Dilute: 2-4 lbs/100 gal Concentrate: 3-5 lbs/A	Apply as a dormant spray Apply only at bud break to control primary infection.
Apricots	Brown rot blossom blight Coryneum blight (shot hole)	<i>Monilinia</i> spp. <i>Coryneum beijerinckii</i>	Dilute: 2-5 lbs/100 gal Concentrate: 6-20 lbs/A Aerial: 8-20 lbs/A	Dormant application: Nov 15-Jan 15 Apply as a bud spray through bud swell and popcorn stage Do not apply after bloom
Avocados	Anthracoese Scab	<i>Glomerella cingulata</i> <i>Spenceroma persea</i>	Dilute: 2-3 lbs/100 gal Concentrate: 8-10 lbs/A	Apply when blossom buds open Repeat at monthly intervals for 5-6 applications.
Bananas	Sigatoka	<i>Cercospora musae</i>	2.5 to 4 lbs/A	Apply every 3-4 weeks
Caneberries	Anthracoese	<i>G. oosporium</i> <i>S. anthracinum</i>	2 lbs/100 gal	Apply when leaf buds open. Repeat when flower buds show white and continue at 10-14 day intervals.
	Leaf and cane spot Yellow rust	<i>Septoria</i> spp. <i>Kuchinoia uredinis</i> <i>Phragmidium rubicandae</i>	2-4 lbs/100 gal	Apply delayed dormant spray in the spring and make a fall application.
Cherries	Brown rot blossom blight Coryneum blight Dead bud	<i>Monilinia</i> spp. <i>Coryneum beijerinckii</i> <i>Pseudomonas syringae</i>	2-5 lbs/100 gal Dilute: 4-6 lbs/100 gal Concentrate: 8-20 lbs/A	Apply at popcorn to full bloom stage Apply before heavy rains in the fall and again in January. If disease is severe, apply a late summer spray after harvest.
Citrus	Melanose Fruit rot Spot anthracnose (scab) Greasy spot (pink pitting) Brown rot	<i>Phomopsis</i> spp. <i>Diplodia natalensis</i> <i>Diaporthe Citri</i> <i>Elainae fawcettii</i> <i>Mycosphaerella citri</i> <i>Phytophthora citrophthora</i>	Dilute: 3/4-3 lbs/100 gal Concentrate: 4-20 lbs/A Aerial: 4-20 lbs/A Dilute: 1/2-2 lbs/100 gal	Florida and Texas Apply as pre-bloom and post bloom sprays. Apply in the fall just before the rain season. Apply as an overall or leaf spray. Apply to the soil beneath the trees. Use with lime in areas where copper injury is known to occur.
Cocoa	Black pod rot	<i>Phytophthora palmivora</i>	2-4 lbs/A	Apply on a 14-21 day schedule in high rainfall areas.
Coffee	Iron spot Pink Disease Leaf rust Leaf spot Berry spot	<i>Cercospora</i> spp. <i>Corticium salmonicolor</i> <i>Hemileia vastatrix</i> <i>Mycena citricola</i> <i>Cercospora coffeicola</i>	2-6 lbs/A 3.5-8 lbs/A	Apply 3 applications at monthly intervals: at the beginning of the wet season Apply as locally recommended usually at 3-4 week intervals depending upon disease severity and rainfall conditions.
Cranberry	Fruit rot	<i>Ceratomyces lunata</i>	8 lbs/A	One application in late bloom; 1-2 additional applications at 10-14 day intervals may be necessary.
Currents and Gooseberries	Leaf spot (cane blight)	<i>Bathyosphaeria ribis</i>	5-10 lbs/A	Make 3 applications starting after harvest, before bloom and after petal fall.
Filberts	Bacterial blight	<i>Pseudomonas</i> spp. <i>Xanthomonas</i> spp.	4-5 lbs/100 gal	Apply in late August or early September. Under severe conditions apply another application when 3/4 of the leaves have dropped.
Grapes	Black rot Downy mildew Powdery mildew	<i>Gulignardia bidwellii</i> <i>Plasmopara viticola</i> <i>Uncinula necator</i>	2 lbs/A 2 lbs/A	Apply the first application when shoots are 6-8 inches long. A second application should be applied just after bloom and a third before fruit changes color or every 4-10 days if disease is severe. Apply the first application when shoots are 6-8 inches long. A second application should be applied just before blooming. Foliar injury occurs on copper sensitive varieties.
Mango	Anthracoese	<i>Glomerella cingulata</i>	Dilute: 2 lbs/100 gal Concentrate: 4-6 lbs/A	Apply monthly after fruit set until harvest.
Olive	Peacock spot Anthracoese	<i>Cyrtospora oleaginum</i> <i>Glomerella oleivora</i>	Dilute: 2-3 lbs/100 gal Concentrate: 8-12 lbs/A	Apply prior to winter rains. A second application in the early spring should be made under severe disease symptoms.
Peaches and Nectarines	Brown rot blossom blight Coryneum blight (shot hole) Leaf curl Bacterial spot	<i>Monilinia</i> spp. <i>Coryneum beijerinckii</i> <i>Taphrina deformans</i> <i>Xanthomonas pruni</i>	Dilute: 2-4 lbs/100 gal Concentrate: 8-20 lbs/A Aerial: 8-20 lbs/A 2 lbs/100 gal 1/4 lbs/100 gal	For leaf curl and shot hole apply as a dormant spray from Nov 15 to Jan 15. For brown rot and shot hole apply before bud swell in the full pink bud stage. Apply as a dormant spray Apply post bloom at first and second cover sprays. Slight injury to the foliage may occur with post bloom sprays.
Pears	Pink blight	<i>Erwinia amylovora</i>	Dilute: 1/4-1/2 lbs/100 gal Concentrate: 1-2 lbs/A	Apply at 10% bloom and repeat at 5-7 day intervals during the bloom period. Do not apply to copper sensitive varieties.
Plums and Prunes	Brown rot blossom blight Coryneum blight (shot hole)	<i>Monilinia</i> spp. <i>Coryneum beijerinckii</i>	Dilute: 4-5 lbs/100 gal Concentrate: 12-20 lbs/A Aerial: 16-20 lbs/A	Apply as a dormant spray before heavy rains begin. For brown rot apply at early green bud to full popcorn stages.
Strawberries	Leaf spot Bacterial leaf blight	<i>Mycosphaerella fragariae</i> <i>Pseudomonas</i> spp.	2-3 lbs/100 gal 2-3 lbs/A	Begin application when plants are established and continue on a weekly schedule.
Tea	Blighter blight	<i>Exobasidium vexans</i>	2-4 lbs/A	Apply on a 7-10 day schedule.
Walnuts	Walnut blight	<i>Xanthomonas juglandis</i>	Dilute: 2-4 lbs/100 gal Concentrate: 8-12 lbs/A Aerial: 8-12 lbs/A	Make first application at early pre-bloom when the catkins are partially expanded. Make additional applications at 7-10 day intervals.

FIELD AND VEGETABLE CROPS

CROP	DISEASE	LATIN NAME	RATE	REMARKS
Alfalfa	Leaf spots	Cercospora spp. Leptosphaeria spp.	2-4 lb/A	Apply by ground or air 10-14 days prior to harvest. Slight injury may occur to sensitive varieties.
Beans	Bacterial blight (halo blight) Anthracnose Downy mildew Cercospora leaf spot	Xanthomonas phaseoli Pseudomonas phaseolicola Colletotrichum dematophanum Peronospora spp. Cercospora spp.	1-4 lb/A	Begin applications by ground or air when plants are 5-8 inches high and repeat at 10-14 day intervals.
Beets	Downy mildew Leaf spot	Peronospora spp. Cercospora beticola	1-3 lb/A	Apply by ground or air when disease first appears and repeat at 7-14 day intervals.
Broccoli, Brussel Sprouts Cabbage Cauliflower	Downy mildew Black rot Black leaf spot	Peronospora brassicae Xanthomonas campestris Alternaria spp.	1-2 lb/A 1-3 lb/A	Apply at 7 day intervals. Apply by ground or air when disease first appears and repeat at 7-14 day intervals.
Carrots	Downy mildew Blight	Peronospora spp. Cercospora spp. Alternaria spp.	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals.
Celery	Bacterial blight Early blight Late blight	Pseudomonas spp. Cercospora spp. Septoria spp.	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals.
Cucurbits (cantaloupes, honey dews, cucumbers, pumpkins, squash, watermelon)	Downy mildew Angular leaf spot Anthracnose Alternaria leaf spot Bacterial wilt Gummy stem blight	Pseudoperonospora cubensis Pseudomonas lachrymans Colletotrichum lagenarium Alternaria cucumerina Erwinia tracheiphila Mycosphaerella melonis	1W-1 lb/A	Apply by ground or air and when disease appears and repeat at 7-10 day intervals.
Eggplant	Alternaria blight Anthracnose Fruit rot	Alternaria spp. Colletotrichum spp. Phomopsis vexans	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals.
Lettuce	Downy mildew	Bremia lactucae	1-2 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals. Slight injury may occur under adverse weather conditions.
Onions	Downy mildew Purple blotch	Peronospora spp. Alternaria porri	2-4 lb/A	Apply by ground or air when plants are 4-6 inches high and repeat at 7-10 day intervals.
Peanuts	Cercospora leaf spot	Cercospora personata	1W-3 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals. May be tank-mixed with sulphur to enhance the control.
Peppers	Bacterial spot Anthracnose Early blight Late blight Leaf spot Downy mildew	Xanthomonas vesicatoria Colletotrichum spp. Alternaria solani Phytophthora infestans Cercospora capsici Peronospora spp.	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals.
Potatoes	Early blight	Alternaria solani	1-6 lb/A	Apply at 7-10 day intervals when plants are 8 inches high up until harvest. Apply by ground or air.
Spinach	Anthracnose Cercospora leaf spot Downy mildew White rust	Glomerella spp. Cercospora spp. Peronospora effusa A. byrrhoidentia s	2-3 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals.
Sugar Beets	Cercospora leaf spot	Cercospora personata	1W-1 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals. May be tank-mixed with sulphur to enhance the control.
Tobacco	Damping off diseases Angular leaf spot Frog's eye disease Brown spot Wild fire Blue mold Bacterial spot	Various Pseudomonas angularis Cercospora nicotiana Alternaria longipes Pseudomonas tabacoi Peronospora tabacina Xanthomonas vesicatoria	4-5 lb/A 4-5 lb/A 4-5 lb/A 2 lb/A 2 lb/A 2 lb/A	Avoid overwatering. Dust the seed bed with fungicide after planting. Apply on a 7-10 day basis when disease appears. Destroy all infected plants. Apply just before transplanting and when topped. Apply every 7-10 days when disease appears. Apply every 7 days from seeding to transplanting. Apply every 7-10 days when disease appears. Apply prior to appearance of the disease by ground or air and

Brussels Sprouts Cabbage Cauliflower	Downy mildew	Peronospora parasitica	1/2-1 lb/A	Apply at 7-day intervals
	Black rot Black leaf spot	Xanthomonas campestris Alternaria spp.	1-3 lb/A	Apply by ground or air when disease first appears and repeat at 7-14 day intervals.
Carrots	Downy mildew Blight	Peronospora spp. Cercospora spp. Alternaria spp.	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals
Celery	Bacterial blight Early blight Late blight	Pseudomonas spp. Cercospora spp. Septoria spp.	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals.
Cucurbits (cantaloupes, honey dews, cucumbers, pumpkins, squash, watermelon)	Downy mildew	Pseudoperonospora cubensis	1 1/2-1 lb/A	Apply by ground or air and when disease appears and repeat at 7-10 day intervals
	Angular leaf spot	Pseudomonas lachrymans		
	Anthrachnose	Colletotrichum lagenarium		
	Alternaria leaf spot	Alternaria cucumerina		
	Bacterial wilt Gummy stem blight	Erwinia tracheiphila Mycosphaerella melonis		
Eggplant	Alternaria blight Anthrachnose Fruit rot	Alternaria spp. Colletotrichum spp. Phomopsis vexans	2-4 lb/A	Apply by ground or air, when disease appears and repeat at 7-10 day intervals.
Lettuce	Downy mildew	Bremia lactucae	1-2 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals. Slight injury may occur under adverse weather conditions.
Onions	Downy mildew Purple blotch	Peronospora spp. Alternaria porri	2-4 lb/A	Apply by ground or air when plants are 4-6 inches high and repeat at 7-10 day intervals
Peanuts	Cercospora leaf spot	Cercospora personata	1 1/2-3 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals. May be tank-mixed with sulphur to enhance the control.
Peppers	Bacterial spot	Xanthomonas vesicatoria	2-4 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals
	Anthrachnose	Colletotrichum spp.		
	Early blight	Alternaria solani		
	Late blight	Phytophthora infestans		
	Leaf spot Downy mildew	Cercospora capsici Peronospora spp.		
Potatoes	Early blight	Alternaria solani	1-6 lb/A	Apply at 7-10 day intervals when plants are 6 inches high up until harvest. Apply by ground or air.
Spinach	Anthrachnose Cercospora leaf spot Downy mildew White rust	Glomerella spp. Cercospora spp. Peronospora effusa Aibyon occidentalis	2-3 lb/A	Apply by ground or air when disease appears and repeat at 7-10 day intervals.
Sugar Beets	Cercospora leaf spot	Cercospora personata	1 1/2-3 lb/A	Apply by ground or air when disease appears and repeat at 7-14 day intervals. May be tank-mixed with sulphur to enhance the control.
Tobacco	Damping off diseases	Various		Avoid overwatering. Dust the seed bed with fungicide after planting.
	Angular leaf spot	Pseudomonas angularis	4-5 lb/A	Apply on a 7-10 day basis when disease appears. Destroy all infected plants.
	Frog-eye disease	Cercospora nicotiana	4-5 lb/A	Apply just before transplanting and when topped.
	Brown spot	Alternaria longipes	4-5 lb/A	Apply every 7-10 days when disease appears
	Wild fire	Pseudomonas tabaci	2 lb/A	Apply every 7 days from seeding to transplanting.
	Blue mold	Peronospora tabacina	2 lb/A	Apply every 7-10 days when disease appears
Tomatoes	Bacterial spot	Xanthomonas vesicatoria	2-4 lb/A	Apply prior to the appearance of the disease by ground or air and repeat at 7-10 day intervals.
	Bacterial speck	Pseudomonas tomato		
	Bacterial canker	Corynebacterium michiganense		
	Anthrachnose	Glomerella phomoides		
	Early blight	Alternaria solani		
	Late blight	Phytophthora infestans		
	Leaf mold	Cladosporium fulvum		
	Septoria leaf spot	Septoria lycopersici		
	Gray leaf spot	Stemphylium solani		
Wheat and Barley	Helminthosporium spot blotch Septoria leaf blotch	Helminthosporium spp. Septoria spp.	1 1/2-2 lb/A	Apply by air or ground. Apply the first application at early heading and follow 10 days later with a second application.

MONERY CHEMICAL