

EFFECTIVE, FAST ACTING INSECTICIDE FOR USE IN HOMES, INSTITUTIONS, RESTAURANTS, FOOD PROCESSING PLANTS, DAIRIES, LIVESTOCK AND OUTDOORS FOR MOSQUITO CONTROL. APPLY WITH CONVENTIONAL MECHANICAL OR COMPRESSED AIR EQUIPMENT.

### PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

DO NOT apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. DO NOT APPLY WHILE FOOD PROCESSING IS UNDER WAY. After space spraying in bakeries, and other food processing plants, wash all equipment with potable water, benches, shelving etc., where exposed food will be handled. ALL food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, and cover fish aquariums before spraying.

### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish. DO NOT APPLY DIRECTLY TO WATER. DO NOT contaminate water supplies by cleaning of equipment or disposal of wastes.

### STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label directions must be disposed of according to applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures. BUYER assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Insecticide, Liquid, n.o.s

# swat #30

### Ready-To-Use Water-Based Insecticide

#### **ACTIVE INGREDIENTS:**

Pyrethrins 0.1%
Piperonyl Butoxide, Technical 1.0%
INERT INGREDIENTS (Contains Petroleum Distillate) 98.9%

TOTAL 100.0%

 Equivalent to 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds.

EPA Reg. No. 48211-49

EPA Est. No. 48211-MO-01

## KEEP OUT OF REACH OF CHILDREN CAUTION

Harmful If swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, IMMEDIATELY flush eyes or skin with planty of water. Obtain medical attention if irritation persists.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

DO NOT TRANSPORT OR STORE UNDER 32°F.

MANUFACTURED BY:

INTERCON CHEMICAL COMPANY

3647 BELL AVENUE

ST. LOUIS, MISSOURI 63108

Not Reviewed. Registrant claims to be in aconfdance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. ALL food processing surfaces should be covered during treatment or thoroughly cleaned before using. Vacate treated area and ventilate before reoccupying. For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control Roaches, Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Meal Worms, Grain Mites, and Cadelles: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray Into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For SILVERFISH, spray bookcases. For ANTS spray trails, nests and points of entry. If surface application ONLY is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control BEDBUGS: Spray mattresses lightly, particularly tufts and seams. Take beds apart and spray in all joints. Repeat treatment as needed. To control FLEAS and BROWN DOG TICKS: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moulding, window and door trames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

**NET CONTENTS:** 

GAL.

2280

DIRE To control CLOTHI storage. Brush thoro ble. Apply spray libe storages, directing si fested articles may from fabric to avoid SPACE SPRAYING: systems. Apply with a For RAPID control of Wasps, Hornets. An-Indian Meal Moths ( angle, distributing It 6 ounces per 1000 cu FOGGING: To kill e crawling insects, use space. CONTACT IN ON ANIMALS: To pri Mosquitoes and Gna of the hair. To contro

tinued protection.
To control Blood-Su using a stiff brush to or 3 weeks IF require OUTDOOR ADULT I thoroughly treat the velocity DOES NOT tion where mosquite ponds and shora line per 1000 square feet Using truck-mount Rotomist® Models toot swath, calibrate

rate of 2 ounces o