

# stomp-out granular

## A GRANULAR NON-SELECTIVE WEED AND GRASS KILLER

Active Ingredient:	
Bromacil (5-bromo-3-sec-butyl-6-methyluracil) .....	1.5%
Inert Ingredients .....	98.5%
TOTAL	100.00%

EPA Reg. No. 48211-73

EPA Est. No. 48211-MO-01

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

Harmful if swallowed or inhaled. Avoid breathing dust. Remove contaminated clothing and wash before reuse. Do not get in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling. See label for additional precautionary statements.

MANUFACTURED BY:

INTERCON CHEMICAL COMPANY

ST. LOUIS

3647 BELLEVUE

ST. LOUIS

ST. LOUIS

MISSOURI 63108

ACCEPTED

MAR - 7 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 48211-73

### USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is specifically designed for quick knock-down and control of general mixed vegetation and such perennial grasses as Johnson Grass, Dallis Grass, and Bermuda Grass on noncropped land and industrial sites. This product may provide season-long control of seedlings on treated areas. Soil type, rainfall, and length of growing season will influence control. Apply on parking lots, storage yards, fence lines, highway and railroad sites, and other similar areas.

**APPLICATION:** For all uses, uniform application is essential. Applications become effective as soil treatments following rainfall or irrigation. Conventional herbicide spreaders may be used for application.

**GENERAL VEGETATION CONTROL:** For DRY granular treatment, apply 1/2 to 2 pounds per 100 sq. ft. The lighter rates should be used only at time of weed emergence on moist soil or when precipitation is anticipated while weeds are still young. The higher rates may be necessary to insure thorough coverage of heavy vegetation or to provide extended residual control.

**PERENNIAL WEED CONTROL:** Persistent perennial weeds are most effectively treated when nearing maturity or during dormant stages in the fall. Spring applications may also be made under moderate rainfall conditions. Summer applications are recommended only where rainfall is dependable for leaching herbicide into the soil. Apply 1 to 3 pounds per 100 sq. ft. On deep-rooted perennials such as Bindweed, Canada Thistle, Bermuda Grass, and Quack Grass the heavier rates may be required.

**JOHNSON GRASS CONTROL:** This product is particularly effective on deep-rooted perennial grasses such as Johnson Grass, Bermuda Grass and Dallis Grass on industrial sites, roadsides, fence lines and storage yards. Apply dry at 2 to 3 pounds per 100 sq. ft. for the control of established perennial grasses. Seedlings can be controlled with 1 to 2 pounds per 100 sq. ft. Maximum efficiency on Johnson Grass is obtained by treating after heading until late fall.

Net weight: 50 lb.

2348

**BEST AVAILABLE COPY**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
STATEMENT OF PRACTICAL TREATMENT  
CAUTION**

**IF SWALLOWED:** Call a physician or Poison Control Center immediately.  
**IF IN EYES:** Flush with plenty of water. See a physician if irritation persists.  
**IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

**STORAGE AND DISPOSAL**

**PESTICIDE DISPOSAL** - Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state, or local procedures.

**CONTAINER DISPOSAL** - Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner and drum in a sanitary landfill or by incineration if allowed by state and local authorities. If burned, stay away from smoke.

**WARRANTY STATEMENT**

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**NOTICE**

Because of the many varying conditions affecting use and application, such as location, seasonal weather, soil composition, and type of vegetation, manufacturer warns buyer that these and other conditions may impair or vary the results or effects of the use of this product. In any event, complete elimination of weeds is not guaranteed. It is advisable to consult your State Extension Service or Agricultural Experiment Station Weed Specialists on specific problems.

**APPLY DRY.**

**FOR GENERAL VEGETATION AND PERENNIAL GRASSES  
ON NONCROPPED LAND AND INDUSTRIAL SITES.**

**IMPORTANT**

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops, or other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlain by roots of adjacent valuable growths. Careless application of this material or washing by water runoff to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution. Use of low trajectory angle sprinklers or drop nozzles will decrease the likelihood of drift.

**BEST AVAILABLE COPY**