

SYMCOAT

DIRECTIONS FOR USE

GENERAL CLASSIFICATION

It is a violation of Federal Law to use this product in a manner inconsistent with the label.

INSTRUCTIONS FOR USE:

Application:

1. Shake Very Well!
2. Pre-planter box seed treater: Calibrate the equipment accurately to apply the recommended rate of SYMCOAT for the respective seeds being coated.
3. Planter box mixing: This can be achieved by pouring or spraying the seeds with SYMCOAT while filling the planter box and using a stick to thoroughly mix the seeds and seed coat. Fill the planter box in quarters and mix at each level.
4. Use with fungicides: If mixed with fungicides for ease of application check compatibility. Thoroughly mix the SYMCOAT with fungicides as recommended to assure even coating of both products on the seed. Read the fungicide label for application instructions and cautions.

AIDS PLANT GROWTH AND INCREASES YIELDS

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)..... 0.01

based on biological activity

Inert Ingredients.....99.99%

Keep Out Of Reach Of Children

CAUTION

NET VOLUME: 32 Fluid Oz.
(.95 Liters)

NET WEIGHT: 2.44 lbs.
(1.09 kg)

SEE ADDITIONAL PRECAUTIONARY STATEMENTS
ON SIDE PANEL

E.P.A. Reg. No.:

Establishment No.:

AGRO-K
CORPORATION
5750 Main Street NE
Minneapolis, Minnesota 55432
(612) 571-6555/Telex No. 290936
AGRO-K CORP ERLY

RATE OF APPLICATION.

Soybeans - 2 fluid ounces Symcoat
per 60 lbs. medium size seeds.
Wheat - 2 fluid ounces mixed with
2 ounces of water per 100 lbs of
seeds.
Corn - 2 fluid ounces Symcoat per
bushel of seed.

PRECAUTIONARY STATEMENTS

ENVIRONMENTAL HAZARDS

Do Not apply directly to water.
Do Not contaminate water by clean-
ing of equipment or disposal of
water.

CONTAINER DISPOSAL

Do not reuse empty container.
Wrap container and put in trash
collection.

