 <p align="center"><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b>  <b>Office of Pesticide Programs</b>  <b>Registration Division (7505P)</b>  <b>Ariel Rios Building</b>  <b>1200 Pennsylvania Ave., NW</b>  <b>Washington, D.C. 20460</b></p> <p align="center">NOTICE OF PESTICIDE:  <input type="checkbox"/> Registration  <input checked="" type="checkbox"/> Reregistration          (under FIFRA, as amended)</p>	EPA Reg. Number:	Date of Issuance:
	48234-2	25 NOV 2008
	Term of Issuance:	
Name of Pesticide Product:		<b>Regal Ronstar Plus</b>

Name and Address of Registrant (include ZIP Code):


**Regal Chemical Company**  
**P.O. Box 900**  
**Alpharetta, GA 30239**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:

- Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.
- Based on toxicity ranking per the acute toxicity review, the First Aid statements should be placed on the label in the following order:  
 "IF ON SKIN OR CLOTHING:...  
 IF INHALED:...  
 IF IN EYES:...  
 IF SWALLOWED:..."
- Per the acute toxicity review, the Hazards to Humans and Domestic Animals must be revised to read:  
 "CAUTION  
 Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation.  
 Avoid contact with skin, eyes, or clothing. Avoid breathing dust."

Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	 Date: 25 NOV 2008
---------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------

4. Based on acute toxicity data for this product and the RED, the PPE section currently on the label should be revised to read:

“Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Loaders, applicators, and other handlers must wear:  
Long-sleeved shirt and long pants,  
Shoes plus socks, and  
Chemical-resistant gloves.”

5. Per the Oxadiazon RED label table, the text in **bold type** below must be added to the following User Safety Requirements:

“...If no such instructions for washables **exist**, use detergent and hot water...”

6. The text in bold type must be added to the User Safety Recommendation text currently on the label:

“User should remove clothing/**PPE** immediately if pesticide gets inside.”

7. Per the RED, the early entry PPE currently on the label (“coveralls over shirt-sleeved shirt and short pants, chemical-resistant gloves made of any waterproof material, chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure”) must be revised to read:

“Coveralls,  
Shoes plus socks, and  
Chemical-resistant gloves made of any waterproof material.”

8. A Non-Agricultural Use Requirements box must be added to label with the following entry restriction text:

**“NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter the treated area until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.”

9. Per the product chemistry review, the statement "Do not contaminate water, food by storage or disposal" must be changed to read "Do not contaminate water, food, or feed by storage or disposal" and the statement must be placed directly under the Storage and Disposal heading. In addition, the subheading "Storage" should be changed to "Pesticide Storage."
10. Under the Storage and Disposal section, change first bullet to read "**Pesticide Storage**".
11. Under the Notice to Buyer, change the last statement to read "**To the extent consistent with applicable law**, seller makes no warranties; express or implied."

A stamped copy of your labeling is enclosed for your records. Please note that the final product reregistration cannot be considered until after all active ingredients in this product are eligible for reregistration. Submit one copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller  
Product Manager (23)  
Herbicide Branch  
Registration Division (7505P)

Enclosure

4/11

NOT FOR USE ON HOME LAWNS. FOR SALE FOR USE AND USE BY PROFESSIONAL APPLICATORS ONLY. NOT FOR SALE TO OR USE BY HOMEOWNERS/CONSUMERS.

**PREEMERGENT HERBICIDE FOR TURF AND ORNAMENTALS**

Selective, preemergent herbicide for the control of annual grasses and broadleaf weeds in turf and woody ornamental shrubs, vines and trees.

**ACCEPTED**  
with **COMMENTS**  
in EPA Letter Dated:  
**25 NOV 2008**

**REGAL RONSTAR PLUS**

Under the Federal Insecticide Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No

48234-2

**ACTIVE INGREDIENT:**

Oxadiazon [2-tert-butyl-4-(2, 4-dichloro-5-isopropoxyphenyl)-delta-1, 3, 4-oxadiazolin-5-one] ..... 2.0%

**OTHER INGREDIENTS:** ..... 98.0%

**TOTAL:** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN  
WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-621-5208 for emergency medical treatment information.	

EPA REG NO. 48234-2

EPA EST NO. 48234-GA-1

Regal Chemical Company  
PO Box 900  
Alpharetta, GA 30239

**PRECAUTIONARY STATEMENTS**

**WARNING**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Causes skin and eye irritation. Harmful if inhaled. Do not get on skin or on clothing. Avoid contact with eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Loaders, applicators, and other handlers must wear:

- Coveralls worn over short-sleeved shirt and short pants
- Chemical resistant gloves made of any waterproof material
- Chemical resistant footwear plus socks
- Protective eye wear
- Chemical resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticides get inside. Then wash body thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Regal Ronstar Plus® should not be used in areas where irrigation or rainfall results in direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxadiazon is absorbed.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance to its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow entry into treated areas during the restricted entry interval (REI) 12 hours.

PPE required for early entry to treated areas, that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is coveralls over short-sleeved shirt and short pants, chemical resistant gloves made of any waterproof material, chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure.

**STORAGE AND DISPOSAL**  
 Storage: Do not contaminate water, food by storage or disposal. Do not store in or around the home. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Completely empty bag into application equipment. Then dispose of bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**GENERAL INFORMATION**  
 Regal Ronstar Plus is a selective, preemergent herbicide for the control of many annual grasses and broadleaf weeds in turf and on a wide variety of woody ornamental shrubs, vines and trees. Regal Ronstar Plus controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination. Apply Regal Ronstar Plus with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly. Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

- Restrictions:**
- Not for use on home lawns
  - Do not apply this product through any type of irrigation system.
  - Do not apply aerially.
  - Do not apply more than 300 pounds of product per acre per year (equivalent to 6 pounds ai/A/year) except in areas of heavy weed infestation.
  - In areas of heavy weed infestation, do not apply more than 375 pounds of product per year (equivalent to 7.5 pounds ai/A/year).
  - Do not apply more than 200 pounds of product per acre (equivalent to 4 pounds ai/A) in a single application.

**TURF**  
 For use in established perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue and Zoysia turf; such as, fairways, parks, golf courses and lawns. Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. Do not apply to wet turf. For best results, water-in the product as soon as practical after application.

**PREEMERGENT WEED CONTROL**

Weeds Controlled	Amount of Regal Ronstar Plus		Directions
	Per acre	Per 1000 sq. ft.	
Crabgrass ( <i>Digitaria spp.</i> ) Goosegrass ( <i>Eleusine indica</i> ) Field Sandbur ( <i>Cenchrus incertus</i> ) Annual Sedge ( <i>Cyperus compressus</i> )	100-200 lbs.	4 1/4 - 4 1/2 lbs.	Apply late winter or early spring prior to seed germination. Application made after March or April may not be fully effective.
Annual Bluegrass ( <i>Poa Annua spp.</i> ) (See note below)	100-200 lbs.	2 1/4 - 4 1/2 lbs.	Apply in late summer to early fall prior to seed germination.
Virginia Buttonweed ( <i>Diodia virginiana</i> ) Pigweed ( <i>Amaranthus spp.</i> ) Florida Pusley ( <i>Richardia scabra</i> ) Oxalis ( <i>Oxalis spp.</i> ) Stinging Nettle ( <i>Urtica dioica</i> ) Carpetgrass ( <i>Axonopus offinis</i> )	200 lbs.	4 1/2 lbs.	Apply in early spring prior to seed germination.

NOTE: In many areas, annual bluegrass exists as at least two biotypes or subspecies of *Poa annua*. The true annual biotype is *Poa annua* spp. annua which germinates every year from seed, while the biotype that behaves like a perennial is *Poa annua* spp. reptans. This perennial biotype

can survive through the summer and established plants will not be controlled by this product. Consult the extension service or University weed specialists in your area for more information.

**FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF**

Weeds Controlled	Amount of Regal Ronstar Plus		Directions
	Per acre	Per 1000 sq. ft	
Crabgrass, Goosegrass, and other annual weeds	150-200 lbs.	3 3/8 – 4 1/2 lbs.	For use in Bluegrass, Bentgrass, Perennial Ryegrass and Tall Fescue turf. Make application in the fall after turf has become dormant. For late, fall applications, seeding into treated areas should be delayed until 4 months after green up.

**ANNUAL BLUEGRASS CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED WITH PERENNIAL RYEGRASS ONLY**

Weeds Controlled	Amount of Regal Ronstar Plus		Directions
	Per acre	Per 1000 sq. ft	
Annual Bluegrass ( <i>Poa Annua</i> spp.) and other annual weeds	100 lbs.	2 1/4 lbs	Apply in late summer to early fall prior to <i>Poa annua</i> spp. seed germination. Any <i>Poa annua</i> spp. established when this product is applied will not be controlled. Regal Ronstar Plus will only control the <i>Poa annua</i> spp. that is established from seed every year. Regal Ronstar Plus should be applied no closer than 60 days prior to overseeding. NOTE: Do not use where <i>Poa trivialis</i> is present in overseed mix.

NOTE: In many areas, annual bluegrass exists as at least two biotypes or subspecies of *Poa annua*. The true annual biotype is *Poa annua* spp. *annua* which germinates every year from seed, while the biotype that behaves like a perennial is *Poa annua* spp. *reptans*. This perennial biotype can survive through the summer and established plants will not be controlled by this product. Consult the extension service or University weed specialists in your area for more information.

**ANNUAL WEED CONTROL DURING BERMUDAGRASS, ZOYSIAGRASS, SEASHORE PASPALUM AND TROPIC LALO (*PASPALUM HIERONYMII*) ESTABLISHMENT FROM SPRIGS**

Weeds Controlled	Amount of Regal Ronstar Plus		Directions
	Per acre	Per 1000 sq. ft	
Crabgrass, Goosegrass, and other annual weeds	100-150 lbs.	2 1/4 - 3 3/8 lbs.	For Bermudagrass, make application immediately prior to or immediately after sprigging. For Seashore Paspalum, make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after Regal Ronstar Plus application may disrupt the herbicide barrier and reduce effectiveness. Irrigate immediately after application for best results.

For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 150 pounds per acre (3 pounds active ingredient) of Regal Ronstar Plus.

Do not apply to Bentgrass turf mowed at less than 3/8 inch.

Regal Ronstar Plus may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.

Avoid contact with flowers and shrubs except as recommended elsewhere on this label. Do not apply Regal Ronstar Plus to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment except as noted above. Do not apply to Dichondra or Centipedegrass. Regal Ronstar Plus should not be used on putting greens or tees because of the varying cultural practices used.

When making applications to Tropic Lalo, applications must be made 0-12 days after sprigging. Do not use in areas where endangered Hawaiian Goose or nene (*Branta sandvicensis*), Hawaiian Duck (*Anas wyvilliana*), Hawaiian Stilt (*Himantopus mexicanus knudseni*), and Hawaiian Moorhen (*Gallinula chloropus sandvicensis*) may feed.

**WOODY ORNAMENTAL SHRUBS, VINES, TREES AND CONIFER NURSERY APPLICATION**

It is impossible to test every species and variety of ornamental or nursery plants. Plant tolerance to Regal Ronstar Plus has been found to be acceptable in the specified genera and species. Neither the manufacturer nor the seller has determined whether or not Regal Ronstar Plus can safely be used on plants not specified on this label. The professional user should determine if Regal Ronstar Plus can be used safely prior to use.

Apply Regal Ronstar Plus anytime during the year, prior to weed seed germination, for weed control. Regal Ronstar Plus can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of Regal Ronstar Plus will improve weed control activity.



Regal Ronstar Plus should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of Regal Ronstar Plus. Do not apply to wet foliage or under conditions in which granules will collect on leaves. Do not apply in greenhouses as plant injury may result from co-distillation of Regal Ronstar Plus active ingredient onto leaf surfaces in condensation.

**Amount To Use**

Ground Application — Apply Regal Ronstar Plus broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 1/4 to 4 1/2 pounds per 1,000 sq. ft.) depending on the weeds to be controlled, as listed on the following page. Remove existing weed growth before application.

**ANNUAL BROADLEAF WEEDS**

Bittercress	Bristly Oxtongue	Carpetweed
Cheeseweed	Common Groundsel	Common Purslane
*Evening Primrose	Fiddleneck	Fireweed
*Galinsoga	Garden Spurge	Golden Ragwort
Lambsquarters	** Liverwort	Niruri
Pennsylvania Smartweed	Petty Spurge	**Prostrate Spurge
Redroot Pigweed	Shepherdspurge	Sensitive Plant
Sow Thistle	Speedwell	Spiny Amaranth
Spotted Catsear	*Swinecress	Yellow Woodsorrel (Oxalis)

\*Use 150 to 200 pounds (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

\*\*Use 200 pounds (4.0 pounds of active ingredient) per acre for less susceptible weeds.

**ANNUAL GRASSES**

Annual Bluegrass	Annual Sedge	*Barnyardgrass
Bristly Foxtail	Crabgrass	Fall Panicum
Green Foxtail	Goosegrass	**Jungle Ricegrass
**Panama Paspalum	Ripgut Bromegrass	Wild Oats

\*Use 150 to 200 pounds (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

\*\*Use 200 pounds (4.0 pounds of active ingredient) per acre for less susceptible weeds.

**CROP TOLERANCE**

Regal Ronstar Plus is recommended for use on a wide variety of ornamental species and ground covers in container, field grown and landscape planting situations. Regal Ronstar Plus may be used at the recommended rates on the following species.

**ESTABLISHED ORNAMENTALS AND GROUND COVERS**

Abutilon	Acacia	Agapanthus
Allamanda, Common	Andromeda	Aralia
Arborvitae	Ash, Green	Azalea
Baby's Breath	Bald Cypress	Bamboo, Feathery
Banyan, Weeping	Barberry	Beafsteak plant
Bearberry	Beech, European	Birch, Paper
Blanket, Flower	Blazing, Star	Boxwood, Japanese
Boxwood, Common	Bougainvillea	Brassaia, DwarfBuddleia
Bugleflower	Callistemon	Camellia, Sasanqua
Candytuft	Carissa	Carnation
Ceanothus	Cedar, Red	Cedar, Atlas
Cistus	Citrus, Ornamental*	Grapefruit
Lemon	Lime	Mandarin

Tangelo  
 Cuban Shaddock  
 Cleyera  
 Cotoneaster  
 Chrysanthemum  
 Dahlia  
 Dieffenbachia  
 Dogwood, Kousa  
 Escallonia  
 Euonymus, Spreading  
 Falsecypress, Hinoki  
 Forsythia  
 Ginko  
 Heath, Spring  
 Hemlock, Canada  
 Holly  
 Ivy, Algerian  
 Jasmine, Primrose  
 Juniper  
 Lamb's Ear  
 Larch, European  
 Leucothoe, Coast  
 Lilac  
 Magnolia  
 Metallic Plant  
 Money Tree  
 Myrtle  
 Ochana  
 Orthosiphon  
 Palm, Bamboo  
 Palm, Macarthur  
 Palm, Parlor  
 Palm, Sugar  
 Periwinkle, Common  
 Pine  
 Podocarpus, Yew  
 Potentilla  
 Pyracantha  
 Rhododendron  
 Sargent Cherry  
 Serviceberry, Saskatoon\*  
 St. Johnswort, Golden  
 Strawberry Tree  
 Syzygium  
 Wedelia  
 Woadwaxen, Common  
 Yew

Tangerine  
 Valencia  
 Coreopsis  
 Crabapple\*  
 Cypress  
 Daphne, Fragrant  
 Dogwood, Common  
 Douglas fir  
 Eucalyptus  
 Euonymus,  
 Falsecypress, Japanese  
 Gardenia  
 Gladiolas  
 Heather, Scotch  
 Hibiscus  
 Ice Plant  
 Ivy, English  
 Jasmine, Star  
 Justicia  
 Lantana, Trailing  
 Laurel, English  
 Leucothoe, Drooping  
 Littleleaf Linden  
 Mahonia, Grape Holly  
 Mexican Orange  
 Murraea  
 Nandina  
 Ohai  
 Osmanthus  
 Palm, Fishtail  
 Palm, Manac  
 Palm, Rattan  
 Paperbark Tree  
 Photinia, Frasier's  
 Pittosporum, Japanese  
 Pomegranite\*  
 Privet  
 Raphiolepis  
 Rose  
 Scotch Broom  
 Spruce  
 St. Johnswort,  
 Sumac, Staghorn  
 Trailing African Daisy  
 Wisteria, Chinese  
 Woolly Thyme

Temple  
 Washington Navel  
 Coromandel  
 Crapemyrtle  
 Cypress, Leyland  
 Deutzia, Slender  
 Dogwood, Redosier  
 Elm  
 Euryops, Gray Leaved  
 Wintercreeper  
 Flowering Quince  
 Gazania, Trailing  
 Golden Raintree  
 Heliotropium  
 Honeysuckle  
 Ilima  
 Jasmine, Orange  
 Jasmine, Showy  
 Kwanzan Cherry  
 Lapalapa  
 Leopards - Bane  
 Lily of the Nile  
 Loquat\*  
 Maple  
 Mock Orange  
 Myoporum  
 Oak  
 Oleander  
 Pachysandra  
 Palm, Licuala  
 Palm, Manila  
 Palm, Sealing Wax  
 Peony  
 Poinciana  
 Plum, Natal  
 Poplar  
 Protea species  
 Redwood, Dawn  
 Russian Olive  
 Sedum  
 Statice  
 Gokdencup  
 Sweet William  
 Viburnum  
 Witchazel, Common  
 Xylosma

**NEWLY TRANSPLANTED ORNAMENTALS AND GROUND COVERS**

Blanket Flower  
 Carnation  
 Dogwood, Flowering  
 Gazania, Trailing  
 Ivy, English  
 Ice Plant

Brassaia, Dwarf  
 Cedar, Red  
 Elm  
 Hackberry  
 Hemlock, Canada  
 Juniper

Camellia  
 Coromandel  
 Forsythia  
 Ivy, Algerian  
 Holly  
 Leopards - Bane

Metallic Plant  
Periwinkle, Large  
Privet  
Trailing African Daisy  
Yew

Oak  
Pine  
Protea Species  
Wedelia

Periwinkle, Common  
Poplar  
Spruce  
Wisteria, Chinese

\*For use only on plants that will not bear edible fruit within one year of Regal Ronstar Plus herbicide application.

**FOREST NURSERY APPLICATION**

Seedlings of the following Species can be treated with Regal Ronstar Plus herbicide when at least 4 weeks old: Pine – Loblolly, Eastern White, Slash

**SENSITIVE VARIETIES**

Under certain growing conditions, the plant varieties listed below may be sensitive to Regal Ronstar Plus herbicide. Growers should evaluate usage on a small scale prior to large scale applications.

**PLANT SPECIES**

Plant Species	Varieties
Andromeda	Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, (Rhododendron spp), Leuchtenfuer, Memoire John Hearrens, Hino- Crimson, White Grandeur, Everest, Girard Roberta
Cotoneaster	Eichholz, Himalayan, Shangri-la (Cotoneaster spp)
Crapemyrtle	Petite Red Imp, Seminole Bush
Heath, Spring	Mediterranean Pink
Hemlock, Canada	Pendula
Hibiscus	Red Heart
Rhododendron	Daphnoides, Nova Zembla, Rhododendron spp), Catawba Album
Spruce, Norway	Pumila
Spruce, White	Densata
Viburnum	Davidii

**NOTICE TO BUYER**

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal conditions. This warranty does not extend to use of this product contrary to label use directions, or under conditions not reasonably foreseeable to seller; buyer assumes all risk of any such use. Seller makes no warranties; expressed or implied.

Ronstar® is a registered trademark of the Bayer CropScience.