



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

January 22, 2026

William Dunnivant
General Manager
Regal Chemical Company
600 Branch Dr.
Alpharetta, GA 30004

Subject: Label Amendment - Registration Review Mitigation for Prodiamine
Product Name: RegalKade G
EPA Registration Number: 48234-16
Case Number: 675053
Application Date: January 15, 2026

Dear William Dunnivant:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Prodiamine Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must submit one copy of the final printed labeling before you release the product for

shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Tracy Jackson by phone at 202-566-2268, or via email at jackson.tracy@epa.gov.

Sincerely,



Maryam K. Muhammad-Perch, Team Lead
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

ENCLOSURE: Stamped label

ACCEPTED**01/22/2026**

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under
EPA Reg. No. 48234-16

PRODIAMINE

GROUP

3

HERBICIDE

RegalKade® G

For pre-emergence control of grass and broadleaf weeds in:

- Container, Field Grown and Landscape Ornamentals
- Conifer and Hardwood Seedling Nurseries
- Established Perennial & Wildflower Plantings
- Plants Grown for Foliage Production (Florida Only)
- Christmas Tree Farms

ACTIVE INGREDIENT:Prodiameine [2,4-dinitro-*N*³,*N*³-dipropyl-6-(trifluoromethyl)-1,3-benzenediamine] 0.5%**OTHER INGREDIENTS:** 99.5%**TOTAL:** 100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-621-5208 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**EPA REG. NO. 48234-16
EPA EST. NO. 48234-GA-1**

Manufactured by:
REGAL CHEMICAL COMPANY
ALPHARETTA, GEORGIA

NET WEIGHT:

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4.6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following directions to minimize off site movement.

AQUATIC ORGANISM ADVISORY: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, whether directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling, and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under

certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

GENERAL INFORMATION

RegalKade G is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Container, Field Grown Ornamentals, Landscapes
- Conifer and Hardwood Seedling Nurseries
- Established Perennial & Wildflower Plantings
- Plants Grown for Cut Foliage Production
- Christmas Tree Farms

RegalKade G controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when RegalKade G is activated by at least 0.5 in. of rainfall or irrigation, or with shallow (1 or 2 in.) incorporation, prior to weed seed germination and within 14 days following application.

Do not graze or feed livestock forage cut from areas treated with RegalKade G.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

WEED RESISTANCE MANAGEMENT: For resistance management, RegalKade G is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to RegalKade G and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of RegalKade G or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and considers mechanical control methods), cultural, biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or pest control advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific plants and weed biotypes.

Contact your local sales representative or extension agent to find out if suspected resistant weeds to this product have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specific for your local conditions.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

LANDSCAPES, CONTAINER, FIELD GROWN & LANDSCAPE ORNAMENTALS (including Christmas Trees, Established Perennial and Wildflower Plantings):

RegalKade G may be applied for residual pre-emergence weed control in the culture of ornamentals.

USE RATES

Apply RegalKade G at 132 to 300 lb./Acre in Fall and/or Spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 300 lb./Acre. All applications must be made prior to germination of the weed seeds.

WEEDS CONTROLLED

When used in Accordance with this label RegalKade G will provide control of the following weeds:

Barnyardgrass	Kochia
Bluegrass, Annual (<i>Poa annua</i>)	Lambsquarter, Common
Carpetweed	Lovegrass
Chickweed, Common	Panicum (Texas, Fall, Browntop)
Chickweed, Mouseear (from seed)	Pigweed
Crabgrass (large, smooth)	Purslane, Common
Crowfootgrass	Pursley, Florida
Cupgrass, Woolly	Rescuegrass
Foxtail, Annual	Shepherd's Purse
Goosegrass	Signalgrass, Broadleaf
Henbit	Speedwell, Persian
Itchgrass	Sprangletop
Johnsongrass (from seed)	Spurge, Prostrate
Junglerice	Witchgrass
Knotweed	Woodsorrel, Yellow (from seed)

APPLICATION TIMING AND INFORMATION

RegalKade G may be applied to newly transplanted and established container, field nursery stock and landscapes as a broadcast, over the top application. Irrigation or rainfall soon after application will wash granules off plant foliage and activate RegalKade G in the soil. Apply RegalKade G to dry foliage only.

RegalKade G may be applied at any time to established plants. Delay application to allow soil to settle around new transplants and water thoroughly before applying RegalKade G. Apply RegalKade G after cuttings from roots and become established. Apply before budding and grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. In addition to outdoor locations, RegalKade G may be applied in shadehouses and uncovered polyhouses. Uncovered polyhouses must remain open for a minimum of 30 days following treatment. RegalKade G may be applied to soil (including mulch, gravel, wood chips, or other permeable base) upon which containerized ornamentals are placed. Only shallow cultivation or hand weeding, if necessary after application, should be performed so as to minimize exposure of untreated soil.

RegalKade G is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when RegalKade G is activated by at least 0.5 in. of rainfall or irrigation, or with shallow (1 or 2 in.) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when RegalKade G is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of RegalKade G, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

CONIFER AND HARDWOOD SEEDLING NURSERIES

RegalKade G may be applied for residual pre-emergence weed control in conifer and hardwood seedling nurseries. RegalKade G may be applied to conifer and hardwood seedling nurseries any time after the soil has settled around newly transplanted seedlings, liners or bare root plants. Apply RegalKade G at 132 to 300 lb./Acre in Fall and/or Spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 300 lbs. of product/Acre. Most effective weed control will be obtained when RegalKade G is activated by at least 0.5 in. of rainfall or irrigation prior to weed seed germination and within 14 days following application.

TOLERANT ORNAMENTAL SPECIES

RegalKade G will not harm most ornamental trees, shrubs and flowers. The species listed below are tolerant to RegalKade G. Best results will be obtained when RegalKade G is uniformly applied to the soil surface. RegalKade G may be applied over the top of these species without injury.

Scientific Name	Common Name
<i>Abies spp.</i>	Fir Species** (balsam, Fraser, Nobel, etc.)
<i>Abelia grandiflora</i>	Abelia: Sherwood
<i>Acer palmatum</i>	Japanese Maple
<i>A. platanoides</i>	Norway Maple
<i>Aciliiea sp.</i>	Yarrow: King Edward
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Agapanthus Orientalis</i>	
<i>Akebia quintata</i>	Five-Leaf or Chocolate Vine
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion
<i>Anemone hybrida</i>	Japanese Anemone
<i>Aquilegia sp.</i>	Aquilegia: Red and Gold
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Artemesia sp.</i>	Wormwood; Silver Mound, Castle
<i>Aster sp.</i>	Aster: Bonny Blue, Purple Dome
<i>Aster X Frikartii</i>	
<i>Athrium Filiz-femina</i>	Lady Fern; Fern Lady
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Begonia sp.</i>	Fibrous Begonia: Hardy Grandis
<i>Berberis gladwynensis</i>	Barberry
<i>B. Julianae</i>	Wintergreen Barberry
<i>B. mentorensis</i>	Mentor Barberry
<i>B. Thunbergii</i>	Japanese Barberry
<i>B. verruculosa</i>	Warty Barberry
<i>Bergenia cordifolia</i>	
<i>Boltonia asteroides</i>	Snowbank
<i>Bougainvillea sp.</i>	Bougainvillea
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue); Royal Red
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)
<i>Campus X Tagliabuana</i>	Trumpet Creeper, Trumpet Flower, Madame Galen
<i>Carpobrotus edulis</i>	Hottentrot Fig (Ice Plant)
<i>Cassia artemisoides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac

Scientific Name	Common Name
<i>Ceratostigma plumbaginoides</i>	
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Chrysanthemum nipponicum</i>	
<i>Cleyera japonica</i>	Cleyera
<i>Citrus spp.</i>	Citrus species*
<i>Coreopsis sp.</i>	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
<i>Cornus florida</i>	Flowering Dogwood
<i>C. stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>C. buxifolius</i>	Cotoneaster
<i>C. Dammeri</i>	Bearberry Cotoneaster
<i>C. microphyllus</i>	Rockspray Cotoneaster
<i>Cretaegus spp.</i>	Hawthorne
<i>Crocosmia sp.</i>	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma sp.</i>	Cooperi Pink
<i>Delphinium sp.</i>	Larkspur: Blue Elf
<i>Dianthus deltoides</i>	Dianthus; Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea purpurea</i>	Coneflower, Purple; Mangus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus Fortunei</i>	Wintercreeper
<i>E. japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia suspensa</i>	Weeping Forsythia
<i>F. viridissima</i>	Greenstem Forsythia
<i>Gaillardia sp.</i>	Gaillardia, Blanket Flower: "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine, Radicans
<i>Gaura sp.</i>	
<i>Gentianna dahurica</i>	Gentian
<i>Geranium Cinereum</i>	Cranesbill
<i>Gladiolus spp.</i>	Gladiolus species**
<i>Gypsophila repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthemum sp.</i>	Sunrose
<i>Hemerocallis sp.</i>	Daylily: Aztec Gold, Stella De Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus sp.</i>	Mallow: Disco Belle White
<i>Hibiscus Rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. Sieboldiana</i>	Hosta, "Searsucker"
<i>Houttuynia cordata var. variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Ilex Bernyi</i>	Holly
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly: Helleri

Scientific Name	Common Name
<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly: Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris spp.</i>	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsoni
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans sp.</i>	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica X Faurei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender sp.</i>	Lavender; Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>Ligustrum amurense</i>	Amur Privet
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lillium sp.</i>	Lily: Jazz
<i>Liriope Muscari</i>	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
<i>Liriope Muscari var. variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower, Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum sp.</i>	Loosestrife; Modern Pink
<i>Magnolia spp.</i>	Magnolia species**
<i>Maleophora luteola</i>	Ice Plant
<i>Malus sp.</i>	Crabapple*
<i>Miscanthus sinensis</i>	Yaku Jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus spp.</i>	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly): Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photinia Fraseri</i>	Frasier's Photinia (Redtip)
<i>Physotegia Virgiana</i>	Dragonhead, False; Vivid
<i>Picea spp.</i>	Spruce species** (Colorado Blue, Norway, etc.)

Scientific Name	Common Name
<i>Pieris japonica</i>	Lily-of-the-valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. elliotti</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. Thunbergiana</i>	Japanese Black Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. virginiana</i>	Virginia Pine
<i>Pistachio</i> sp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus Laurocerasus</i>	English Laurel
<i>Prunus</i> sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga Menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. Fortuneana</i>	Firethorn
<i>P. Koidzumii</i>	Forethorn
<i>Pyrus</i> sp.	Bradford Pear sp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus</i> spp.	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Raphiolepsis umbellata</i>	Yedda Hawthorn
<i>Rhododendron</i> (including azalea)	'Coral Bells' 'Delaware Valley White' 'Flame Creeper' 'Formosa' 'Girard Crimson' 'George L. Tabor' 'Hino-crimson' 'PJM' 'Roseum Elegans' 'Wakeiebsiu' 'White Gumbo'
<i>Rosa Banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia</i> sp.	Black-Eyed Susan; Goldstrum
<i>Santolina</i> virens	Santolina
<i>Saxifraga</i> sp.	Saxifrage; Purple Dome
<i>Scabiosa</i> sp.	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum cauticola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea Bumalda</i>	Spirea: Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushberry
<i>Taxus cuspidata</i>	Japanese Yew

Scientific Name	Common Name
<i>T. media</i>	Yew
<i>Teucrium sp.</i>	Germander
<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiatum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa spp.</i>	Tulip species
<i>Veronica sp.</i>	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca minor</i>	Common Periwinkle, Vinca
<i>Vitis sp.</i>	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filamentosa</i>	Adam's Needle, Yucca

*Ornamental species only. Do not use on food producing trees or vines.

NEW PLANTINGS, REPLANTING AND CROP ROTATION

If areas are to be seeded, do not apply RegalKade G until seeded plants are up and well established. Areas treated with RegalKade G should be rotated only to species listed on this label for 1 year following application unless the following test has shown species safety: Before planting a species not listed on this label it is recommended that several test strips of an indicator plant such as wheat, sorghum or corn be sown into the treated areas. If the indicator plants germinate and grow normally to a height of 12 inches with normal root development, it is safe to plant. In areas disturbed by new plantings or replanting of labeled species, it may be necessary to retreat exposed soil to maintain satisfactory weed control.

PRECAUTIONS AND LIMITATIONS

Do not apply to wet or moist foliage.

Do not apply in wind conditions that will affect product distribution. Uneven distribution will result in erratic weed control.

Do not use on any varieties or cultivars unless user is familiar with the results on specific plant materials under his growing conditions.

NOTICE TO USER: Plant tolerance to RegalKade G has been found to be acceptable on all ornamentals that it has been tested on. Due to the large number of species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to RegalKade G. Neither the manufacturer nor the seller has determined whether or not RegalKade G can be used safely on ornamental plants not specified on this label. The professional user should determine if RegalKade G can be used safely prior to commercial use. In a small area test the recommended rates on a small number of plants for phytotoxicity prior to widespread use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of this product, which are beyond the control of REGAL CHEMICAL COMPANY, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User, and Buyer and User agree to hold REGAL and Seller harmless for any claims relating to such factors.

REGAL warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or REGAL, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, REGAL MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither REGAL or Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF REGAL AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF REGAL OR SELLER, THE REPLACEMENT OF THE PRODUCT.

REGAL and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of REGAL.

REGAL CHEMICAL COMPANY
ALPHARETTA, GA 30004