

48273-7

8/18/2011

10946



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

AUG 18 2011

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mr. Matthew Granahan  
Nufarm Americas Inc.  
150 Harvester Dr., Suite 200  
Burr Ridge, IL 60527

Subject: Notification per PR Notice 2007-4 and 1998-10  
EPA Reg. No. 48273-7  
Marman Atila, Glyphosate  
Application dated July 29, 2011; revised 8570-1 form dated August 16, 2011

Dear Mr. Granahan:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 and 1998-10 of the subject product. The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and 1998-10 and finds that the action requested falls within the scope of PRN 2007-4 and 1998-10. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please contact Hope Johnson at 703-305-5410.

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis  
Product Manger 25  
Herbicide Branch  
Registration Division (7505P)



Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060

 <b>United States</b> <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	--	-----------------------

**Application for Pesticide - Section I**

<b>1. Company/Product Number</b> 48273-7	<b>2. EPA Product Manager</b> Bo Davis	<b>3. Proposed Classification</b> <input type="checkbox"/> None <input type="checkbox"/> Restricted
<b>4. Company/Product (Name)</b> Marman Atila, Glyphosate	<b>PM#</b> 25	
<b>5. Name and Address of Applicant (Include ZIP Code)</b> Marman USA, Inc. c/o Nufarm Americas Inc. 150 Harvester Drive, Suite 200 Burr Ridge, IL 60477  <input type="checkbox"/> Check if this is a new address		<b>6. Expedited Review.</b> In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: <b>EPA Reg. No.</b> <u>NOTIFICATION</u>  <b>Product Name</b> <u>AUG 18 2011</u>

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4 and PR Notice 1998-10. This notification is consistent with the guidance in PR Notice 2007-4 and PR Notice 1998-10 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, 156.156 and 152.46. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of PR Notice 1998-10 and 40 CFR §§ 156.10, 156.140, 156.144, 156.146, 156.156 and 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

<b>1. Material This Product Will Be Packaged In:</b>				<b>2. Type of Container</b>	
<b>Child-Resistant Packaging</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<b>Unit Packaging</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	<b>Water Soluble Packaging</b> <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	<input type="checkbox"/> Plastic
<b>* Certification must be submitted</b> If "Yes" Unit Packaging wgt.    No. per container    If "Yes" Package wgt    No. per container				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
<b>3. Location of Net Contents Information</b>		<b>4. Size(s) Retail Container</b>		<b>5. Location of Label Directions</b>	
<input checked="" type="checkbox"/> Label <input checked="" type="checkbox"/> Container		30, 55, 220, 250, 260, Gallons, Bulk		<input checked="" type="checkbox"/>	
<b>6. Manner in Which Label is Affixed to Product</b>				<input type="checkbox"/> Other Self-Adhesive Integrated Label/Booklet	
<input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

**Section - IV**

<b>1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)</b>		
<b>Name</b> Matthew Granahan matthew.granahan@us.nufarm.com	<b>Title</b> Regulatory Manager	<b>Telephone No. (Include Area Code)</b> (630) 455-2048
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		<b>6. Date Application Received (Stamped)</b>
<b>2. Signature</b> 	<b>3. Title</b> Regulatory Manager	
<b>4. Printed Name</b> Matthew Granahan	<b>5. Date</b> 08/16/2011	



30x46

**Nufarm Americas Inc.**

150 Harvester Drive, Suite 200

Burr Ridge, IL 60527

Telephone: (630) 455.2000 Facsimile: (630) 455.2001

www.us.nufarm.com

July 29, 2011

Via Overnight Courier

Bo Davis (PM-25)  
U. S. Environmental Protection Agency (7504P)  
Document Processing Desk (NOTIF)  
Room S4900, One Potomac Yard  
2777 S. Crystal Drive  
Arlington, VA 22202

**NOTIFICATION**

**AUG 18 2011**

**Subject: Marman Atila, Glyphosate  
EPA Reg. No. 48273-7  
Label Notification per PRN 07-04 & 98-10**

Dear Mr. Davis:


Marman USA would like to change Storage and Disposal text on above referenced label. It is believed this can be revised via notification and is consistent with PRN 07-04. Additionally Nufarm has updated the Warranty, changed the PHYSICAL OR CHEMICAL HAZARDS Section, added emergency contact information, and added the section HOT LINE NUMBER per PRN 98-10

To process this request please find enclosed the following:


- Application for Pesticide Registration EPA form 8570-1
- Certification with Respect to Label Integrity
- Revised labeling with areas of change clearly identified (1 copy)
- Revised labeling – clean (1 copy)
- CD containing the proposed labeling

If you should have any questions regarding this matter, please feel free to contact me at (630) 455-2048 or [matthew.granahan@us.nufarm.com](mailto:matthew.granahan@us.nufarm.com).

Sincerely,



Matthew Granahan  
Regulatory Manager  
Marman USA



{Container Label}

# MARMAN ATILA, GLYPHOSATE

**Water Soluble Herbicide for Nonselective Control of Many Annual and Perennial Weeds**

**Avoid Contact with Foliage, Green Stems, Exposed Non-Woody Roots or Fruit of Crops, Desirable Plants and Trees, Since Severe Injury or Destruction May Result**

**Active Ingredient:**

\*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt ..... 41%

**Other Ingredients\*** ..... 59%

Total ..... 100%

\*\*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate. This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

**KEEP OUT OF REACH OF CHILDREN**

## WARNING AVISO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

SEE [BACK] [SIDE] [INSIDE] [PANEL] [BOOKLET] FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

Net \_\_\_\_\_ gal.

EPA Reg. No. 48273-7      EPA Est. \_\_\_\_\_

Manufactured For  
**MARMAN USA, INC.**  
P.O. Box 22829  
Tampa, FL 33622-2829

**NOTIFICATION**

**AUG 18 2011**

{Container Label Back Panel}

### PRECAUTIONARY STATEMENTS

**Hazards to Humans & Domestic Animals**

**WARNING! AVISO!:** Causes substantial but temporary eye injury. Harmful if swallowed or inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, shoes plus socks and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

In case of an emergency involving this product, call collect, day or night (314) 694-4000.

**FIRST AID**

<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Immediately call a poison control center or doctor for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

{Note: The first aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

**Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**Physical or Chemical Hazards**

Solutions of this product should be mixed and stored using only stainless steel, aluminum, fiberglass, plastic and plastic-lined containers.

DO NOT MIX, STORE OR UTILIZE THIS PRODUCT OR SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR TANKS. This product or solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Read the entire label before using this product.**

**Read Warranty Disclaimer and Limitation of Liability before buying or using.**

**If terms are not acceptable, return at once unopened.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Refer to Booklet for complete Directions for Use.**

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep container closed to prevent spills and contamination. Store in a locked, cool, dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes of this product may cause irreversible eye damage and may be dangerous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to the label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING AND DISPOSAL:**

(This product is available in multiple containers. Follow the container disposal instructions below that apply to your container type.)

**[Nonrefillable containers larger than 5 gallons:]** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or

by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**[Refillable containers larger than 5 gallons:]** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Agricultural Use Requirements**


Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks and protective eyewear.

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried.



7 of 46

{Booklet Label}

# MARMAN ATILA, GLYPHOSATE

**Water Soluble Herbicide for Nonselective Control of Many Annual and Perennial Weeds**

**Avoid Contact with Foliage, Green Stems, Exposed Non-Woody Roots or Fruit of Crops, Desirable Plants and Trees, Since Severe Injury or Destruction May Result**

**Active Ingredient:**

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt .....	41%
<b>Other Ingredients*</b> .....	59%
<b>Total</b> .....	100%

\*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate. This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patents.

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail).

SEE [BACK] [SIDE] [INSIDE] [PANEL] [BOOKLET] FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

### Complete Directions for Use

EPA Reg. No. 48273-7      EPA Est. \_\_\_\_\_

Manufactured For  
**MARMAN USA, INC.**  
P.O. Box 22829  
Tampa, FL 33622-2829

This is an end-use product. Marman USA, Inc. does not intend and has not registered it for reformulation.  
See Individual Container Label for Repackaging Limitations.

### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF

8046

PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV072911)

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans & Domestic Animals**

**WARNING! AVISO!** Causes substantial but temporary eye injury. Harmful if swallowed or inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

<b>User Safety Recommendations</b>	
Users should:	
<ul style="list-style-type: none"> <li>Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> </ul>	
In case of an emergency involving this product, call collect, day or night (314) 694-4000.	

**FIRST AID**

<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

{Note: The first aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

**Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**Physical or Chemical Hazards**

Solutions of this product should be mixed and stored using only stainless steel, aluminum, fiberglass, plastic and plastic-lined containers.

DO NOT MIX, STORE OR UTILIZE THIS PRODUCT OR SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR TANKS. This product or solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Read the entire label before using this product. Use only according to label instructions.**

NO  
1A1



**Read Warranty Disclaimer and Limitation of Liability before buying or using.  
If terms are not acceptable, return at once unopened.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Keep container closed to prevent spills and contamination. Store in a locked, cool, dry place inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes of this product may cause irreversible eye damage and may be dangerous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to the label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER HANDLING AND DISPOSAL:

(This product is available in multiple containers. Follow the container disposal instructions below that apply to your container type.)

**[Nonrefillable containers larger than 5 gallons:]** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning is not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**[Refillable containers larger than 5 gallons:]** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box apply only to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, chemical resistant gloves Category A such as butyl rubber, natural rubber, neoprene rubber or nitrile rubber  $\geq 14$  mls., shoes plus socks and protective eyewear.

**Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried.

**General Information**

Do not apply this product using aerial spray equipment except under conditions as specified within this label.

This product, a water-soluble liquid, mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and through mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant that advances to complete browning of the above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the **Weeds Controlled** section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the herbicide and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust. Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut, and have not been allowed to regrow to the recommended stage for treatment. Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

**Note:** Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, other than unintended consequences.

**Mixing, Additives and Application Instructions**

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes. Do not apply when wind or other conditions favor drift. Hand-gun applications should be properly directed to avoid spraying desirable plants. NOTE: Reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

### Mixing

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of this product (see the **Directions for Use** and **Weeds Controlled** sections of this label) near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

### Tank Mixtures

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance. Mix labeled tank mixtures of this product with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it **slowly** through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture **slowly** through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Slowly add diluted mixture through the screen into the tank. Continue agitation.
6. Continue filling the spray tank with water and add the required amount of this product near the end of the filling process.
7. Where nonionic surfactant is recommended, add this to the spray tank before completing the filling process.
8. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

Immediately clean sprayer and parts after using this product by thoroughly flushing with water.

### Additives Surfactants

Nonionic surfactants that are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactant, use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants that contain at least 70 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 70 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

### Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, and this product plus 2,4-D, Banvel™ or residual herbicide tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve, which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactant. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

**Note:** The use of ammonium sulfate as an additive does not preclude the need for additional surfactant. Do not use herbicide rates lower than recommended on this label.

### Colorants or Dyes

Agriculturally-approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacture's recommendations.

### Application Equipment and Techniques

Do not apply this product through any type of irrigation system. This product may be applied with the following application equipment:

**Aerial:** Fixed Wing and Helicopter

#### Broadcast Spray

**Controlled Droplet Applicator (CDA):** Hand-held or boom-mounted applicators that produce a spray consisting of a narrow range of droplet sizes.

**Hand-Held and High-Volume Spray Equipment:** Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers\*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. \*This product is not registered in California or Arizona for use in mistblowers.

**Selective Equipment:** Recirculating sprayers, shielded sprayers and wiper applicators.

See the appropriate part of this section for specific instructions and application rates.

### Spray Drift Management:

**Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops.** Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

### Aerial Equipment

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. See the **Weeds Controlled** section of this label for specific rates. Unless otherwise specified, do not exceed 1 quart per acre. **Aerial applications of this product may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems and preharvest applications. Refer to the individual use area sections of this label for recommended volumes and application rates.**

### Aerial Spray Drift Management

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

**Importance of droplet size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduced drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity** and **Temperature Inversion** sections of this label).

### Controlling droplet size:

- **Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- **Pressure:** Use the lower spray pressures recommended for the nozzle. High pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of nozzles:** Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle orientation:** Orienting nozzles so that the spray is released backward, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- **Nozzle type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.
- **Boom length:** For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

- **Application height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, that causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should be applied only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

**Avoid direct application to any body of water.**

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

**Ensure uniform application:** To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **Prolonged exposure of this product to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.** The maintenance of an organic coating (paint), that meets aerospace specification MIL-C-38413, may prevent corrosion.

This product plus Oust™, Banvel™ or 2,4-D tank mixtures may not be applied by air in California.

**Broadcast Equipment**

**For control of annual or perennial weeds listed on this label using broadcast equipment:** Use the recommended rates of this product in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the **Weeds Controlled** section of this label for specific rates. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**Controlled Droplet Application (CDA)**

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre. For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of this product at a flow rate of 2 fluid ounces per minutes and a walking speed of 1.5 mph (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of this product at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 mph (2 to 4 quarts per acre). Controlled droplet application equipment

140x46

produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

**Hand-Held and High-Volume Equipment**

Use only coarse sprays.

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. For control of annual weeds listed on this label, apply a 0.5 percent solution of this product plus nonionic surfactant to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. Allow three or more days before tillage or mowing. For annual weeds over 6 inches tall, or when not using additional surfactant, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control perennials, such as bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle. When using application methods that result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees. Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

**Spray Solution**

Desired Volume	Amount of This Product					
	1/2%	1%	1-1/2%	2%	5%	10%
1 gal.	2/3 oz.	1-1/3 oz.	2 oz.	2-2/3 oz.	6-1/2 oz.	13 oz.
25 gal.	1 pt.	1 qt.	1-1/2 qt.	2 qt.	5 qt.	10 qt.
100 gal.	2 qt.	1 gal.	1-1/2 gal.	2 gal.	5 gal.	10 gal.

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

**Selective Equipment**

This product may be applied through a recirculating spray system, a shielded applicator, or a wiper applicator after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping system. A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse. A shielded applicator directs the herbicide solution onto weeds, while shielding desirable vegetation from the herbicide. A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution. **Avoid contact with desirable vegetation.**

Contact of the herbicide solution with the desirable vegetation may result in damage or destruction. Adjust applicators used above desired vegetation so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction. Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

**Shielded Applicators**

When applied as directed under conditions described for shielded applicators, this product will control those weeds listed in the **Weeds Controlled** section of this label. Use the following equation to convert from a broadcast rate per acre to a band rate per acre:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Herbicide Broadcast Rate per Acre} = \text{Herbicide Band Rate per Acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Volume of Solution per Acre} = \text{Band Volume of Solution per Acre}$$

Use nozzles that provide uniform coverage within the treated area. Keep shields on shielded sprayers adjusted to protect desirable vegetation. **Extreme care must be exercised to avoid contact with desirable vegetation.** For specific application rates and instructions for control of various annual and perennial weeds, see the **Weeds Controlled** section of this label.

**Wiper Applicators**

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions. Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator. Do not use wiper equipment when weeds are wet. Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water. Do not add surfactant to the herbicide solution.

**For Rope or Sponge Wick Applicators:** Mix 1 gallon of this product in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this *Wiper Applicators* section.

**For Porous-Plastic Applicators:** Solutions ranging from 33 to 100 percent of this product in water may be used in porous-plastic wiper applicators.

When applied as recommended under the conditions described for *Wiper Applicators*, this product **controls** the following weeds:

#### Annual Grasses

Corn	Rye, common
<i>Zea mays</i>	<i>Secale cereale</i>
Texas Panicum	Shattercane
<i>Panicum texanum</i>	<i>Sorghum bicolor</i>

#### Annual Broadleaves

Sicklepod	Starbur, bristly
<i>Cassia obtusifolia</i>	<i>Acanthospermum hispidum</i>
Spanishneedles	
<i>Bidens bipinnata</i>	

When applied as recommended under the conditions described for *Wiper Applicators*, this product **suppresses** the following weeds:

#### Annual Broadleaves

Beggarweed, Florida	Ragweed, giant
<i>Desmodium tortuosum</i>	<i>Ambrosia trifida</i>
Dogfennel	Sunflower
<i>Eupatorium capilliflorum</i>	<i>Helianthus annuus</i>
Pigweed, redroot	Thistle, musk
<i>Amaranthus retroflexus</i>	<i>Carduus nutans</i>
Ragweed, common	Velvetleaf
<i>Ambrosia artemisiifolia</i>	<i>Abutilon theophrasti</i>

#### Perennial Grasses

Bermudagrass	Smutgrass
<i>Cynodon dactylon</i>	<i>Sporobolus poiretii</i>
Guineagrass	Vaseygrass
<i>Panicum maximum</i>	<i>Paspalum urvillei</i>
Johnsongrass	
<i>Sorghum halepense</i>	

#### Perennial Broadleaves

Dogbane, hemp	Nightshade, silverleaf
<i>Apocynum cannabinum</i>	<i>Solanum elaeagnifolium</i>
Milkweed	Thistle, Canada
<i>Asclepias syriaca</i>	<i>Cirsium arvense</i>

#### Weeds Controlled

This herbicide controls many annual and perennial grasses and broadleaf weeds.

#### Annual Weeds

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage.
- For maximum agronomic benefit, apply when weeds are 6 inches or less in height.

- To prevent seed production, apply prior to seedhead formation.
- This product does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

**Low-Volume Broadcast Application  
(Low-Rate Technology)**

When applied as directed under the conditions described, this product will control the weeds listed below when:

1. Water carrier volumes of 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications are recommended. (See the **Aerial Application and Techniques** section of this label for approved sites.)
2. A nonionic surfactant is added at 0.5 to 1 percent by total spray volume. Use 0.5 percent surfactant concentration when using surfactants that contain at least 70 percent active ingredient or a 1 percent surfactant concentration for those surfactants containing less than 70 percent active ingredient.

**Note:**

- The addition of 2 percent dry ammonium sulfate by weight of 17 pounds per 100 gallons of water may increase the performance of this product on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the **Mixing, Additives and Application Instructions** section of this label.
- Do not tank-mix with soil residual herbicides when using these rates unless otherwise specified.
- For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.
- Refer to the **Tank Mixtures** portion of this section for control of additional broadleaf weeds.

Weed Species	Maximum Height/Length	Rate per Acre* (Fluid Ounces)
Foxtail ( <i>Setaria</i> spp.)	12"	8 oz.
Barnyardgrass ( <i>Echinochloa crus-galli</i> )	6" (0 to 4" 4 to 6")	12 oz. 16 oz. <sup>1)</sup> 24 oz. <sup>1)</sup>
Bluegrass, annual ( <i>Poa annua</i> )		
Brome, downy** ( <i>Bromus tectorum</i> )		
Mustard, blue ( <i>Chorispora tenella</i> )		
Mustard, tansy ( <i>Descurainia pinnata</i> )		
Mustard, tumble ( <i>Sisymbrium altissimum</i> )		
Mustard, wild ( <i>Sinapis arvensis</i> )		
Spurry, umbrella ( <i>Holosteum umbellatum</i> )		
Barley ( <i>Hordeum vulgare</i> )	12"	
Rye ( <i>Secale cereale</i> )		
Sandbur, field ( <i>Cenchrus</i> spp.)		
Shattercane ( <i>Sorghum bicolor</i> )		
Stinkgrass ( <i>Eragrostis ciliaris</i> )		
Wheat ( <i>Triticum aestivum</i> )	18"	
Morningglory ( <i>Ipomoea</i> spp.)	2"	16 oz.
Sicklepod ( <i>Casia obtusifolia</i> )		
Bluegrass, bulbous ( <i>Poa bulbosa</i> )	6"	
Cheat ( <i>Bromus secalinus</i> )		
Chickweed, common ( <i>Stellaria media</i> )		
Chickweed, mouseear ( <i>Cerastium vulgatum</i> )		
Corn ( <i>Zea mays</i> )		
Goatgrass, jointed ( <i>Aegilops cylindrica</i> )		
Groundsel, common ( <i>Senecio vulgaris</i> )		
Henbit ( <i>Lamium amplexicaule</i> )		
Horseweed/Marestail ( <i>Conyza canadensis</i> )		
Lambsquarters, common ( <i>Chenopodium album</i> )		
Pennycress, field ( <i>Fanweed, Thlaspi arvense</i> )		
Rocket, London ( <i>Sisymbrium irio</i> )		
Ryegrass, Italian ( <i>Lolium multiflorum</i> )		
Shepherdspurse ( <i>Capsella bursa-pastoris</i> )		
Spurge, annual ( <i>Euphorbia</i> spp.)		
Buttercup ( <i>Ranunculus</i> spp.)	12"	



Cocklebur ( <i>Xanthium strumarium</i> )		
Crabgrass ( <i>Digitaria</i> spp.)		
Dwarf dandelion ( <i>Krigia cespitosa</i> )		
False flax, smallseed ( <i>Camelina microcarpa</i> )		
Foxtail, Carolina ( <i>Alopecurus carolinianus</i> )		
Johnsongrass, seedling ( <i>Sorghum halepense</i> )		
Oats, wild ( <i>Avena fatua</i> )		
Panicum, fall ( <i>Panicum dichotomiflorum</i> )		
Panicum, Texas ( <i>Panicum texanum</i> )		
Pigweed, redroot ( <i>Amaranthus retroflexus</i> )		
Pigweed, smooth ( <i>Amaranthus hybridus</i> )		
Witchgrass ( <i>Panicum capillare</i> )		
Sicklepod ( <i>Cassia obtusifolia</i> )	3 to 4"	24 oz.
Signalgrass, broadleaf ( <i>Brachiaria platyphylla</i> )	4"	
Horseweed/Marestail ( <i>Conzya canadensis</i> )	7 to 12"	
Lambsquarters, common ( <i>Chenopodium album</i> )		
Spurge, annual ( <i>Euphorbia</i> spp.)		
Rice, red ( <i>Oryza sativa</i> )	4"	32 oz.
Teaweed ( <i>Sida spinosa</i> )		
Sprangletop ( <i>Leptochloa</i> spp.)	6"	
Geranium, Carolina ( <i>Geranium carolinianum</i> )	12"	
Goosegrass ( <i>Eleusine indica</i> )		
Primrose, cutleaf evening ( <i>Oenothera laciniata</i> )		
Pusley, Florida ( <i>Richardia scabra</i> )		
Sicklepod ( <i>Cassia obtusifolia</i> )	5 to 12"	
Spanishneedles ( <i>Bidens bipinnata</i> )		
Filaree ( <i>Erodium</i> spp.)	12"	48 oz.
Sprangletop ( <i>Leptochloa</i> spp.)		

Use these rates to control barnyardgrass in Alabama, Arkansas, Mississippi, Missouri, Louisiana and Texas for pre-plant treatments.

\*For those rates less than 32 fluid ounces per acre, this product at rates up to 32 fluid ounces per acre may be used where heavy weed densities exist.

\*\*For control in no-till systems, use 16 ounces per acre.

**Tank Mixtures**

This Product plus Banvel plus Nonionic Surfactant  
 This product plus 2,4-D plus Nonionic Surfactant

**Do not apply Banvel or 2,4-D Tank Mixtures by Air in California**

These tank mixtures are recommended only for use in fallow and reduced tillage areas. Follow use directions as given in the **Low-Volume Broadcast Application** section. This product plus Banvel or 2,4-D will control the annual grasses and broadleaf weeds listed for this product alone at the indicated heights (except 8 fluid ounces per acre applications), plus the following broadleaf weeds. For those weeds previously listed at 8 fluid ounces of this product alone per acre, use 12 fluid ounces in these tank mixtures.

**Note:** Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if Banvel is applied within 45 days of planting. The addition of Banvel in a mixture with this product may provide short-term residual control of selected weed species.

Apply 12 to 16 fluid ounces of this product plus 0.25 lb. a.i. of Banvel or 0.5 pound a.i. of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre to control dense populations of the following annual broadleaf weeds when less than the height indicated:

- |   |   |
|---|---|
| Cocklebur (12")<br><i>Xanthium strumarium</i>   | Morningglory (6")<br><i>Ipomoea</i> spp.                |
| Kochia* (6")<br><i>Kochia scoparia</i>          | Pigweed, redroot (12")<br><i>Amaranthus retroflexus</i> |
| Lambsquarters (12")<br><i>Chenopodium album</i> | Pigweed, smooth (12")<br><i>Amaranthus hybridus</i>     |
| Lettuce, lettuce (6")                           | Thistle, Russian (12")                                  |

*Lactuca serriola*  
Marestail/Horseweed (6")  
*Conzya canadensis*  
\*Controlled only with Banvel tank mixture.

*Salsola kali*

Apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre to control the following annual broadleaf weeds when less than 6 inches in height:

Ragweed, common <i>Ambrosia artemisiifolia</i>	Smartweed, Pennsylvania <i>Polygonum pensylvanicum</i>
Ragweed, giant <i>Ambrosia trifida</i>	Velvetleaf <i>Abutilon theophrasti</i>

### High-Volume Broadcast Applications

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes are 10 to 40 gallons per acre for ground applications. Apply 1 to 1.5 quarts of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume. Use 1 quart per acre if weeds are less than 6 inches tall and 1.5 quarts per acre if weeds are over 6 inches tall. If weeds have been mowed, grazed or cut, allow adequate time for new growth to reach recommended stages prior to treatment. These rates will also provide control of weeds listed in the **Low-Volume Broadcast Application** section.

### Weed Species

Balsamapple* <i>Momordica charantia</i>	Panicum <i>Panicum</i> spp.
Brassia, fivehook <i>Brassi hyssopifolia</i>	Ragweed, common <i>Ambrosia artemisiifolia</i>
Brome <i>Bromus</i> spp.	Ragweed, giant <i>Ambrosia trifida</i>
Fiddleneck <i>Amsinckia</i> spp.	Smartweed, Pennsylvania <i>Polygonum pensylvanicum</i>
Fleabane, hairy <i>Conzya bonariensis</i>	Sowthistle, annual <i>Sonchus cleraceus</i>
Fleabane <i>Erigeron</i> spp.	Sunflower <i>Helianthus annuus</i>
Kochia <i>Kochia scoparia</i>	Thistle, Russian <i>Salsola kali</i>
Lettuce, prickly <i>Lactuca serriola</i>	Velvetleaf <i>Abutilon theophrasti</i>

\*Apply only with hand-held equipment.

### Perennial Weeds

Apply this product as follows to control or destroy most perennial weeds:

**Note:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence. The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the **Mixing, Additives and Application Instructions** section of this label. When applied as recommended under the conditions described, this product will control the following perennial weeds:

Alfalfa <i>Medicago sativa</i>	Knapweed <i>Centaurea repens</i>
Alligatorweed* <i>Alternanthera philoxeroides</i>	Lantana <i>Lantana camara</i>
Anise (fennel) <i>Foeniculum vulgare</i>	Lespedeza <i>Lespedez</i> spp.
Artichoke, Jerusalem <i>Helianthus tuberosus</i>	Milkweed <i>Asclepias</i> spp.
Bahiagrass*	Muhly, wirestem

<i>Paspalum notatum</i>	<i>Muhlenbergia frondosa</i>
Bentgrass	Mullein, common
<i>Agrostis</i> spp.	<i>Verbascum thapsus</i>
Bermudagrass	Napiergrass
<i>Cynodon dactylon</i>	<i>Pennisetum purpureum</i>
Bermudagrass, water (Knotgrass)	Nightshade, silverleaf
<i>Paspalum distichum</i>	<i>Solanum elaeagnifolium</i>
Bindweed, field	Nutsedge, purple and yellow
<i>Convolvulus arvensis</i>	<i>Cyperus rotundus</i> and <i>Cyperus esculentus</i>
Bluegrass, Kentucky	Orchardgrass
<i>Poa</i> spp.	<i>Dactylis glomerata</i>
Blueweed, Texas	Pampasgrass
<i>Helianthus ciliaris</i>	<i>Cortaderia</i> spp.
Brackenfern	Paragrass
<i>Pteridium aquilinum</i>	<i>Brachiaria mutica</i>
Bromegrass, smooth	Phragmites*
<i>Bromus inermis</i>	<i>Phragmites</i> spp.
Bursage, woollyleaf	Poison hemlock
<i>Franseria tomentosa</i>	<i>Conium maculatum</i>
Canarygrass, reed	Quackgrass
<i>Phalaris arundinacea</i>	<i>Agropyron repens</i>
Cattail	Redvine*
<i>Typha</i> spp.	<i>Burnnichia ovata</i>
Clover, red	Reed, giant
<i>Trifolium pratense</i>	<i>Arundo donax</i>
Clover, white	Ryegrass, perennial
<i>Trifolium repens</i>	<i>Lolium perenne</i>
Cogongrass	Smartweed, swamp
<i>Imperata cylindrica</i>	<i>Polygonum coccineum</i>
Dallisgrass	Spurge, leafy*
<i>Paspalum dilatatum</i>	<i>Euphorbia esula</i>
Dandelion	Starthistle, yellow
<i>Taraxacum officinale</i>	<i>Centaurea solstitialis</i>
Dock, curly	Sweet Potato, wild*
<i>Rumex crispus</i>	<i>Ipomoea pandurata</i>
Dogbane, hemp	Thistle, artichoke
<i>Apocynum cannabinum</i>	<i>Cyanara cardunculus</i>
Fescues	Thistle, Canada
<i>Festuca</i> spp.	<i>Cirsium arvense</i>
Fescue, tall	Timothy
<i>Festuca arundinacea</i>	<i>Phleum pratense</i>
Guineagrass	Torpedograss*
<i>Panicum maximum</i>	<i>Panicum repens</i>
Horsenettle	Trumpet creeper*
<i>Solanum carolinense</i>	<i>Campsis radicans</i>
Horseradish	Vaseygrass
<i>Armoracia rusticana</i>	<i>Paspalum urvillei</i>
Ice Plant	Velvetgrass
<i>Mesembryanthemum crystallinum</i>	<i>Holcus</i> spp.
Johnsongrass	Wheatgrass, western
<i>Sorghum halepense</i>	<i>Agropyron smithii</i>
Kikuyugrass	
<i>Pennisetum clandestinum</i>	
*Partial control	

This product is not registered in California for use on water bermudagrass.

See **Directions for Use** and **Mixing, Additives and Application Instructions** sections of this label for labeled uses and specific application instructions.

**Alfalfa:** Apply 1 quart of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Make application after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Follow applications with deep tillage at least 7 days after treatment, but before soil freeze-up.

**Alligatorweed:** Apply 4 quarts of this product per acre or apply a 1.5 percent solution with hand-held equipment to provide partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.

**Anise (fennel) / Poison Hemlock:** Apply a 1 to 2 percent solution of this product as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Repeat applications may be needed in succeeding years to control plants arising from seeds.

**Bentgrass:** For suppression in grass seed production areas. Only for ground applications, apply 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should be actively growing and have at least 3 inches of growth. Avoid tillage prior to treatment. Tillage 7 to 10 days after application is recommended for best results. Failure to use tillage after treatment may result in unacceptable control.

**Bermudagrass:** For control, apply 5 quarts of this product per acre. For partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control. Allow 7 or more days after application before tillage.

**Bermudagrass, water (knotgrass):** Apply 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Apply when water bermudagrass is actively growing and 12 to 18 inches in length. Allow 7 or more days before tilling, flushing or flooding the field.

Fall applications only: Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 5 to 10 gallons of water per acre. Till fallow fields prior to application. Apply prior to frost on water bermudagrass that is actively growing and 12 to 18 inches in length. Allow 7 or more days before tillage.

**Bindweed, field:** For control apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts each of the Mississippi River. Apply when the weeds are actively growing and are at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. For best results, apply in late summer or fall. Fall treatment must be applied before a killing frost. Allow 7 or more days after application before tillage. Also for control, apply 2 quarts of this product plus 0.5 pound a.i. of Banvel in 10 to 20 gallons of water per acre. At these rates, apply using only ground application.

The following tank mixtures with 2,4-D may be applied using aerial application equipment (except in California) only in fallow and reduced tillage systems. For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product plus 1 pound a.i. of 2,4-D in 10 to 20 gallons of water per acre with only ground equipment. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth. For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

**Only in California,** apply 1 to 5 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending upon local conditions.

For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. Allow 3 or more days after application before tillage.

**Bluegrass, Kentucky / Bromegrass, smooth / Orchardgrass:** Apply 2 quarts of this product in 10 to 40 gallons of water per acre when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.

**Orchardgrass (sods going to no-till corn):** Apply 1 to 1.5 quarts of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine is necessary for optimum results.

**Blueweed, Texas:** Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.

**Brackenfern:** Apply 3 to 4 quarts of this product per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment. Apply to fully expanded fronds that are at least 18 inches long.

**Bursage, woollyleaf:** For control, apply 2 quarts of this product plus 1 pint of Banvel per acre. For partial control, apply 1 quart of this product plus 1 pint of Banvel per acre. Add 0.5 to 1 percent nonionic surfactant by total spray volume and apply in 3 to 20 gallons of water per acre. Apply when plants are producing new active growth that has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

**Canarygrass, reed / Timothy / Wheatgrass, western:** Apply 2 to 3 quarts of this product per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.

**Cogongrass:** Apply 3 to 5 quarts of this product plus 0.5 to 1 percent nonionic surfactant in 10 to 40 gallons of water per acre. Apply when Cogongrass is at least 18 inches tall and actively growing in last summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

**Dandelion / Dock, curly:** Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage. Also for control, apply 16 fluid ounces of this product plus 0.5 pound a.i. 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre.

**Dogbane, hemp:** Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage. For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay applications until maximum emergence of dogbane has occurred.

**Fescue, tall:** Apply 3 quarts of this product in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development. Fall applications only: Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. Allow 7 or more days after application before tillage. A sequential application of 1 pint per acre of this product plus nonionic surfactant will improve long-term control and control seedlings germinating after fall treatments of the following spring.

**Guineagrass:** Apply 3 quarts of this product per acre or use a 1 percent solution with hand-held equipment. Apply to actively growing guineagrass when most has reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.

**Johnsongrass / Ryegrass, perennial:** Apply 1 to 3 quarts of this product per acre. In annual cropping systems apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In noncrop, or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank-mix with residual herbicides when using the 1 quart per acre rate. For burndown of Johnsongrass, apply 1 pint per acre plus 0.6 to 1 percent nonionic surfactant in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage. Spot treatment (partial control or suppression): Apply a 1 percent

solution of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

**Kikuyugrass:** Apply 2 to 3 quarts of this product per acre. Spray when most kikuyugrass is at least 8 inches in height (3- or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

**Knapweed / Horseradish:** Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

**Latana:** Apply this product as a 1 to 1.25 percent solution using only hand-held equipment. Apply to actively growing latana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.

**Milkweed, common:** Apply 3 quarts of this product per acre. Apply when actively growing and most of the milkweed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.

**Muhly, wirestem:** Apply 1 to 2 quarts of this product per acre. Use 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture, sod or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. This product will not provide residual control of wirestem muhly from seeds that germinate after application of this product. Do not tank mix with residual herbicides when using the 1 quart per acre rate.

**Nightshade, silverleaf:** For control, apply 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant total spray volume in 30 to 10 gallons of water per acre. Apply when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth.

**Nutsedge, purple, yellow:** Apply 3 quarts of this product per acre as a broadcast spray, or apply a 1 to 2 percent solution from hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers. Sequential applications of 1 to 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre will provide control. Apply when a majority of the plants are in the 3- to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control. For suppression to partial control of existing plants, apply 1 pint to 2 quarts of this product per acre, plus 0.5 to 1 percent nonionic surfactant in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants. Wait 7 days after treatment before tillage or mowing.

**Pampasgrass / Ice Plant:** Apply this product as a 1.5 to 2 percent solution using hand-held equipment. Apply to plants that are actively growing. Pampas grass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

**Phragmites:** For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment. In other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

**Quackgrass – In Annual Cropping Systems or in Pastures and Sods Followed by Deep Tillage:** Apply 1 to 2 quarts of this product per acre. For the 1 quart rate, apply 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. For the 2 quart rate, apply in 10 to 40 gallons of water per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 6 to 8 inches in height and actively growing. Do not till

between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, for best results, use a moldboard plow.

**Quackgrass – Pasture or Sod or Other Noncrop Areas where Deep Tillage is not Planned Following Application:** Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.

**Redvine:** For suppression, apply 24 fluid ounces of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre plus 0.5 to 1 percent nonionic surfactant by total volume. Apply to actively growing plants, in late September or early October, that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

**Reed, giant:** For control of giant reed, apply a 2 percent solution of this product when plants are actively growing. Best results are obtained when applications are made in late summer to fall.

**Smartweed, swamp:** Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage. Also for control, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D plus 0.5 to 1 percent nonionic surfactant by total volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.

**Spurge, leafy:** For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. 2,4-D plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.

**Starthistle, yellow:** Best results are obtained when applications are made during periods of active growth, including the rosette, bolting and early flower stages. For spray-to-wet applications, apply this product as a 2 percent solution. For broadcast applications, apply 2 quarts per acre in 10 to 40 gallons per acre of water carrier.

**Sweet Potato, wild / Thistle, artichoke:** Apply this product as a 2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.

**Thistle, Canada:** Apply 2 to 3 quarts of this product per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage. For suppression of Canada thistle, apply 1 quart per acre of this product, or 1 pint of this product plus 0.5 pound a.i. 2,4-D per acre, plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

**Torpedograss:** Apply 4 to 5 quarts of this product per acre to provide partial control of torpedograss. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.

**Trumpet creeper:** For control, apply 2 quarts of this product per acre in 5 to 10 gallons of water per acre. Apply to actively growing plants, in late September or October, that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

**Other Perennials Listed on this Label:** Apply 3 to 5 quarts of this product per acre. Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

#### Woody Brush and Trees

When applied as recommended under the conditions described, this product controls or partially controls the following woody brush, plants and trees:

Alder  
*Alnus* spp.  
Ash\*  
*Fraxinus* spp.  
Aspen, quaking  
*Populus tremuloides*  
Bearmat (Bearclover)  
*Chamaebatia foliolosa*  
Beech  
*Fagus grandifolia*  
Birch  
*Betula* spp.  
Blackberry  
*Rubus* spp.  
Blackgum  
*Nyssa* spp.  
Bracken  
*Peridium* spp.  
Broom, French  
*Cytisus monspessulanus*  
Broom, Scotch  
*Cytisus scoparius*  
Buckwheat, California\*  
*Eriogonum fasciculatum*  
Cascara\*  
*Rhamnus purshiana*  
Catsclaw\*  
*Acacia greggi*  
Ceanothus\*  
*Ceanothus* spp.  
Chamise  
*Adenostoma fasciculatum*  
Cherry, Bitter  
*Prunus emarginata*  
Cherry, Black  
*Prunus serotina*  
Cherry, Pin  
*Prunus pensylvanica*  
Coyote brush  
*Baccharis consanguinea*  
Creeper, Virginia\*  
*Parthenocissus quinquefolia*  
Dewberry  
*Rubus trivialis*  
Dogwood  
*Cornus* spp.  
Elderberry  
*Sambucus* spp.  
Elm\*  
*Ulmus* spp.  
Eucalyptus  
*Eucalyptus* spp.  
Gorse  
*Ulex europaeus*  
Hasardia\*  
*Haplopappus squamosus*  
Hawthorn  
*Crataegus* spp.  
Hazel  
*Corylus* spp.  
Hickory\*  
Maple, Red\*\*  
*Acer rubrum*  
Maple, Sugar  
*Acer saccharum*  
Maple, Vine\*  
*Acer circinatum*  
Monkey Flower\*  
*Mimulus guttatus*  
Oak, Black\*  
*Quercus velutina*  
Oak, Northern Pin  
*Quercus palustris*  
Oak, Post  
*Quercus stellata*  
Oak, Red  
*Quercus rubra*  
Oak, Southern Red  
*Quercus falcata*  
Oak, White\*  
*Quercus alba*  
Permisson\*  
*Diospyros* spp.  
Pine  
*Pinus* spp.  
Poison Ivy  
*Rhus radicans*  
Poison Oak  
*Rhus toxicodendron*  
Poplar, yellow\*  
*Liniodendron tulipifera*  
Raspberry  
*Rubus* spp.  
Redbud, eastern  
*Ceris canadensis*  
Rose, multiflora  
*Rosa multiflora*  
Russian-olive  
*Elaeagnus angustifolia*  
Sage, black, white  
*Salvia* spp.  
Sagebrush, California  
*Artemisia californica*  
Salmonberry  
*Rubus spectabilis*  
Salt cedar  
*Tamarix* spp.  
Sassafras  
*Sassafras aibidum*  
Sourwood  
*Oxydendrum arboreum*  
Sumac, Poison\*  
*Rhus vernix*  
Sumac, Smooth\*  
*Rhus glabra*  
Sumac, Winged\*  
*Rhus copallina*  
Sweetgum  
*Liquidambar styraciflua*  
Swordfern\*  
*Polystichum munitum*  
Tallowtree, Chinese



*Carya* spp.  
 Holly, Florida / Brazilian Peppertree  
*Schinus terebinthifolius*  
 Honeysuckle  
*Lonicera* spp.  
 Hornbeam, American\*  
*Carpinus caroliniana*  
 Kudzu  
*Pueraria lobata*  
 Locust, black\*  
*Robinia pseudoacacia*  
 Madrone  
*Arbutus menziesii*  
 Manzanita  
*Arctostaphylos* spp.

*Sapium sebiferum*  
 Tan Oak  
*Lithocarpus densiflorus*  
 Thimbleberry  
*Rubus parviflorus*  
 Tobacco, tree\*  
*Nicotiana glauca*  
 Trumpet creeper  
*Campsis radicans*  
 Waxmyrtle, southern\*  
*Myrica cerifera*  
 Willow  
*Salix* spp.

\*Partial control

\*\*See below for control or partial control instructions.

**Note:** If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth.

Apply this product when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments. Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See **Directions for Use** and **Mixing, Additives and Application Instructions** sections of this label for labeled uses and specific application instructions. Apply this product as follows to **control** or **partially control** the following woody brush and trees:

**Alder / Dewberry / Honeysuckle / Post Oak / Raspberry:** For control, apply 3 to 4 quarts per acre of this product as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.

**Aspen, quaking / Cherry: bitter, black, pin / Hawthorn / Oak, southern red / Sweetgum / Trumpet creeper:** For control, apply 2 to 3 quarts of this product per acre as a broadcast spray or as a 1 to 1.5 percent solution with hand-held equipment.

**Birch / Elderberry / Hazel / Salmonberry / Thimbleberry:** For control, apply 2 quarts per acre of this product as a broadcast spray or as a 1 percent solution with hand-held equipment.

**Blackberry:** For control, apply 3 to 4 quarts per acre of this product as a broadcast spray, or 1 to 1.5 percent solution with hand-held equipment. Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a ¾ percent solution of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.

**Broom: French, Scotch:** For control, apply a 1.5 to 2 percent solution with hand-held equipment.

**Buckwheat, California / Hasardia / Monkey Flower / Tobacco, tree:** For partial control of these species, apply a 1 to 2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

**Catsclaw:** For partial control, apply as a 1 to 1.5 percent solution with hand-held equipment.

**Coyote Brush:** For control, apply a 1.5 to 2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

**Eucalyptus:** For control of eucalyptus resprouts, apply a 2 percent solution of this product with hand-held equipment when resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants.

**Kudzu:** For control, apply 4 quarts of this product per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.

**Madrone resprouts:** For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.

**Maple, red:** For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre as a broadcast spray.

**Maple, sugar / Oak, northern pin / Oak, red:** For control, apply as a 1 to 1.5 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

**Poison Ivy / Poison Oak:** For control, apply 4 to 5 quarts of this product per acre as a broadcast spray or as a 2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.

**Rose, multiflora:** For control, apply 2 quarts of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply prior to leaf deterioration by leaf-feeding insects.

**Sage, black / Sagebrush, California / Chamise / Tallowtree, Chinese:** For control of these species, apply a 1 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.

**Tan oak resprouts:** For suppression or partial control, apply a 2 percent solution of this product to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.

**Willow:** For control, apply 3 quarts of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment.

**Other Woody Brush and Trees Listed on this Label:** For partial control, apply 2 to 4 quarts of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment.

#### Noncrop Uses

See **General Information** and **Mixing, Additives and Application Instructions** sections of this label for essential product performance information and the following **Noncrop** sections for specific recommended uses. **Extreme care must be exercised to avoid contact of spray with foliage, green stems, exposed non-woody roots or fruit or crops, desirable turfgrasses, trees, shrubs or other desirable vegetation since severe damage or destruction may result.**

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year. The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate- or sulfosate-containing products does not exceed the stated maximum use rate.

This product does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

#### Industrial, Recreational and Public Areas

When applied as directed for **Noncrop Uses**, under conditions described, this product controls annual and perennial weeds listed on this label growing in areas such as airports, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumber yards, parking areas, parks, petroleum tank farms and pumping installations,

pipelines, power and telephone rights-of-way, railroads, roadsides, schools, storage areas, utility substations, other public areas and similar industrial or noncrop areas.

For specific application rates and instructions for control of various annual and perennial weeds and woody brush and trees, see the **Weeds Controlled** section of this label. This product may be applied with recirculating sprayers, shielded applicators or wiper applicators in any noncrop site specified on this label. See the **Selective Equipment** part of **Application Equipment and Techniques** section of this label for information on proper use and calibration of this equipment.

**Tank Mixtures for Industrial Sites and Forestry Site Preparations**  
**This Product Plus Oust™**

Use on industrial sites including airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, pipelines, railroads, roadsides, storage areas or other similar sites where bare ground is desired. This tank mixture may also be used as a site preparation treatment for sites to be planted to jackpine, loblolly pine, red pine, slash pine and Virginia pine. When applied as directed for **Noncrop Uses** under the conditions described, this product plus Oust provides control of annual weeds listed in the **Weeds Controlled** section of the label for this product and Oust, and control or partial control of the perennial weeds listed below. Apply 1 to 2 quarts of this product with 2 to 4 ounces of Oust in 10 to 40 gallons of spray solution per acre as a broadcast spray to actively growing weeds. This mixture may be applied by aerial equipment in site prep operations. When applied by air, use the recommended rates in 5 to 15 gallons of spray solution per acre. This product plus Oust tank mixtures may not be applied by air in California. For control of annual weeds, use the lower rates of these products. For control of the listed perennial weeds, use the higher rates of both products. For partial control, use the lower rates.

- |  |  |
|--|--|
| Bahiagrass<br><i>Paspalum notatum</i>        | Johnsongrass**<br><i>Sorghum halepense</i> |
| Bermudagrass*<br><i>Cynodon dactylon</i>     | Poorjoe**<br><i>Diodia teres</i>           |
| Broomsedge<br><i>Andropogon virginicus</i>   | Quackgrass<br><i>Agropyron repens</i>      |
| Dock, curly<br><i>Rumex crispus</i>          | Trumpetcreeper*<br><i>Campsis radicans</i> |
| Dogfennel<br><i>Eupatorium capilliflorum</i> | Vaseygrass<br><i>Paspalum urvillei</i>     |
| Fescue, tall<br><i>Festuca arundinacea</i>   | Vervain, blue<br><i>Verbena hastata</i>    |

\*Suppression only at the higher rates.  
\*\*Control at the lower rates.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

**Tank Mixtures**  
**Noncrop Sites**

When applied as a tank mixture, this product provides control of the emerged annual weeds and partial control of the emerged perennial weeds listed in this label. When applied as a tank mixture, the following residual herbicides will provide preemergence control of the weeds listed in the individual product labels.

- This Product Plus Diuron**
- This Product Plus Krovar™ I**
- This product Plus Krovar II**
- This product Plus Ronstar™ 50WP**
- This Product Plus Simazine Princep Caliber™ 90**
- This Product Plus Simazine 4L**
- This Product Plus Simazine 80W**
- This Product Plus Surflan™ 75W**
- This Product Plus Surflan AS**

When tank mixing with residual herbicides, add an agriculturally approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution. See the **Mixing, Additives and Application Instructions** section of this label before preparing these tank mixtures. Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

**Note:** For backpack sprayer and handgun applications, see the *Hand-Held and High Volume Equipment* section for recommended rates.

**Annual Weeds:** Apply 1 quart per acre of this product in these tank mixtures when weeds are less than 6 inches tall and 1.5 quarts per acre when weeds are more than 6 inches tall.

**Perennial Weeds:** For partial control of perennial weeds using these tank mixtures, apply 2 to 5 quarts per acre of this product. Follow the recommendations in the *Weeds Controlled* section of this label for stage of growth and application rates for specific perennial weeds.

#### Preemergence Weed Control

For preemergence weed control, refer to the individual product labels for specific noncrop sites, rates, carrier volumes and precautionary statements. Mix only the quantity of spray solution that can be used during the same day. Do not allow tank mixtures to stand overnight as this may result in reduced weed control.

#### Farmstead Weed Control

When applied as directed for *Noncrop Uses* under conditions described, this product controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts and for general nonselective farmstead weed control. For specific application rates and instructions for control of various annual and perennial weeds, see the *Weeds Controlled* section of this label.

#### Farm Ditches

This product will suppress perennial grasses along farm ditches. Apply this product at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces per acre when treating tall (coarse) fescue, fine fescue, orchardgrass or quackgrass covers. For best suppression of these species, add ammonium sulfate at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces per acre without ammonium sulfate when treating Kentucky bluegrass. Apply treatments in 10 to 20 gallons of spray solution per acre to actively growing perennial grass covers. For best spray distribution and coverage, use flat fan nozzles. Add a nonionic surfactant at a rate of 0.5 percent of the spray solution. Where broadleaf weed control or suppression is desired, tank mix this product with an appropriate, labeled broadleaf weed herbicide.

#### Conservation Reserve Program (CRP Acres)

This product can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres. For specific application rates for various annual and perennial weeds, see the *Weeds Controlled* section of this label. CPR applications may be made with wiper applicators or conventional spray equipment. For selective applications with broadcast spray equipment, apply 12 to 16 ounces per acre of this product in early spring before desirable CPR grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if applications are made when plants are not dormant.

#### Habitat Management

This product is recommended for the restoration and/or maintenance of native habitats and in wildlife management areas. Apply as recommended in the *Noncrop Uses* Section of this label.

**Habitat Restoration and Maintenance:** When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

**Wildlife Food Plots:** This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling.

#### Ornamentals and Christmas Trees

**This product is not recommended for use as an over-the-top broadcast spray in ornamentals and Christmas trees.** **Note:** Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

When applied as instructed for the conditions described for *Noncrop Uses* this product controls undesirable vegetation listed on this label prior to planting within and around greenhouses and shadehouses, and as a postdirected spray around established ornamentals and Christmas trees. For specific application rates and instructions for control of various annual and perennial weeds, see the *Weeds Controlled* section of this label. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

**Site Preparation:** Following preplant applications of this product, any ornamental, nursery species or Christmas tree species may be replanted. Precautions should be taken to protect nontarget plants during site preparation applications.

**Greenhouse/Shadehouse Use:** This product may be used to control weeds listed on this label that are growing in greenhouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

**Postdirected Spray:** Use as a postdirected spray around established woody ornamental species, nursery species or Christmas trees such as those listed below. Care must be exercised to avoid contact of spray, drift or mist with foliage of or green bark of established ornamental species.

- |  |                       |
|--|-----------------------|
| Arborvitae                                 | Lilac                 |
| <i>Thuja</i> spp.                          | <i>Syringa</i> spp.   |
| Azalea                                     | Magnolia              |
| <i>Rhododendron</i> spp.                   | <i>Magnolia</i> spp.  |
| Boxwood                                    | Maple                 |
| <i>Buxus</i> spp.                          | <i>Acer</i> spp.      |
| Crabapple                                  | Oak                   |
| <i>Malus</i> spp.                          | <i>Quercus</i> spp.   |
| Euonymus                                   | Privet                |
| <i>Euonymus</i> spp.                       | <i>Ligustrum</i> spp. |
| Fir  | Pine                  |
| <i>Abies</i> spp., <i>Pseudotsuga</i> spp. | <i>Pinus</i> spp.     |
| Jobba                                      | Spruce                |
| <i>Simmondsia chinensis</i>                | <i>Picea</i> spp.     |
| Hollies                                    | Yew                   |
| <i>Ilex</i> spp.                           | <i>Taxus</i> spp.     |

**Silvicultural Sites and Rights-of-Way**

**Note: Not recommended for use as an over-the-top broadcast spray in silvicultural nurseries.**

When applied as directed for **Noncrop Uses** under conditions described, this product controls undesirable vegetation listed on this label. This product also suppresses or controls undesirable vegetation listed on this label when applied at recommended rates for release of established coniferous species listed on this label. For specific application rates and instructions for control of various brush, annual and perennial weeds, see the **Weeds Controlled** section of this label. For specific application rates for release of listed coniferous species, see the **Conifer Release** part of this section of the label. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

**Aerial Application:** This product may be applied using aerial spray equipment for silvicultural site preparation, conifer release and rights-of-way treatments. See the **Application Equipment and Techniques** part of the **Mixing, Additives and Application Instructions** section of this label for information on how to apply this product by air. **Do not apply this product by air to rights-of-way in the state of California.**

**Site Preparation**

Any silvicultural species may be planted following preplant applications of this product.

**Postdirected Spray**

In established silvicultural sites, use as a spray on the foliage of undesirable vegetation. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

**Conifer Release**

For release, apply only where conifers have been established for more than one year. Do not disturb vegetation prior to treatment or until visual symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late fall. Injury may occur to conifers treated for release, especially where spray patterns overlap, the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.**

Applications must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the **Weeds Controlled** section of this label.

For release of the following conifer species:

- |                              |                   |
|------------------------------|-------------------|
| Douglas fir                  | Pines*            |
| <i>Pseudotsuga menziesii</i> | <i>Pinus</i> spp. |
| Fir                          | Spruce            |

*Abies* spp.  
Hemlock  
*Tsuga* spp.

*Picea* spp.

\*Includes all species except eastern white pine, loblolly pine or slash pine.

Apply 1.5 to 2 quarts of this product per acre except in Washington and Oregon, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of this product per acre before conifer bud swell for control of annual weeds. For fall treatment in Washington and Oregon, west of the crest of the Cascade mountains, apply 1 to 1.5 quarts of this product per acre before any major leaf drop of deciduous species.

For release of western hemlock, apply 1 quart of this product per acre.

For release of the following conifer species:

Loblolly pine  
*Pinus taeda*

Eastern white pine  
*Pinus strobus*

Slash pine  
*Pinus elliottii*

**Late Season Application:** Apply 1.5 to 2 quarts of this product in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will create the potential for increased injury in the form of tip and/or needle burn. Injury may decrease with later applications. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants. Applications made according to label directions will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash  
*Fraxinus* spp.  
Cherry, black  
*Prunus serotina*  
Cherry, Pin  
*Prunus pensylvanica*  
Elm  
*Ulmus* spp.  
Hawthorn  
*Crataegus* spp.  
Locust, black  
*Robina pseudoacacia*  
Maple, red  
*Acer rubra*  
Oak, black  
*Quercus velutina*  
Oak, post  
*Quercus stellata*  
Oak, southern red  
*Quercus falcata*

Oak, white  
*Quercus alba*  
Persimmon  
*Diospyros* spp.  
Poplar, yellow  
*Liriodendron tulipifera*  
Sassafras  
*Sassafras albidum*  
Sourwood  
*Oxydendrum arboreum*  
Sumac, poison  
*Rhus vernix*  
Sumac, smooth  
*Rhus glabra*  
Sumac, winged  
*Rhus copallina*  
Sweetgum  
*Liquidambar styraciflua*

Apply only to those sites where woody brush and trees listed in this label constitute the majority of the undesirable species.

#### **This Product Plus Oust Tank Mixtures for Conifer Release from Herbaceous Weeds**

To release loblolly pines from herbaceous weeds, tank mixtures of this product with Oust will provide control of annual weeds listed in the **Weeds Controlled** section of this and the Oust label, and partial control of the perennial weeds listed below.

Apply 16 to 24 fluid ounces of this product with 2 to 4 ounces of Oust in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top of the young loblolly pines. This product plus Oust tank mixtures may not be applied by air in California. This tank mixture may be applied using aerial equipment. When applying by air, use the recommended rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inch height (or runner length on annual vines), use the lower rates of both products. Use the higher rates of both products when annual weeds are in more advanced stages of growth and

approaching flower or seed formation. Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

- |  |   |
|--|---|
| Bahiagrass<br><i>Paspalum notatum</i>        | Johnsongrass*<br><i>Sorghum halepense</i>   |
| Broomsedge<br><i>Andropogon virginicus</i>   | Poorjoe*<br><i>Diodia teres</i>             |
| Dock, curly<br><i>Rumex crispus</i>          | Trumpetcreeper**<br><i>Campsis radicans</i> |
| Dogfennel<br><i>Eupatorium capilliflorum</i> | Vaseygrass<br><i>Paspalum urville</i>       |
| Fescue, tall<br><i>Festuca arundinacea</i>   | Vervain, blue<br><i>Verbena hastata</i>     |

\*Control only at the higher rates.  
\*\*Suppression only at the higher rates.

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought-flood water, insects or disease. Read and observe the cautionary statements and all other information appearing on the labels of all herbicides used.

**Note to User:** This product must not be used in areas where adverse impact on federally designated endangered/threatened plant or aquatic species is likely. Prior to making applications, the user of this product must determine that no such species are located in or immediately adjacent to the area to be treated.

### Cut Stump Treatments

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, apply during periods of active growth and full leaf expansion. When used according to directions for cut stump application, this product will **control, partially control or suppress** many types of woody brush and tree species, some of which are listed below:

- |                                      |  |
|--------------------------------------|--|
| Alder<br><i>Alnus spp.</i>           | Saltcedar<br><i>Tamarisk spp.</i>          |
| Eucalyptus<br><i>Eucalyptus spp.</i> | Sweetgum<br><i>Liquidambar styraciflua</i> |
| Madrone<br><i>Arbutus menziesii</i>  | Tan Oak<br><i>Lithocarpus densiflorus</i>  |
| Oak<br><i>Quercus spp.</i>           | Willow<br><i>Salix spp.</i>                |
| Reed, giant<br><i>Arundo donax</i>   |  |

### Injection and Frill Applications

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 1 ml of this product per each 2 to 3 inches of trunk diameter (DBH). This is best achieved by applying a 50 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as this, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, apply during periods of active growth and after full leaf expansion. This treatment will **control** the following woody species:

- |                               |  |
|-------------------------------|--|
| Oak<br><i>Quercus spp.</i>    | Sweetgum<br><i>Liquidambar styraciflua</i> |
| Poplar<br><i>Populus spp.</i> | Sycamore<br><i>Platanus occidentalis</i>   |

This treatment will **suppress** the following woody species:

320946

Black gum  
*Nyssa sylvatica*  
Dogwood  
*Cornus* spp.

Hickory  
*Carya* spp.  
Maple, red  
*Acer rubrum*

**Turfgrasses and Grasses for Seed Production  
Preplant and Renovation**

When applied as directed for **Noncrop Uses** under conditions described, this product controls most existing vegetation prior to the planting or renovation of either turfgrasses or grass seed production areas. For specific application rates and instructions for control of various annual and perennial weeds, and woody brush and trees, see the **Weeds Controlled** section of this label. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control. **Do not disturb soil or underground plant parts before treatment.** Delay tillage or renovation techniques such as vertical mowing, coring or slicing for 7 days after application to allow proper translocation into underground plant parts.

**Turfgrasses**

Where existing vegetation is growing in a field or unmowed situation, apply this product to actively growing weeds at the stages of growth listed in the **Weeds Controlled** section of this label. Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray. Desirable turfgrasses may be planted following the above procedures.

**Grasses for Seed Production**

Apply this product to actively growing weeds at the stages of growth recommended in the **Weeds Controlled** section of this label prior to planting or renovation of turf or forage grass areas grown for seed production. **Do not feed or graze** treated areas within 8 weeks after application.

**Annual Weed Control in Dormant Bermudagrass and Bahiagrass Turf**

When applied as directed for **Noncrop Uses** under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release or dormant bermudagrass and bahiagrass turf. Refer to the rate table for this product alone under the **Release of Bermudagrass and Bahiagrass** section of this label for recommended rates and volumes on the species to be suppressed or controlled. Treat only when turf is dormant and prior to spring greenup. Spot treatments or broadcast applications of this product in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained turfgrass areas, i.e., golf courses, lawns, etc. **Do not apply tank mixtures** of this product plus Oust in highly maintained turfgrass areas.

**Release of Bermudagrass or Bahiagrass**

**Note:** Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. Use tank mixtures of this product plus Oust only on railroads, highways, utility plant sites or other right-of-way areas.

When applied as directed for **Noncrop Uses** under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. This product may be tank-mixed with Oust as recommended for residual control. Make applications to dormant bermudagrass or bahiagrass. Tank mixtures of this product plus Oust may delay greenup. To avoid delays in greenup and minimize injury, do not add more than 1 ounce per acre of Oust on bermudagrass or more than 0.5 ounce per acre on bahiagrass, or treat when these grasses are in a semi-dormant condition. For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4- to 6-leaf stage.

**Weeds Controlled**

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below. Apply the recommended rates of this product alone or as a tank mixture in 10 to 25 gallons of water, plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre. For the best recommendation for the mixture of weeds within your geographic area, contact your Marman USA sales representative.

**Weeds Controlled or Suppressed with This Product Alone\***

Note: C = Control S = Suppression

Weed Species	This Product Fluid Oz./Acre					
	8	12	16	24	32	64

SIC  
T



Barley, little ( <i>Hordeum pusilium</i> )	S	C	C	C	C	C
Bedstraw, catchweed ( <i>Galium aparine</i> )	S	C	C	C	C	C
Bluegrass, annual ( <i>Poa annua</i> )	S	C	C	C	C	C
Chervil ( <i>Chaerophyllum tainturieri</i> )	S	C	C	C	C	C
Chickweed, common ( <i>Stellaria media</i> )	S	C	C	C	C	C
Clover, crimson ( <i>Trifolium incarnatum</i> )	•	S	S	C	C	C
Clover, large hop ( <i>Trifolium campestre</i> )	•	S	S	C	C	C
Fescue, tall ( <i>Festuca arundinaceae</i> )	•	•	•	•	S	S
Geranium, Carolina ( <i>Geranium carolinianum</i> )	•	•	S	S	C	C
Henbit ( <i>Lamium amplexicaule</i> )	•	S	C	C	C	C
Ryegrass, Italian ( <i>Lolium multiflorum</i> )	•	•	S	C	C	C
Speedwell, corn ( <i>Veronica arvensis</i> )	S	C	C	C	C	C
Vetch, common ( <i>Vicia sativa</i> )	•	•	S	C	C	C

\*These rates apply only to sites where an established competitive turf is present.

**Weeds Controlled or Suppressed with This Product Plus Oust\***

Note: C = Control S = Suppression

Weed Species	This Product (Fl. oz. / A)	This Product + Oust						
	Oust (Oz. / A)	8 +	12 +	12 1/2	16 +	16 1/2	12 +	16 +
		1/4	1/4	1/2	1/4	1/2	1	1
Barley, little ( <i>Hordeum pusilium</i> )		C	C	C	C	C	C	C
Bedstraw, catchweed ( <i>Galium aparine</i> )		C	C	C	C	C	C	C
Bluegrass, annual ( <i>Poa annua</i> )		S	C	C	C	C	C	C
Chervil ( <i>Chaerophyllum taintuieri</i> )		C	C	C	C	C	C	C
Chickweed, common ( <i>Stellaria media</i> )		S	C	C	C	C	C	C
Clover, crimson ( <i>Trifolium incarnatum</i> )		S	S	S	S	C	C	C
Clover, large hop ( <i>Trifolium campestre</i> )		•	•	S	S	S	C	C
Fescue, tall ( <i>Festuca arundinaceae</i> )		•	•	•	•	•	S	S
Geranium, Carolina ( <i>Geranium carolinianum</i> )		•	S	S	C	C	C	C
Henbit ( <i>Lamium amplexicaule</i> )		•	S	C	C	C	C	C
Ryegrass, Italian ( <i>Lolium multiflorum</i> )		•	S	S	C	C	C	C
Speedwell, corn ( <i>Veronica arvensis</i> )		S	C	C	C	C	C	C
Vetch, common ( <i>Vicia sativa</i> )		C	C	C	C	C	C	C

\*These rates or mixtures of rates apply only to sites where an established competitive turf is present.

**Release of Actively Growing Bermudagrass**

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the **Weeds Controlled** section of this and the Oust label, and suppression or partial control of certain perennial weeds. For control or suppression of those annual species listed on this label, use 1 to 3 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. Use the higher rate of this product for partial control of the following perennial species. Use the lower rates for suppression of growth. For best results, see the **Weeds Controlled** section of this label for proper stage of growth.

- |  |   |
|--|---|
| Bahiagrass<br><i>Paspalum notatum</i>              | Johnsongrass*<br><i>Sorghum halepense</i>   |
| Bluestem, silver<br><i>Andropogon saccharoides</i> | Trumpetcreeper**<br><i>Campsis radicans</i> |
| Fescue, tall<br><i>Festuca arundinacea</i>         | Vaseygrass<br><i>Paspalum urvillei</i>      |

\*Control at the higher rates.  
\*\*Suppression only at the higher rates.

This product may be tank-mixed with Oust. If tank-mixed, use no more than 1 to 2 pints per acre of this product with 1 to 2 ounces of Oust per acre. Use the lower rates of both mixtures to control annual weeds below 6 inches in height (or runner length in annual vines) that are listed in the **Weeds Controlled** section of this booklet and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. Use the higher rates of this product to provide partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass <i>Paspalum notatum</i>	Johnsongrass* <i>Sorghum halepense</i>
Bluestem, silver <i>Andropogon saccharoides</i>	Poorjoe** <i>Diodia teres</i>
Broomsedge <i>Andropogon virginicus</i>	Trumpetcreeper* <i>Campsis radicans</i>
Dock, curly <i>Rumex crispus</i>	Vaseygrass <i>Paspalum urvillei</i>
Dogfennel <i>Eupatorium capilliflorum</i>	Vervain, blue <i>Verbena hastata</i>
Fescue, tall <i>Festuca arundinacea</i>	

\*Suppression only at higher rates.  
\*\*Control at the higher rates.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

### Cool Season Turf Growth Regulation

When applied as directed, this product will suppress growth and seedhead development of listed turf species in industrial sites. This product is recommended for management of coarse turf on roadside rights-of-way or other industrial areas. Do not use on high-quality turf or other areas where some turf color changes cannot be tolerated. Slight turf discoloration may occur but turf will regreen and regrow under moist conditions as effects of this product wear off. Apply 4 to 6 fluid ounces of this product per acre alone or in a recommended tank mixture. Spray volumes of 10 to 40 gallons per acre are recommended. When using this product, mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

This product can be used for growth and seedhead suppression of **Tall Fescue** and **Smooth Brome**. For best results, apply this product in a recommended tank mixture to actively growing turfgrasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury. After mowing or removal of seedheads, this product in a recommended tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

**Annual Grasses:** For growth suppression of some annual grasses such as annual ryegrass, wild barley and wild oats, apply 3 to 4 fluid ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

### Tank Mixtures

For the following tank mixtures, consult each product label for weeds controlled and the correct stage of application. Do not treat turf under stress.

**Tank Mixtures plus 2,4-D Amine:** For additional weed control benefits, up to 1 pound a.i. per acre of 2,4-D amine may be added to the following tank mixtures. Consult the 2,4-D amine label for weeds controlled.

### Tall Fescue:

**This Product plus Telar™:** For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.5 ounce of Telar per acre. This tank mixture can also be applied after mowing or removal of tall fescue seedheads to turf growth suppression. Make only one of the above applications per growing season.

**This Product plus Oust:** For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-in-stage of development. Use up to 0.25 ounce of Oust per acre.

**This Product plus Escort™:** This tank mixture can be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control or partial control of some annual weeds. Use up to 1/3 ounce of Escort per acre.

**Smooth Brome:**

**This Product plus Oust:** For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of Oust per acre.

### Bahiagrass Seedhead and Vegetative Suppression

When applied as directed in the indicated noncrop areas (roadsides, airports, golf course roughs and plant sites), this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications. Apply this product 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces per acre of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 4 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

A tank mixture of this product plus Oust **may be applied only on roadsides** for seedhead inhibition and vegetative suppression. Apply 6 fluid ounces per acre of this product plus 0.25 ounce per acre of Oust, plus 0.5 to 1 percent nonionic surfactant by total spray volume 1 to 2 weeks following an initial spring mowing. When using this product plus Oust for suppression of bahiagrass, make only 1 application per year.

### Cropping Systems

When applied as directed for **Cropping Systems**, under the conditions described, this product controls annual and perennial weeds listed on this label, prior to the emergence of direct seeded crops or prior to transplanting of crops listed on this label. See **General Information** and **Mixing, Additives and Application Instructions** sections of this label for essential product performance information. See the following **Cropping Systems** sections for specific recommended uses.

**Extreme care must be exercised to avoid contact of spray with foliage, green stems or fruit of desirable crops, plants, trees or other desirable vegetation since severe damage or destruction may result.** Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Except as otherwise specified on this label, repeat treatments must be made before the crop emerges in accordance with the instructions of this label. Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts per acre of this product per year.

The maximum use rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate- or sulfosate-containing products does not exceed the stated maximum use rate.

For any crop not listed below, applications must be made at least 30 days prior to planting.

Do not harvest or feed treated vegetation for 8 weeks following application. Following spot treatment or selective equipment use, allow 14 days before grazing domestic livestock or harvesting forage grasses and legumes.

**Row Crops:** \*Corn (all), \*Cotton, Peanuts, \*Sorghum (Milo), \*Soybeans, \*Sugarcane

**Cereal Grains:** \*Barley, \*Buckwheat, \*Millet (Pearl, Proso), \*Oats, \*\*Rice, \*Rye, \*Triticale, \*Wheat (all), \*Wild Rice

**Citrus:** Calamondin, Chironja, Citron, Grapefruit, Kumquat, Lemon, Lime, Mandarin Orange, Orange (all), Pummelo, Tangelo, Tangerine, Tangors

**Tree Nuts:** Almond, Beechnut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (Black, English)

**Vine Crops:** Grapes, Kiwi Fruit

**Tree Fruits:** Apple, Apricots, Cherry (Sweet, Sour), Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince

**Vegetables:** Artichoke (Jerusalem), \*Asparagus, Beans (all), Beet Greens, Beets (Red, Sugar, Broccoli (all), Brussels Sprouts, Cabbage (all), Cabbage (Chinese), \*\*\*Cantaloupe, Carrot, Cauliflower, \*\*\*Casaba Melon, Celery, Chard (Swiss), Chicory, Collards, \*\*\*Crenshaw Melon, \*\*\*Cucumber, \*\*\*Eggplant, Endive, \*\*\*Garlic, \*\*\*Gourds, \*\*\*Ground Cherry, \*\*\*Honeydew Melon, \*\*\*Honey Ball Melon, Horseradish, Kale, Kohlrabi, Leek, Lentils, Lettuce, \*\*\*Mango Melon, \*\*\*Melons (all), \*\*\*Muskmelon, Mustard Greens, Okra, Onion, Parsley, Parsnips, Peas (all), \*\*\*Pepper (all), \*\*\*Persian Melon, Potato (Irish, Sweet), \*\*\*Pumpkin, Radish, Rape Greens, Rhubarb, Rutabaga, Shallot, Spinach (all), \*\*\*Squash (Summer, Winter), \*\*\*Tomatillo, \*\*\*+Tomato, Turnip, \*\*\*Watercress, \*\*\*Watermelon, Yams

**Small Fruits and Berries:** Blackberry, Blueberry, Boysenberry, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Huckleberry, Loganberry, Olallieberry, Raspberry (Black, Red)

**Forage Crops and Legumes:** \*Alfalfa, \*Forage Grasses, \*Forage Legumes

**Tropical Crops:** Acerola, Atemoya, Avocado, Banana, Breadfruit, Canistel, Carambola, Cherimoya, Cocoa Beans, Coffee, Dates, Figs, Genip, Guava, Jaboticaba, Jackfruit, Longan, Lychee, Mango, Papaya, Passion Fruit, Persimmons, \*\*\*\*Pineapple, Plantains, Pomegranate, Sapodilla, Sapote (Black, Mamey, White), Soursop, Sugar Apple, Tamarind, Tea

- \*Spot treatments may be applied in these crops.
- \*\*Do not treat rice fields or levees when the fields contain flood water.
- \*\*\*Apply only prior to planting. Allow at least 3 days between application and planting.
- \*\*\*\*Do not feed or graze treated pineapple forage following application.
- +Use is restricted only to direct seeded crops.

When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove residues of this product, that could cause crop injury, from the plastic prior to transplanting. Residues can be removed by a single 0.5-inch application of water either by natural rainfall or via a sprinkler irrigation system. Applications made at emergence will result in injury or death to emerged seedlings.

**Spot Treatment** (Only those crops with "\*" can be spot treated): Applications in growing crops must be made prior to heading of small grains and milo, initial pod set in soybeans, silking of corn or boll opening on cotton.

For forage grasses and forage legumes see **Spot Treatment** in the **Pastures** section of **Cropping Systems** in this label. For dilution and application rates using boom or hand-held equipment, see **Mixing, Additives and Application Instructions** and **Weeds Controlled** sections of this label.

**Note:** For forage grasses and forage legumes, no more than one-tenth of any acre should be treated at one time. For all other crops, do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

**Selective Equipment:** This product may be applied through recirculating sprayers, shielded applicators or wiper applicators in cotton and soybeans. Shielded and wiper applicators may also be used in tree crops and grapes. Wiper applicators may be used in wheat, rutabagas, forage grasses and forage legumes, including pasture sites and grain sorghum (milo). See the **Selective Equipment** part of the **Application Equipment and Techniques** section of this label for information on proper use and calibration of this equipment.

Allow at least the following time intervals between application and harvest:

Cotton, Soybeans	7 days
Apples, Citrus, Pear	1 day
Atemoya, Avocado, Breadfruit, Canistel, Carambola, Cherry, Grapes, Dates, Jaboticaba, Jackfruit, Logan, Lychee, Passion Fruit, Persimmons, Rutabagas, Sapodilla, Sapote, Soursop, Sugar Apple, Tamarind	14 days
Stone Fruit	17 days
Nut Crops	3 days
Wheat <sup>1</sup>	35 days
Sorghum (milo) <sup>1,2</sup>	40 days

<sup>1</sup>Do not use roller applicators.  
<sup>2</sup>Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

**Asparagus**

When applied as directed for **Cropping Systems** under the conditions described, this product controls weeds listed on this label in asparagus. For specific application rates and instructions for control of various annual and perennial weeds, see the **Weeds Controlled** section of this label.

**Prior to Crop Emergence:** Apply this product prior to crop emergence for the control of emerged labeled annual and perennial weeds. **Do not apply within a week before the first spears emerge.**

**Spot Treatment:** Apply this product immediately after cutting, but prior to the emergence of new spears. Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

**Postharvest:** Apply this product after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatment should be applied as directed or shielded spray to avoid contact of the spray with ferns, stems or spears. Direct contact of the spray with the asparagus may result in serious crop injury.

**Note:** Select and use recommended types of spray equipment for postemergence postharvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

### Berries and Small Fruits

Wiper applicators may be used in cranberries in accordance with instructions in this section. For other berries, apply as a preplant broadcast application, or as a directed spray or wiper application post-planting. See **General Information** and **Mixing, Additives and Application Instructions** sections of this label for essential product performance information. See the **Selective Equipment** part of the **Application Equipment and Techniques** section of this label for information on recommended use and calibration of this equipment. Allow a minimum of 30 days between last application and harvest of cranberries. For other small fruits and berries, allow a minimum of 14 days between last application and harvest.

**For Wick or other Wiper Applicators:** Mix 1 gallon of this product in 4 gallons of water to prepare a 20 percent solution. In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this product are wiped on the weeds. A second treatment in the opposite direction may be beneficial.

Do not permit herbicide solution to contact desirable vegetation, including green shoots, canes or foliage.

### Corn

**Hooded Sprayers:** This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bounding or raising the hoods of the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground. When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 1 quart of this product per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Leave at least an 8-inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 mph.
- Maximum wind speed: 10 mph.
- Use low-drift nozzles

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction. Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. For specific application rates and instructions for control of various annual and perennial weeds, see the **Weeds Controlled** section of this label. Do not graze or feed corn forage or fodder following application of this product through hooded sprayers. Do not apply more than 3 quarts of this product per acre per year for hooded sprayer applications.

### Fallow and Reduced Tillage Systems

Use this product in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the **Weeds Controlled** Section of this label for specific rates and instructions. This product may be

applied using ground or aerial spray equipment. See the **Application Equipment and Techniques** section of this label for instructions.

**Tank Mixtures**  
**This Product plus Banvel plus Nonionic Surfactant**  
**This Product plus 2,4-D plus Nonionic Surfactant**

**Do not apply Banvel or 2,4-D tank mixtures by air in California.** Applications of 2,4-D or Banvel must be made at least 7 days prior to planting corn. The addition of Banvel in a mixture with this product may provide short-term residual control of selected weed species. Some crop injury may occur if Banvel is applied within 45 days of planting. Refer to the Banvel and 2,4-D labels for cropping restrictions and other use instructions.

**This Product plus Goal plus Nonionic Surfactant**

This product alone or in tank mixtures with Goal plus 0.5 to 1 percent nonionic surfactant by total spray volume will provide control of those weeds listed below. Make applications when weeds are actively growing and at the recommended stages of growth. Avoid spraying when weeds are subject to moisture stress, when dust is on the foliage or when straw canopy covers the weeds.

This Product 12 fluid oz./A		This Product 16 fluid oz./A	
Wheat	18"*	Annual grasses at left plus:	
Barley	12"	Ryegrass, annual	6"
Bluegrass, annual	6"	Chickweed	6"
Barnyardgrass	6"	Groundsel	6"
Rye	6"	Marestail	6"
*Maximum height or length in inches		Rocket, London	6"
		Shepherdspurse	6"
		Crabgrass	12"
		Johnsongrass, seedling	12"
		Lambsquarters	12"
		Oats, wild	12"
		Pigweed, redroot	12"
		Mustards	12"

**Note:** Use 32 fluid ounces of this product per acre where heavy weed densities exist.

This Product 12 fluid oz./A + Goal** 2 to 4 fluid oz./A		This Product 16 fluid oz./A + Goal** 2 to 4 fluid oz./A	
Annual grasses above plus:		Annual weeds above plus:	
Cheeseweed, common	3"	Cheeseweed, common	6"
Chickweed	3"	Groundsel	6"
Groundsel	3"	Chickweed	12"
Rocket, London	6"	Rocket, London	12"
Shepherdspurse	6"	Shepherdspurse	12"

**Note:** Use 32 fluid ounces of this product per acre in mixtures with 2 to 4 fluid ounces of Goal per acre where heavy weed densities exist.

\*Maximum height or length in inches.

\*\*Use the higher rate of Goal when weeds approach maximum recommended height or stands are dense.

These recommended tank mixtures may be applied using ground or aerial spray equipment. Refer to the **Weeds Controlled** section of this label for specific rates and instructions.

**Ecofarming Systems**

The recommendations made in this section are not registered for use in California. The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow. Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

- This Product at 16 to 20 fluid ounces per acre**
- Plus**
- 2,4-D at 0.375 to 0.5 pound a.i. per acre**
- plus**
- Atrazine at 0.75 to 1 pound a.i. per acre**
- Plus**
- Lasso® at 2.5 to 3 quarts per acre**

Apply the above tank mixture in a 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the desired carrier volume.

**Weeds Controlled:** The following weeds, up to a maximum height of 4 inches, will be controlled:

Brome, downy <i>Bromus tectorum</i>	Lettuce, prickly <i>Lactuca serriola</i>
Cheat <i>Bromus secalinus</i>	Pigweed, redroot <i>Amaranthus retroflexus</i>
Foxtail, green <i>Setaria viridis</i>	Thistle, Russian <i>Salsola kali</i>
Foxtail, yellow <i>Setaria letuscens</i>	Wheat, volunteer <i>Triticum aestivum</i>
Kochia* <i>Kochia scoparia</i>	

\*For improved control of kochia, add 4 fluid ounces per acre (0.125 pound a.i. per acre) of Banvel to the above tank mixture.

Risk of crop injury from 2,4-D or Banvel can be reduced by applying this treatment 7 to 14 days before planting. Refer to the label booklet for Lasso herbicide for preemergence weed control achieved by this tank mixture. Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products used in these tank mixtures.

#### Aid to Tillage

This product, when used in conjunction with preplant tillage practices, will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 8 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Make applications when weeds are actively growing and before they are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage. Tank mixtures with residual herbicides may result in reduced performance.

#### Postharvest Grain Sorghum, Sorghum Regrowth Control

This product may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of this product per acre for control, or 1.5 pints of this product per acre for suppression. Use 0.5 percent nonionic surfactant in 3 to 10 gallons of spray solution per acre.

#### Pastures

Apply this product prior to planting forage grasses and legumes.

**Pasture or Hay Crop Renovation:** When applied as a broadcast spray, this product controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes. Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

**Spot Treatment:** When applied as a spot treatment as recommended, this product controls annual and perennial weeds listed in this label that are growing in pastures, forage grasses and forage legumes composed of bahiagrass, bermudagrass, bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa or clover.

**Wiper Application:** When applied as directed, this product controls or suppresses the weeds listed under **Wiper Applicators** and the **Selective Equipment** section of this label.

For spot treatment and wiper application, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Further applications may be made in the same area at 30-day intervals. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

#### Sugarcane

When applied as directed for **Cropping Systems**, under the conditions described, this product controls those emerged annual and perennial weeds listed on this label growing in or around sugarcane or in fields prior to the emergence of plant cane. This product will also control undesirable sugarcane. **Note:** Where repeat treatments are necessary, do not exceed a total of 10.6 quarts of this product per acre per year. Do not apply to vegetation on or around ditches, canals or ponds containing water to be used for irrigation.

**Broadcast Treatment:** Apply this product in 10 to 40 gallons of water per acre on emerged weeds prior to the emergence of plant cane. For specific application rates and instructions for control of various annual and perennial weeds, see the **Weeds Controlled** section of this label.

For removal of last stubble or ratoon cane, apply 4 to 5 quarts of this product in a 10 to 40 gallons of water per acre to new growth having at least 7 or more new leaves. Allow 7 or more days after application before tillage.

**Spot Treatment in or Around Sugarcane Fields:** For dilution and application rates using hand-held equipment, see **Mixing, Additives and Application Instructions** and **Weeds Controlled** sections of this label.

For control of volunteer or diseased sugarcane, make a 1 percent solution of this product in water and spray to wet the foliage of vegetation to be controlled. **Note:** When spraying volunteer or diseased sugarcane, the plants should have at least 7 new leaves.

Avoid spray contact with healthy can plants since severe damage or destruction may result. Do not feed or graze treated sugarcane forage following application.

## Conservation Tillage, Minimum Tillage and No-Till Systems

### Corn and Soybeans

#### Tank Mixtures

The recommendations made in this section are not registered for use in California.

When applied as recommended under the conditions described, the tank mixtures listed in this section control many emerged weeds, and give preemergence control of many annual weeds where corn or soybeans will be planted directly into a cover crop, established sod or in previous crop residues. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures. For mixing instructions, see the **Mixing, Additives and Application Instructions** section of this label.

Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre before, during or after planting. Do not apply these mixtures after crop emergence. When tank mixing with residual herbicides, add an agriculturally-approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution. The addition of 1 to 2 percent dry ammonium sulfate by weight may increase the performance of this product. **Note:** When using these tank mixtures, do not exceed 4 quarts of this product per acre.

#### Corn

For residual control, this product may be tank-mixed with the following herbicides or combination of herbicides:

Lasso/Alachlor	Atrazine
Lariat <sup>®</sup>	Cyanazine
Bullet <sup>®</sup>	Simazine
Dual Magnum <sup>™</sup>	Prowl <sup>™</sup>
Bicep Magnum <sup>™</sup>	Micro-Tech <sup>®</sup>
Partner <sup>®</sup>	

For improved burndown, this product may be tank-mixed with 2,4-D or dicamba. Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn. See the **Weeds Controlled** section for specific rate information.

#### Soybeans

For residual control, this product may be tank-mixed with the following herbicides or combination of herbicides:

Canopy <sup>™</sup>	Lorox <sup>™</sup> Plus
Command <sup>™</sup>	Preview <sup>™</sup>
Dual Magnum	Prowl
Gemini <sup>™</sup>	Turbo <sup>™</sup>
Lasso/Alachlor	Scepter <sup>™</sup>
Lexone <sup>™</sup>	Sencor <sup>™</sup>
Linuron	Squadron <sup>™</sup>
Pursuit <sup>™</sup>	Pursuit Plus <sup>™</sup>
Partner	Micro-Tech

For improved burndown, this product may be tank-mixed with the following herbicides:

2,4-DB

2,4-D\*

\*See the label for 2,4-D for intervals between application and planting.

#### Corn and Soybeans

**Annual Weeds:** For difficult-to-control weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply this product at 2 pints per acre in these tank mixtures. For other listed annual weeds, apply 1 to 1.5 pints of this product per acre when weeds are less



than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. For a complete list of annual weeds controlled, see the **Weeds Controlled** section of this label.

**Perennial Weeds:** At normal application times in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the **Weeds Controlled** section of this label for proper stage of growth for perennial weeds.

Use of 2 to 4 quarts of this product per acre in the tank mixtures mentioned above, under these conditions, provides top kill and reduces competition from many emerged perennial grass and broadleaf weeds. For emerged perennial weeds controlled, see **Weeds Controlled** section of this label. To obtain the desired stage of growth, it may be necessary to apply this product alone in the late summer or fall and then follow with a label-approved, seedling weed control program at planting.

**Use of these tank mixtures for bermudagrass or Johnsongrass control in minimum tillage systems is not recommended.** For bermudagrass control, follow the instructions under **Control of Perennial Weeds** section of this label and then use a label-approved, seedling weed-control program in a minimum tillage or conventional tillage system. For Johnsongrass control, follow instructions under **Control of Perennial Weeds** section of this label, and then use a label-approved, seedling weed-control program with conventional tillage.

### Preharvest Applications

When applied as directed under the conditions described, this product controls annual and perennial weeds listed on this label prior to the harvest of cotton, soybeans, grain sorghum (milo) and wheat. For specific rates and application instructions for control of various annual and perennial weeds see the **Weeds Controlled** section of this label. This product may be applied by both ground and aerial application equipment. **Do not apply more than 1 quart per acre of this product by air.** See the **Application Equipment and Techniques** section of this label for instructions for ground and aerial applications. **Note:** Do not apply to crops grown for seed. Reduction in germination or vigor may occur. The use of this product for preharvest grain sorghum (milo) is not registered in California.

#### Soybeans

Apply after pods have set and lost all green color. Allow a minimum of 7 days between application and harvest of soybeans. Avoid excessive seed shatter loss due to ground application equipment. Do not graze or harvest treated crop for livestock feed within 25 days of last preharvest application. **Do not apply more than 6 quarts per acre of this product for preharvest applications.**

#### Cotton

**Broadcast Applications:** This product may be applied using either aerial or ground spray equipment. For ground applications with broadcast equipment, apply this product in 10 to 20 gallons of water per acre. For aerial applications, apply this product in 3 to 10 gallons of water per acre. This product provides weed control and cotton regrowth inhibition when applied prior to the harvest of cotton. Apply 1 to 2 quarts of this product in 3 to 10 gallons of water per acre for cotton regrowth inhibition. Do not apply more than 2 quarts of this product per acre for preharvest applications. **The use of additives for preharvest application to cotton is prohibited.**

This product may be tank mixed with DEF™ 6, Folex™ or Prep™ to provide additional enhancement of cotton leaf drop. Allow a minimum of 7 days between application and harvest of cotton. Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential. Do not feed or graze treated cotton forage or hay following preharvest applications.

#### Grain Sorghum (Milo)

Make applications at 30 percent grain moisture or less and at least 7 days prior to harvest.

#### Wheat

Apply after the hard-dough stage of grain (30 percent or less grain moisture) and at least 7 days prior to harvest. **Do not apply more than 1 quart per acre of this product for preharvest applications to wheat.**

#### Tree and Vine Crops

This product is recommended for weed control in established groves, vineyards and orchards, or for site preparation prior to transplanting crops listed in this section. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high-volume wands, lances, orchard guns, or with wiper applicator equipment, except as directed in this section. See the **Application Equipment and Techniques** section of this label for specific information on use of equipment. When applying this product, refer to the **Weeds Controlled** section of this label and to specific recommendations in this section for rates to be used.

**Note:** Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual weed control. For subsequent weed control, use repeated applications of this product. Do not apply more than 10.6 quarts of this product per acre per year.

**Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of tress or vines. Contact of this product with other than matured brown bark can result in serious crop damage. Avoid painting out stumps with this product as injury resulting from root grafting may occur in adjacent trees.**

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment. For specific application rates and instructions, see the **Weeds Controlled** section of this label and to specific recommendations which follow.

**Middles Management**

(For Annual Weeds in Middles between Rows of Tree and Vine Crops)  
For citrus crops treat uniformly between trees.

**This Product  
or  
This Product plus Goal**

This product alone or in mixtures with Goal will control or suppress the annual weeds listed below. Apply the recommended rates of this product, either alone or in mixtures with Goal, plus 0.5 to 1 percent nonionic surfactant by spray volume in 3 to 10 gallons of water per acre. Apply when weeds are actively growing and less than 6 inches in height or diameter. If weeds are under drought stress, irrigate prior to application. Reduced control may occur if weeds have been mowed prior to application. Up to 48 fluid ounces per acre of this product may be used to control weeds that have been mowed are stressed, or are growing in dense populations.

Weed Species	Maximum Height/Diameter (Inches)	Rate Per Acre	
		This Product (Fl. Oz.)	Goal (Fl. Oz.)
Barley ( <i>Hordeum vulgare</i> )	6	8	-
Bluegrass, annual ( <i>Poa annua</i> )			
Barnyardgrass ( <i>Echinochloa crus-galli</i> )	6	12	-
Chickweed, common ( <i>Stellaria media</i> )			
Red Maids ( <i>Calandrinia ciliata</i> )			
Crabgrass ( <i>Digitaria spp.</i> )	6	16	or
Fleabane, hairy ( <i>Conyza bonariensis</i> )		16 to 32	+
Groundsel, common ( <i>Senecio vulgaris</i> )			4 to 16**
Junglerice ( <i>Echinochloa colonum</i> )			
Lambsquarters, common ( <i>Chenopodium album</i> )			
Pigweed, redroot ( <i>Amaranthus retroflexus</i> )			
Rocket, London ( <i>Sisymbrium irio</i> )			
Ryegrass, common ( <i>Lolium multiflorum</i> )			
Shepherdspurse ( <i>Capsella bursa-pastoris</i> )			
Sowthistle, annual ( <i>Sonchus oleraceus</i> )			
Cheeseweed, common ( <i>Malva spp.</i> )	3	12 to 32	+
Cheeseweed, common ( <i>Malva spp.</i> )	6	16 to 32	+
Filaree* ( <i>Erodium spp.</i> )			4 to 16
Horseweed/Marestail ( <i>Conyza canadensis</i> )			
Nettle, stinging ( <i>Urtica dioica</i> )			
Purslane, common* ( <i>Purtulaca oleracea</i> )			

\*Suppression only.

\*\*The mixture of this product plus Goal is recommended when weeds are stressed or growing in dense populations.

**Strips**

(For Annual and Perennial Weeds in Strips of Tree and Vine Crops)

**Tank Mixtures with Residual Herbicides**

When applied as a tank mixture, this product provides control of the emerged annual weeds and control or suppression of emerged perennial weeds listed in this label. The following residual herbicides will provide preemergence control of those weeds listed in the individual product labels.

- This Product plus Goal 2XL
- This Product plus Karmex™ DF
- This Product plus Krovar I
- This Product plus Krovar II
- This Product plus Simazine Princep Caliber 90
- This Product plus Simazine 4L
- This Product plus Simazine 80W
- This Product plus Solicam™ 80DF
- This Product plus Surflan AS
- This Product plus Surflan 75W
- This Product plus Simazine (80W, 4L or Princep Caliber 90) plus Surflan (AS or 75W)
- This Product plus Goal 2XL Plus Surflan (AS or 75W)
- This Product plus Goal (2XL) plus Simazine (80W, 4L or Princep Caliber 90)
- This Product plus Goal (2XL) plus Surflan (AS or 75W) plus Simazine (80W, 4L or Princep Caliber 90)

Do not apply these tank mixtures in Puerto Rico.

When tank-mixing with residual herbicides, add an agriculturally-approved nonionic surfactant at 0.5 to 1 percent by volume of spray solution. Refer to the individual product labels for specific crops, rates, geographical restrictions and precautionary statements. Read and carefully observe the label claims, cautionary statements, rates and all other information on the labels of all products.

**Recommended Rates**

**Annual Weeds:** Apply 1 to 5 quarts per acre of this product in these tank mixtures. Use rates at the higher end of the recommended range when weeds are stressed, growing in dense populations or are greater than 12 inches tall.

**Perennial Weeds:** Apply 1 pint to 5 quarts per acre of this product in these tank mixtures to control or suppress perennial weeds. Follow the recommendations in the **Weeds Controlled** section of this label for stage of growth and application rates for specific perennial weeds.

**This Product plus Goal plus Simazine/Surflan**

This product plus low rates of Goal in 3-way or 4-way mixtures with simazine and/or Surflan will provide postemergence control of the weeds listed below. Refer to the individual simazine and Surflan labels for preemergence rates, weeds controlled, precautionary statements and other important information. Apply these tank mixtures in 3 to 40 gallons of water. Add 0.5 to 1 percent nonionic surfactant by total spray volume to the spray solution. Apply 1 to 5 quarts per acre of this product plus 4 to 48 fluid ounces per acre of Goal plus labeled rates of simazine and/or Surflan to control the following weeds:

- |  |  |
|--|--|
| Barley, wild<br><i>Hordeum leporinum</i>     | Horseweed/Marestail<br><i>Conyza canadensis</i>  |
| Bluegrass, annual<br><i>Poa annua</i>        | Nettle, stinging<br><i>Urtica dioica</i>         |
| Cheeseweed, common<br><i>Malva</i> spp.      | Pineappleweed<br><i>Matricaria matricariodes</i> |
| Chickweed, common<br><i>Stellaria media</i>  | Rocket, London<br><i>Sisymbrium irio</i>         |
| Filaree*<br><i>Erodium</i> spp.              | Shepherdspurse<br><i>Capsella bursa-pastoris</i> |
| Fleabane, hairy<br><i>Conyza bonariensis</i> | Sowthistle, annual<br><i>Sonchus oleraceus</i>   |
| Groundsel, common<br><i>Senecio vulgaris</i> |  |

\*Use a minimum of 1.5 quarts of this product in these mixtures.

**Note:** This recommendation does not preclude the use of Goal in these mixtures at higher, labeled rates for preemergence weed control.

**Perennial Grass Suppression  
Orchard Floors**

When applied as directed, this product will suppress vegetative growth as indicated below.

**Bahiagrass:** This product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with a single application and approximately 120 days with sequential applications. Apply this product 1 to 2 weeks after full green-up or after mowing to a uniform height of 3 to 4 inches. Applications must

be made prior to seedhead emergence. Apply 6 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of this product plus nonionic surfactant. A second sequential application of 2 to 4 fluid ounces may be made approximately 45 days after the last application.

**Bermudagrass:** For burndown, apply 1 to 2 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre. Use 1 quart of this product in 3 to 20 gallons of water per acre east of the Rocky Mountains. Use 1 to 2 quarts of this product in 3 to 10 gallons of water per acre west of the Rocky Mountains. Use this treatment only if reduction of the bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

**Suppression only (east of the Rocky Mountains):** Apply 6 to 16 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 20 gallons of water per acre no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Use rates of 6 to 10 fluid ounces of this product plus nonionic surfactant in shaded conditions or where a lesser degree of suppression is desired. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

**Suppression only (west of the Rocky Mountains):** Apply 16 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre to bermudagrass up to 6 inches in height and no sooner than 1 to 2 weeks after full green-up. Mowing prior to application may occur provided a minimum height of 3 inches is maintained. Sequential applications may be made when regrowth occurs and bermudagrass injury and stand reduction can be tolerated.

**Cool Season Grass Covers:** For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. For best suppression, add ammonium sulfate to the spray solution at a rate of 2 percent by weight of 17 pounds per 100 gallons of spray solution. For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of this product plus 0.5 to 1 percent nonionic surfactant. Do not add ammonium sulfate. For best results, mow cool-season grass covers in the spring to even their height and apply the recommended rate of this product 3 to 4 days after mowing. Avoid treating cool season grass covers under poor growing conditions, such as drought stress (drip irrigation), disease or insect damage.

**Low Volume Application  
(Florida and Texas)**

For burndown or control of the weeds listed, apply the recommended rates of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 30 gallons of water per acre. When weed foliage is dense, use 10 to 30 gallons of water per acre.

**Annual Weeds**

**Goatweed:** Apply 2 to 3 quarts per acre of this product plus 17 pounds of ammonium sulfate per 100 gallons of water plus 0.5 to 1 percent nonionic surfactant by total spray volume. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches. If goatweed is greater than 8 inches tall, the addition of Krovar II or Karmex may improve control. Use labeled rates for these residual products. Read and carefully observe the label claims, cautionary statements, rates and all other information on the Krovar II and Karmex labels.

**Perennial Weeds**

Apply when weeds are actively growing and at the growth stages listed in the **Perennial Weeds Controlled** section of this label. If perennial weeds are mowed, allow weeds to regrow to the recommend stage of growth.

S = Suppression B = Burndown PC = Partial Control C = Control

Weed Species	This Product Rate Per Acre			
	1 Qt.	2 Qts.	3 Qts.	5 Qts.
Bermudagrass	B	•	PC	C
Guineagrass Texas and Florida Ridge	B	C	C	C

Florida Flatwoods	•	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	•	PC	C

**Tree Crops**

**Citrus\*\*\*\*:** calamondin, chironja, citron, grapefruit, kumquat, lemon, lime, mandarin orange, orange, pummelo, tangelo, tangerine, tangors.

**Nuts\*\*:** almond, beechnut, Brazil nut, butternut, cashew, chestnuts, chinquapin, filbert, hazel nut, hickory nut, macadamia, pecan, pistachio, walnut.

**Pome Fruit\*\*\*\*:** apple, loquat, mayhaw, pear, quince.

**Stone Fruit\*\*\*:** apricots, cherries, nectarines, olives, peaches, plums/prunes.

For cherries, any application equipment listed in this section may be used in all states. For citron and olives, apply only as a directed spray. Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in the states specified in the following paragraph. In all other states use only wiper equipment.

For **peaches** grown only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee, apply with a shielded boom sprayer or shielded wiper applicator, that prevents any contact of this product with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. **Extreme care must be taken to ensure no part of the peach tree is contacted.**

**Tropical Fruit:** acerola\*, atemoya\*, avocado\*, banana\*\*\*\*, breadfruit\*, canistel\*, carambola\*, cherimoya\*, cocoa beans\*, coffee\*\*\*\*, dates\*, figs\*, genip\*, guava\*\*\*\*, jaboticaba\*, jackfruit\*, logan\*, lychee\*, mango\*, mayhaw\*, papaya\*\*\*\*, passion fruit\*, persimmons\*, plantains\*\*\*\*, pomegranate\*, sapodilla\*, sapote\*, soursop\*, sugar apple\*, tamarind\*, tea\*. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

- Note:**
- \*Allow a minimum of 14 days between last application and harvest.
  - \*\*Allow a minimum of 3 days between last application and harvest of these crops
  - \*\*\*Allow a minimum of 17 days between last application and harvest.
  - \*\*\*\*Allow a minimum of 28 days between last application and harvest.
  - \*\*\*\*\*Allow a minimum of 1 day between last application and harvest.

**Vine Crops**

**Kiwi Fruit**

**Grapes:** Any variety of table, wine or raisin grape may be treated with any equipment listed in this section. Do not apply when green shoots, canes or foliage are in the spray zone. Allow a minimum of 14 days between last application and harvest. In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury.

- Bullet, Lariat, Lasso, Micro-Tech and Partner are registered trademarks of Monsanto Company.
- Oust, Krovar, Escort, Telar, Canopy, Gemini, Lexone, Lorox Plus, Diruon, Preview and Karmex are trademarks of E.I. duPont de Nemours and Company.
- Dual, Dual Magnum, Bicep, Bicep Magnum, Princep and Caliber are trademarks of Syngenta Corporation.
- Surflan is a trademark of Dow AgroSciences LLC
- Banvel and Solicam are trademarks of BASF, Inc.
- Ronstar, Folex and Prep are trademarks of Aventis Environmental Science
- Goal is a trademark of Rohm and Haas Company.
- Sencor and Turbo are trademarks of Bayer AG.
- Prowl, Pursuit, Pursuit Plus, Scepter and Squadron are trademarks of American Cyanamid Company.
- Command is a trademark of FMC Corporation.
- DEF is a trademark of Mobay Chemical Company.

[ ] Denotes alternate/optional language  
 { } Denotes language that does not appear on market labeling

See  
 Pass  
 SS  
 and F

### LABEL HISTORY

FILE NAME	REVISION DATE	COMMENTS
035935-00007.20110729.EPA Not	(RV072911)	PRN 98-10 & 07-04