

48273-11

08/13/2001

1/19

AUG 13 2001

Ms. Kim Davis
Consultant/Agent
MARMAN USA, INC.
36856 Rocky Road
Greeley, CO 80631-9375

SUBJECT: Label Amendment
Atrazine 4L Herbicide, EPA Reg. 48273-11
Your Application Dated July 17, 2001

Dear Ms. Davis:

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provide that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. It is recommended that all "Precautions" and "Note" be changed to "Use Precaution and Restrictions"
2. Add the following under "PPE" "Some materials that are chemical-resistant to this product are listed below. If you want more options, follow instructions for category A on an EPA chemical resistant category selection chart."

"Under "applicators and other handlers must wear" and in the "Agricultural Use Requirements" box change "waterproof gloves" to "Chemical-resistant gloves, such as _____ or _____ or _____ or _____."

CONCURRENCES

SYMBOL ▶	75050							
SURNAME ▶	MINOR, E.							
DATE ▶	Aug 8, 2001							

{Booklet}

RESTRICTED USE PESTICIDE
(Ground and Surface Water Concerns)

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification. This product is a restricted-use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

ATRAZINE 4L HERBICIDE

For Season-Long Weed Control in Corn, Sorghum and Certain Other Crops

ACCEPTED
with **COMMENTS**
In EPA Letter Dated

AUG 13 2001

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Active Ingredients:	
Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	40.8%
Related compounds	2.2%
Other Ingredients	57.0%
Total	100.0%

48273-11

Contains 4 pounds active ingredients per gallon.
Shake well before using.

Keep Out of Reach of Children

CAUTION

See additional precautionary statements and directions for use inside booklet.

_____ Gallons
U.S. Standard Measure

EPA Reg. No. 48273-11 EPA Est. _____

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire *Directions for Use* and the *Conditions of Sale and Warranty* before using this product. If terms are not acceptable return unopened product container at once.

Conditions of Sale and Warranty

The *Directions for Use* of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Marman USA, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Marman USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the *Directions for Use* subject to the inherent risks referred to above. **Marman USA, Inc. makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Marman USA, Inc. or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.** Marman USA, Inc. and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing *Conditions of Sale and Warranty*, which may be varied only by agreement in writing and signed by a duly authorized representative of Marman USA, Inc.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Waterproof gloves
- Chemical-resistant footwear plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury or illegal residues. Do not apply this product through any type of irrigation system.

General Information

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane and certain other crops specified on this label. This product may be applied before or after weeds emerge.

Following many years of continuous use of this product and chemically-related products, biotypes of some of the weeds listed on this label have been reported that cannot be effectively controlled by this and related herbicides. Where this is known or suspected, and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of this product in combinations or in sequence with other registered herbicides that are not triazines. If only resistant biotypes are expected to be present, use a registered non-triazine herbicide. Consult with your State Agricultural Extension Service for specific recommendations.

Since this product acts mainly through root absorption, its effectiveness depends on moisture to move it into the root zone. If weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control. This product is noncorrosive to equipment and metal surfaces, nonflammable and has low electrical conductivity.

Avoid using near adjacent desirable plants or in greenhouses, or injury may occur. To avoid spray drift, do not apply under windy conditions. Avoid spray overlap as crop injury may result. Where the use directions give a range of rates, use the lower rate on coarse-textured soil and soil low in organic matter; use the higher rate on fine-textured soil and soil high in organic matter. **Note:** Marman USA, Inc. does not recommend applications in combination with other herbicides or oils, except as specifically described on this label or in literature published by Marman USA, Inc.

Application Procedures

Ground Application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate the sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all preplant incorporated, preplant surface, preemergence and postemergence applications (with or without oil or surfactant) with ground equipment.

Use a pump with capacity to: (1) maintain 35-40 psi at nozzles; (2) provide sufficient agitation in tank to keep mixture in suspension; and (3) provide a minimum of 20% bypass at all times. Use centrifugal pumps that provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons per minute per 100 gallons tank size circulated through a correctly-positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

Aerial Application: Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 quart of water for each quart of this product applied per acre. For postemergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gallons of water per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

Application in Water or Liquid Fertilizer: Nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence, preplant incorporated or preplant surface ground application on corn and sorghum. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below before use. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges or crop injury may occur.

Compatibility Test: Since liquid fertilizers can vary, even within the same analysis, always **check compatibility with herbicide(s) each time before use**. Be especially careful when using **complete** suspension or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gallons per acre. For other spray volumes, make appropriate changes in the ingredients. Check compatibility using this procedure:

1. Add 1 pint of fertilizer to each of 2 one-quart jars with tight lids.
2. To **one** of the jars add ¼ tsp. or 1.2 milliliters of a compatibility agent approved for this use, such as Compex® or Unite® (¼ tsp. is equivalent to 2 pts. per 100 gals. spray). Gently shake or stir to mix.
3. To **both** jars add the appropriate amount of herbicide(s). If more than one herbicide is used, add them separately with dry herbicides first, flowables next and emulsifiable concentrates last. After each addition, gently shake or stir to thoroughly mix. The appropriate amount of herbicides for this test follows:
 - Dry Herbicides: For each pound to be applied per acre, add 1.5 level teaspoon to each jar.
 - Liquid Herbicides: For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.
4. After adding all ingredients, put lids on a tighten, and invert each jar 10 times to mix. Let the mixtures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be readily remixed, the mixture can be sprayed provided good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility:
 - a. slurry the dry herbicide(s) in water before addition, or
 - b. add ½ of the compatibility agent to the fertilizer and the other ½ to the emulsifiable concentrate or flowable herbicide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

Application in Water Plus Emulsifiable Oil or Oil Concentrate: Adding emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate or single or mixed crop derived oil concentrate) to postemergence water-based sprays in corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe the following directions:

Use one of the following properly emulsified:

1. A suitable oil concentrate containing at least 1% but not more than 20% suitable emulsifier or surfactant blend.
2. Petroleum-derived oil containing at least 1% suitable emulsifier.

Note: In the event of a compatibility problem when mixing oil with this product and water, a compatibility agent such as Compex or Unite should be used. Any of the above oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

Mixing Procedures – All Uses:

1. Be sure sprayer is clean and not contaminated with any other materials, or crop injury or sprayer clogging may result.

2. Fill tank ¼ full with clean water, nitrogen solution or complete liquid fertilizer.
3. Start agitation
4. Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface.
5. Pour product directly into tank.
6. Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action.
7. Add emulsifiable oil, oil concentrate or tank mix herbicide(s) after this product is thoroughly suspended.
8. Finish filling tank.
9. Empty tank as completely as possible before refilling to prevent buildup of oil or emulsifiable concentrate residue. Maintain agitation to avoid separation of materials.
10. If an oil or emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent.
11. Immediately after use, thoroughly clean sprayer by flushing system with water containing a detergent.

Rotational Crops – All Uses:

1. Do not rotate to any crop except corn or sorghum until the following year or injury may occur.
2. If applied after June 10, do not rotate with crops other than corn or sorghum the next year or crop injury may occur.
3. In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to follow corn or sorghum or when a crop of untreated corn or sorghum is to precede other rotational crops.
4. In eastern parts of the Dakotas, KS, western MN and NE, do not rotate to soybeans if the rate applied to corn or sorghum was more than 4 pts./A or equivalent band application rate, or soybean injury may occur.
5. Injury may occur to soybeans planted the year following application on soils having a calcareous surface layer.
6. Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains or small-seeded legumes and grasses the year following application or injury may occur.

Atrazine 4L Applied Alone Corn or Grain Sorghum*

Preplant Surface-Applied, Preplant Incorporated or Preemergence (or Postemergence at 4 pts./A with Oil)

Broadleaf and Grass Weeds Controlled

Annual Morningglory	Mustards
Barnyardgrass (watergrass)***	Nightshade
Cocklebur**	Pigweed
Giant Foxtail**	Purslane
Green Foxtail***	Ragweed
Groundcherry	Sicklepod**
Jimsonweed	Velvetleaf (buttonweed)***
Kochia	Wild Oats
Lambsquarters	Witchgrass (<i>Panicum capillare</i>)***
Large (hairy) crabgrass**	Yellow Foxtail***

Postemergence with Emulsifiable Oil or Oil Concentrate in Water (at 2.4 pts./A)

Broadleaf Weeds Controlled

Annual Morningglory	Pigweed
Cocklebur	Ragweed
Jimsonweed	Smartweed
Lambsquarters	Velvetleaf**
Mustards	Wild Buckwheat

*Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or greater setbacks) that are different from the label the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

**Only partial control.

***Only partial control on medium- and fine-textured soils.

CORN

Preplant Surface-Applied (Broadleaf and Grass Control): Use on medium- and fine-textured soil with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI and WY. Apply the recommended rate of this product 10 to 14 days up to 45 days prior to planting. On coarse-textured soils, do not apply more than two

weeks prior to planting. If an unsatisfactory length of weed control results from adverse environmental conditions following early treatment, a follow-up application of an appropriately labeled herbicide may be used. If the follow-up treatment includes atrazine do not exceed the labeled rate for corn indicated in Table 1. If weeds are present at the time of treatment, apply in tank mix combination with a contact herbicide (for example, Gramoxone® Extra or Roundup®). Observe directions for use, precautions and restrictions on the label of the contact herbicide.

Note: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated (Broadleaf and Grass Control): Broadcast in spring after plowing at rate in Table 1. Apply to the soil and incorporate before, during or after final seedbed preparation. Avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence or At-Planting (Broadleaf and Grass Control): Apply during or shortly after planting before weed emergence, at rate in Table 1.

Postemergence (Broadleaf and Grass Control): Apply before weeds exceed 1.5 inches in height and before corn exceeds 12 inches in height at rate in Table 1.

**Table 1
Broadleaf and Grass Weed Control in Corn***

For All Soil Applications Prior to Crop Emergence	
<ul style="list-style-type: none"> On Highly Erodible Soils (as defined by SCS): If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at planting, apply a maximum of 4 pts./A as a broadcast spray. If the soil coverage with plant residue is less than 30% at planting, a maximum of 3.2 pts./A may be applied. On soils Not Highly Erodible: Apply 4 pts./A as a broadcast spray 	
For Postemergence Application:	
If not atrazine was applied prior to corn emergence, apply a maximum of 4 pts./A broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. active ingredient (5 pts. of this product) per acre per calendar year.	

***Broadleaf Control (eastern CO, western KS, western NE, NM, OK Panhandle, west TX and eastern WY):** On sand, loamy sand, sandy loam, mild to strongly alkaline soil and all recently leveled soil, apply no more than 2.4 pts./A, either preplant surface, preplant incorporated or preemergence. On other soils in these areas, apply rate in Table 1 for broadleaf and grass control.

Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or greater setbacks) that are different from the label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

Postemergence with Emulsifiable Oil or Oil Concentrate in Water: Add the following volume of one of the type oils indicated for aerial or ground application unless the oil label specifies otherwise:

Type Oil	Ground Application	Aerial Application
Oil concentrate (crop or petroleum derived)	1 qt./A	½ = 1 qt./A
Petroleum-derived oil	1 gal./A	2 qts./A

Note: Crop-derived or petroleum-derived oil concentrates should contain at least 1% but not more than 20% suitable emulsifier or surfactant blend. Petroleum-derived oils should contain at least 1% suitable emulsifier.

Broadleaf and Grass Control: For postemergence control of those weeds listed under *Preplant Incorporated and Preemergence*, broadcast 4 pts./A plus emulsifiable oil or oil concentrate after weed emergence, but before weeds reach 1.5 inches in height and before corn exceeds 12 inches in height.

Broadleaf Control: For postemergence control of those weeds listed under *Postemergence with Emulsifiable Oil or Oil Concentrate in Water*, broadcast 2.4 pts./A plus emulsifiable oil or oil concentrate before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

Precautions For Applications with Emulsifiable Oil or Oil Concentrate in Water:

1. Inbred lines or any breeding stock may be severely injured by applications with emulsifiable oil or oil concentrate.
2. Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended, because they may cause compatibility problems or crop injury.
3. Carefully store and handle emulsifiable oil and oil concentrate. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
To avoid crop injury,
4. Do not apply when crop is under stress from prolonged cold, wet weather, poor fertility or other factors, or when crop is wet and succulent from recent rainfall.
5. Do not exceed 2.5 lbs. active ingredient (or 5 pts. of this product) per acre per calendar year.
6. Postemergence applications to corn must be made before corn exceeds 12 inches in height.

Tank Mixtures for Corn

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

Dual® (metolachlor)	Lasso or Lasso EC + Roundup
Dual + Gramoxone Extra	Lasso or Lasso EC + Gramoxone Extra
Dual + Roundup (glyphosate)	Gramoxone Extra
Dual + Princep®	Princep
Dual + Princep + Gramoxone Extra	Princep + Gramoxone Extra
Dual + Princep + Roundup	Princep + Roundup
Bexton® or Ramrod® (propachlor)	Roundup
Lasso® or Lasso EC (alachlor)	Sutan+®

Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of products used in a particular tank mix. **Note:** When the labels of the above herbicides refer to atrazine 80W, use equivalent rates of this product. One pound of 80W equals 1.6 pints of this product.

Princep 80W, Princep 4L or Princep Caliber 90®

In addition to the weeds listed under *Atrazine 4L Applied Alone – Corn or Grain Sorghum: Preplant Surface-Applied, Preplant Incorporated or Preemergence*, this combination also controls crabgrass, fall panicum and carpetweed. Broadcast tank mix before planting, at planting or after planting, but before crop and weeds emerge, at rates in Table 2. Use the 1:1 ratio for control of most weeds. Use the 1:2 ratio for expected heavy infestations of crabgrass and fall panicum. Cultivate shallowly if weeds develop.

Preplant Surface-Applied: Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI and WY. Apply the recommended rate of this product and Princep shown in Table 2 up to 45 days prior to planting. Refer to the *Atrazine 4L Alone* section for information if weeds should develop following the early treatment. On coarse-textured soils, do not apply more than two weeks prior to planting. Refer to the *Atrazine 4L Applied Alone – Preplant Surface-Applied* section of the corn label for additional details. If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, Gramoxone or Roundup). Observe directions for use, precautions and restrictions on the label of the contact herbicide. **Note:** To the extent possible, do not more treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated: Apply to the soil and incorporate in the spring before, during or after final seedbed preparation. Avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence: Apply during or shortly after planting, but before crop and weeds emerge.

Refer to the corn sections of this label and to Princep 80W, Princep Caliber 90 or Princep 4L labels for further directions, limitations and precautions.

Table 2
Tank Mixtures with Princep on Corn

Soil Texture	Broadcast Rate Per Acre			
	1:1 Ratio*		1:2 Ratio**	
	This Product	Princep 80W ¹	This Product	Princep 80W ¹
Sand, loamy sand, sandy loam	2 pts.	1.25 lbs.	1.32 pts.	1.67 lbs.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay or silty clay with low organic matter	2.4 pts.	1.5 lbs.	1.6 pts.	2 lbs.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	3 pts.	1.8 lbs.	1.92 pts.	2.4 lbs.

*For control of most weeds.

**For control of expected heavy infestations of crabgrass and fall panicum.

¹When using Princep Caliber 90 or Princep 4L, use equivalent rates. One pound of Princep 80W equals 0.9 pounds of Princep Caliber 90 or 1.6 pints of Princep 4L.

Princep 80W, Princep 4L or Princep Caliber 90 plus Roundup: Use as tank mixture for preemergence and postemergence control of certain broadleaf and grass weeds where corn will be planted directly into a cover crop, established sod or in previous crop residues. Refer to Roundup label for all directions, weeds controlled, precautions and limitations.

Princep 80W, Princep 4L or Princep Caliber 90 plus Gramoxone Extra: use as tank mixture with Princep and Gramoxone Extra to kill existing vegetation and for residual weed control where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Add this product and Princep to water in spray tank, agitating until thoroughly mixed. Then add Gramoxone Extra and a nonionic surfactant, such as X-77[®]. Continue agitation during application. Broadcast 2-4 pts. of this product plus 1.25-2.5 lbs. of Princep 80W (or 2-4 pts. Princep 4L, or 1.1-2.2 lbs. Princep Caliber 90) plus a suitable amount of Gramoxone Extra in 20-60 gals. of water per sprayed acre. Refer to the Gramoxone Extra label for the appropriate rates to utilize in this tank mixture. Apply before, during or after planting, but before corn emerges. Add 0.5 pt. of a nonionic surfactant, such as X-77, per 100 gals. of spray mixture. Use the higher rate of Gramoxone Extra specified on the label if existing vegetation is 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Refer to further limitations and precautions on labels for this product, Princep and Gramoxone.

Precautions for All Applications to Corn:

1. To avoid crop injury and illegal residues, do not apply more than 2.5 lbs./A active ingredient (5 pts. of this product) per calendar year.
2. For best control of velvetleaf and cocklebur, the application rate cannot be less than 2 lbs./A active ingredient, either alone or in tank mix combinations.
3. Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used.

Note for All Applications to Corn: Do not graze or feed forage from treated areas for 21 days following application, or illegal residues may result.

SORGHUM AND SORGHUM-SUDAN HYBRIDS
(Grain and Forage Types)

Preplant Surface-Applied (Broadleaf and Grass Control): Use on medium- and fine-textured soil with minimum-tillage or no-tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI and WY. Apply the recommended rate of this product shown in Table 3 up to 45 days prior to planting. If an unsatisfactory length of weed control results from adverse environmental conditions following early treatment, a follow-up application of an appropriately labeled herbicide may be used. If the follow-up treatment includes atrazine, do not exceed the labeled rate for corn indicated in Table 1. Under dry conditions, irrigation after application is recommended to move this product into the soil.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide (for example, Gramoxone Extra or Roundup). Observe directions for use, precautions and restrictions on the label of the contact herbicide. **Note:** To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting or weed control will be diminished.

9/19

Preplant Incorporated (Broadleaf and Grass Control): Broadcast in spring after plowing at rate shown in Table 3. Apply before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence or At-Planting (Broadleaf and Grass Control): Apply during or shortly after planting, but prior to weed or crop emergence at rate shown in Table 3.

Postemergence (Broadleaf and Grass Control): Apply at rate shown in Table 3 before weeds exceed 1.5 inches in height and before sorghum exceeds 12 inches in height.

**Table 3
Broadleaf and Grass Weed Control in Sorghum^{1,2}**

<p>For All Soil Applications Prior to Crop Emergence</p> <ul style="list-style-type: none"> • On Highly Erodible Soils (as defined by SCS): If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at planting, apply a maximum of 4 pts./A as a broadcast spray. If the soil coverage with plant residue is less than 30% at planting, a maximum of 3.2 pts./A may be applied. • On soils Not Highly Erodible: Apply 4 pts./A as a broadcast spray <p>For Postemergence Application: If not atrazine was applied prior to sorghum emergence, apply a maximum of 4 pts./A broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. active ingredient (5 pts. of this product) per acre per calendar year.</p>

¹Do not apply preplant surface or preplant incorporated in AL, AR, FL, GA, LA, MS, NM, NC, OK, SC, TN or TX. Do not apply preemergence in NM, OK or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas.

²Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or greater setbacks) that are different from the label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

In case of planting failure, sorghum or corn may be replanted. Do not make a second broadcast application or injury may occur. If originally applied in a band and sorghum or corn is replanted in untreated row middles, this product may be applied in a band to the second planting, provided the maximum application rate of 2.5 lbs. a.i./A of atrazine per calendar year is not exceeded.

Preemergence Broadleaf Weed Control in Furrow-Irrigated Bedded Sorghum (Only AZ and CA): For preemergence control of many broadleaf weeds, broadcast 1.6 to 2.4 pts./A after bed preparation, during or after planting, but before sorghum and weeds emerge and before the first furrow irrigation. Follow with several regular irrigations, making sure to thoroughly wet all soil.

Precautions for Preemergence Application to Furrow-Irrigated Bedded Sorghum in AZ and CA: To avoid possible sorghum injury, do not use on sand or loamy sand soil or on sorghum planted in furrows. Applications to sorghum growing on alkali soils or where cuts, fills or erosion have exposed calcareous or alkali subsoils may cause crop injury. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence Broadleaf Weed Control with Emulsifiable Oil or Oil Concentrate in Water: Broadcast 2.4 pts./A for control of many broadleaf weeds. Apply before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX and desert regions of AZ and CA, apply when sorghum

is 6 to 12 inches in height, but before it reaches boot stage. In all other areas, apply after sorghum reaches the 3-leaf stage, but before it exceeds 12 inches in height. Add 1 gallon of emulsifiable oil per acre for ground application and 0.5 gallon per acre for aerial application, or add 1 quart per acre of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

For the list of weeds controlled, see *Atrazine 4L Applied Alone – Corn or Grain Sorghum – Postemergence with Emulsifiable Oil or Oil Concentrate in Water*.

Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water: See *Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water* in *Corn* section.

Postemergence Broadleaf Weed Control with Surfactant (Only CO, western KS, NM, OK, TX and desert regions of AZ and CA): Broadcast 2.4 pts./A plus 0.75 to 1.5 pts. of surfactant after sorghum reaches 6 inches in height, but before weeds exceed 1.5 inches in height. Apply only on sandy loam and fine-textured soil.

Precautions for all Applications to Sorghum:

1. Heavy rain immediately following application tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Do not apply to furrow-planted sorghum until furrows are leveled (plowed in). Level deep planter marks or seed furrows before application.
2. Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury.
3. Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used.
4. Injury may occur if both this herbicide, preplant surface, preplant incorporated or preemergence, and an at-planting systemic insecticide applied in-furrow are used.
5. Do not apply more than 2.5 lbs. active ingredient (5 pts. of this product) per acre per calendar year.
6. For all soil applications prior to crop emergence (except for preemergence use on bedded sorghum in AZ and CA) do not apply to coarse-textured soils, i.e., sand, loamy sand, sandy loam, or to medium- and fine-textured soils having less than 1% organic matter or injury may occur.
7. For postemergence applications, do not apply to sand or loamy sand or injury may occur.

Note: Do not graze or feed forage from treated areas for 21 days following application or illegal residues may result.

TANK MIXTURES FOR GRAIN SORGHUM

Dual 8E: Use as tank mixture with Dual 8E for control of those weeds listed on the Dual 8E label, as well as on this label. Use this tank mixture only on sorghum seed treated with Concep[®]. Refer to the Dual 8E label for all directions, precautions and limitations.

WINTER WEED CONTROL IN TEXAS

Only for postemergence control of winter weeds, such as henbit, seedling dock and annual thistle on fall bedded land in the Gulf Coast and Backlands of Texas. Apply 0.8 to 1.0 qt./A postemergence to the weeds in November or December to land that will be planted to corn, grain sorghum or forage sorghum the following spring. For best results add a suitable surfactant, such as X-77, at the rate of 0.5% of the spray volume, an emulsifiable oil at the rate of 1.0% of the spray volume or an oil concentrate at the rate of 1 qt./A. Normal weed control programs may be used in the following corn, grain, sorghum or forage sorghum crop. **Note:** Do not plant any crops, except corn, grain sorghum or forage sorghum, the spring following this treatment or illegal residues may result.

ATRAZINE 4L ALONE – CHEMICAL FALLOW

Wheat-Sorghum-Fallow: To control annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum tillage, broadcast 6 pts./A to wheat stubble immediately following wheat harvest. If weeds are present at application, remove them with a sweep plow or other suitable implement after application, or use an approved contact herbicide before or after the application of this product. Plant sorghum into wheat stubble the following spring with minimum soil disturbance. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see *Atrazine 4L Applied Alone – Corn or Grain Sorghum – Preplant Surface-Applied, Preplant Incorporated or Preemergence*.

Precautions:

1. Use only on sandy loam or fine-textured soil or crop injury may result.

2. Wheat-sorghum-fallow cropping sequence must be followed.
3. Do not apply following sorghum harvest.

Note: To avoid illegal residues, do not graze or feed forage from treated area to livestock. To avoid illegal residues and crop injury, do not plant any crop other than those on this label within 18 months following treatment.

Wheat-Corn-Fallow (CO, KS, NE, ND, SD and WY): This product controls cheatgrass (downy brome, chess), kochia, mustards, pigweed, Russian thistle, wild lettuce, wild sunflower and volunteer wheat during the period after wheat harvest. Weed control may extend into following corn crop grown under minimum tillage.

On soils in ND and SD with a pH greater than 7.5, do not exceed 1.5 lbs. a.i./A. For soils with a pH less than 7.5 in ND and SD apply 1 to 2 lbs. a.i./A. Use the higher rate on fine-textured soils and where heavy weed infestations are expected. Use the lower rate on coarse-textured soils and where light weed infestations are expected. In the event grasses are present in the following spring, use a grass herbicide registered for use on corn.

Follow directions for use, notes and precautions in the *Wheat-Sorghum-Fallow* section above, substituting corn for references to sorghum.

Wheat-Fallow-Wheat (CO, KS, NE, ND, SD and WY): For preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, broadcast 1 to 2 pts./A on all soils except those listed under *Precautions*. For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

TANK MIXTURES FOR CHEMICAL FALLOW
Wheat-Sorghum-Fallow or Wheat-Corn-Fallow (KS, NE)

Gramoxone Extra: If weeds are present at application, a tank mix with Gramoxone Extra may be used. Broadcast 6 pts. of this product plus a suitable amount of Gramoxone Extra in 20 to 60 gallons of water per acre by ground equipment. Add 0.5 to 1 pint of a nonionic surfactant, such as X-77, per 100 gallons of spray mixture. Add this product to the spray tank first and thoroughly mix with water, then add Gramoxone Extra, followed by surfactant. Use the higher rate of Gramoxone Extra specified on the label if weeds are 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to the Gramoxone Extra label for further directions, precautions and limitations.

Wheat-Fallow-Wheat (CO, KS, NE, ND, SD and WY)

Gramoxone Extra: If weeds are present at application, a tank mix with Gramoxone Extra may be used. Broadcast 1 to 2 pints of this product plus a suitable amount of Gramoxone Extra in 20 to 60 gallons of water per acre by ground equipment. Add 0.5 to 1.0 pint of a nonionic surfactant, such as X-77, per 100 gallons of spray mixture. Add this product to spray tank first and thoroughly mix with water, then add Gramoxone Extra, followed by surfactant. Use the higher rate of the Gramoxone Extra specified on the label if weeds are 4 to 6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to Gramoxone Extra label for further directions, precautions and limitations.

If weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment. Use tillage to control weeds that escape during fallow period. Till before planting. For this product applied alone or in tank mixture with Gramoxone Extra, plant at least 2 inches deep and 12 months or more after application.

Precautions: To avoid crop injury,

1. Do not use on sandy soil.
2. Do not treat eroded hillsides, caliche and rocky outcroppings or exposed calcareous subsoil.
3. Do not treat soils of the Rosebud and Canyon Series in western NE and adjoining counties in CO and WY.
4. Do not treat soils with calcareous surface layers.
5. Avoid spray overlap.

Note: Do not graze treated areas within 6 months after application or illegal residues may result.

Aerial Application: To assure that the spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 feet, using low-drift nozzles at a maximum pressure of 40 psi and restrict application to periods when wind speed does not exceed 10 mph. To assure that the drift will not adversely affect

adjacent sensitive nontarget plants, apply this product alone by aircraft at a minimum upwind distance of 400 feet from sensitive plants.

ROADSIDES

To control certain annual weeds in established perennial grasses along roadsides in CO, KS, MT, NE, ND, SD and WY, including cheatgrass (downy brome, chess), common (annual) broomweed, little barley, medusahead, sagewort and tumble mustard, broadcast 2 pts./A in a minimum of 10 gallons of water by ground equipment in the fall before the ground freezes, or after thawing in the spring, but before established grasses green-up and weeds emerge. Examples of desirable established grasses include big bluestem, bluegrama, bromegrass, buffalograss, crested wheatgrass, indiagrass, little bluestem, side-oats grama, switchgrass and western wheatgrass. Apply only once per year. Temporary discoloration or other form of injury to the desirable perennial grasses may occur following application. **Notes:** To avoid illegal residues:

- 1. Do not cut or feed roadside grass hay.
- 2. Do not allow livestock to graze treated areas.

CONSERVATION RESERVE PROGRAM (CRP) (NE, OK, OR and TX)

For control or suppression of annual ragweeds, barnyardgrass, black nightshade, cheat, cocklebur, downy brome, fall panicum, field pennycress, giant foxtail, yellow foxtail, Japanese brome, Kentucky bluegrass, kochia, lambsquarters, little barley, marehail, pigweed, prickly lettuce, smooth brome and sunflower refer to the directions, notes and precautions below:

Pure Stands of Newly Seeded Big Bluestem, Switchgrass and Eastern Gamagrass: Use only on loam, silt loam, silty clay loam, clay loam and silty clay soils with at least 1% organic matter.

Establishment: Broadcast 2 qts./A of this product preplant incorporated or preemergence at time of seeding and prior to emergence of weeds. Prepare a good firm seedbed. Plant 1/2 inch deep with a grassland drill (preferred method) or a conventional drill. If a conventional drill is used on prepared seedbeds remove all tension from the disk openers. For best results cultipack or roll after planting.

Renovation of Existing Stands of Big Bluestem and Switchgrass Planted on CRP Acres: Broadcast 1 to 2 quarts per acre to existing stands of big bluestem and switchgrass prior to the emergence of weeds. Use the low rate on soils containing from 1 to 2% organic matter. Use the high rate on soils with 2% or more organic matter.

Renovation of Existing Stands of the following perennial range grasses planted on CRP acres: blue grama, indiagrass, little bluestem, sand lovegrass, sideoats grama and western wheatgrass. Broadcast 0.5 to 1.0 quart per acre in the spring prior to weed emergence or in the fall before the ground freezes and prior to weed emergence after these species have been established for at least one growing season for control or partial control of the weeds listed above. Use the lower rate for weeds easily controlled or suppressed. Use the higher rate on other weeds claimed in an earlier section of this label.

Aerial Application: Make applications at a maximum height of 10 feet above vegetation. Use low-drift nozzles at a maximum pressure of 40 psi. To control drift, restrict application to periods when wind speed does not exceed 10 mph. To assure that drift will not adversely affect adjacent sensitive nontarget plants, apply this product by aircraft at a minimum upwind distance of 400 feet from sensitive plants. Use 3 to 5 gallons per acre total water volume; use the higher water volume when a dense, heavy ground cover is present.

Notes:

- 1. Do not cut or feed grass hay to livestock.
- 2. Do not graze treated areas.
- 3. Do not use seeds for bird food.
- 4. Do not dump or spill product or dispose of containers within reach of livestock.
- 5. Follow all applicable restrictions for the Conservation Reserve Program.

Precaution: To avoid crop injury, make only one application per year. Slight discoloration of desirable grasses may occur following treatment. Injury may be enhanced when used on neutral or alkaline soils.

Severe Drought Conditions

Do not graze forage or cut forage for hay. Under severe drought conditions, the Conservation Reserve Program allows grazing and making of hay from CRP acres, as so specified by the local ASCS (Agricultural Stabilization & Conservation Service) office. This label does not allow under any circumstances grazing or making of hay from CRP acres that have been treated with atrazine.

SUGARCANE

General Use Directions for All States

For control of many broadleaf and grass weeds, including amaranths, crabgrass, fireweed, Flora's paintbrush, foxtails, junglerice and wiregrass, broadcast 4-8 pts./A of this product at time of planting or ratooning, but before sugarcane emerges. Broadcast aerially in a minimum of 5 gals. of spray per acre, or broadcast or band by ground equipment in a minimum of 20 gals. per acre, unless otherwise indicated. One additional application may be made over the sugarcane as it emerges, and two additional applications may be made interline after emergence as directed sprays. Repeat treatments, where needed, may be applied broadcast, band or interline as suggested with the final application being prior to close-in. Do not exceed the rate of herbicide suggested for any one crop of sugarcane. **Note:** Where high rates of this product are used alone, apply in a minimum of 1 qt. of water for each 1 qt. of this product applied per acre.

Aerial Application: To assure that the spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low-drift nozzles at a maximum pressure of 40 psi and restrict application to periods when wind speed does not exceed 10 mph. To assure that drift will not adversely affect adjacent sensitive nontarget plants, apply this product alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

Precautions for All States and Uses:

1. Injury to sugarcane may occur when under moisture stress, when soil is of low adsorptive capacity or when land is first cropped to sugarcane.
2. Do not apply after close-in.
3. Do not apply more than 20 pints per acre to any one crop of sugarcane or crop injury may result.

For specific weed problems the following directions may be used. Other rate and application timings may be used for other weed spectrums and cultural practices, provided they are within the above *General Use Directions for All States* and are consistent with the *Precautions for All States and Uses*.

Florida

For control of emerged pellitory weed: Apply 0.8-1.2 pts./A in at least 40 gals. of water as a directed spray by ground equipment prior to close-in. Add 4 qts. of surfactant for each 100 gals. of spray. Thoroughly cover weed foliage.

For control of alexandergrass, large crabgrass, pellitory (artillery) weed and spiny amaranth, use one of the following methods at planting or ratooning:

1. Apply 8 pts./A preemergence. Follow with one or two applications, as needed, postemergence to sugarcane and weeds, at 4 pts./A. Treat before weeds exceed 1.5 inches in height.
2. Apply 1 to 3 times, as needed, at 4 pts./A postemergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

Louisiana

For control of annual weeds during the summer fallow period, apply 4 pints per acre to weed-free beds immediately after bed formation. Follow normal weed control program after planting. **Precautions:** To avoid crop injury:

1. Do not apply more than 10 quarts per acre to any one crop of sugarcane.
2. If making 4 pints per acre application during summer fallow period, do not exceed 8 quarts per acre during the remainder of the growing season or illegal residues may result.

Texas

Use this product for control of barnyardgrass, pigweed, purslane and sunflower in plant or ratoon sugarcane. Apply 8 pints per acre of this product preemergence. Follow with 1 or 2 applications, as needed, at 6 pints per acre postemergence to sugarcane and weeds. For best results when weeds are emerged, add a nonionic surfactant at a concentration of 2 quarts per 100 gallons to the spray and apply before weeds exceed 1.5 inches in height.

TURFGRASSES FOR SOD (Florida Only)

St. Augustinegrass, Centipedegrass and Zoysiagrass

Broadcast 4-8 pts./A according to soil texture to control those weeds listed under *Gladezine 4L Applied Alone – Corn or Grain Sorghum: Preplant Surface-Applied, Preplant Incorporated or Preemergence*.

Muck or peat	8 pts.	Old beds: within 2 days after lifting sod
		New beds: 3-4 days after sprigging or plugging
Sandy soil	4 pts.	Old beds: Within 2 days after lifting sod
		New beds: 7-10 days after sprigging or plugging

If weeds regrow, apply an additional 4 pts./A on a muck or peat, or 2 pts./A on sandy soil.

Precautions: To avoid crop injury,

1. Do not apply within 30 days prior to cutting or lifting.
2. Do not apply in combination with surfactants or other spray additives.
3. Use only on turfgrass reasonably free of infestations of insects, nematodes and diseases.
4. On newly sprigged turfgrass, temporary slowing of growth may follow application.

TURFGRASS FOR FAIRWAYS, LAWNS, SOD PRODUCTION* AND SIMILAR AREAS

*In states other than FL. For use on turfgrass for sod in FL, see *Turfgrasses for Sod (Florida Only) section*.

Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass

Apply this product after October 1 before emergence of winter annual weeds for control of annual bluegrass, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover and spurweed. This product will control annual bluegrass even if it is emerged at time of treatment. For control of summer annual weeds listed in the preemergence section of the **Atrazine 4L Applied Alone – Corn or Grain Sorghum** section of this label, also apply this product in late winter before the weeds emerge. Apply in a minimum of 15 gallons of water per acre or 1 gallon per 1,000 square feet.

Where annual bluegrass is the major weed use 1 quart per acre (22 ml or 0.75 fl. oz./1,000 sq. ft.). Use 2 quarts per acre (44 ml or 1.5 fl. oz./1,000 sq. ft.) for control of the other weeds named above. Do not exceed 1 quart per acre per treatment on newly sprigged turfgrass or on hybrid bermudagrass, such as Tiflawn, Tifway and Ormond. For continued summer annual weed control apply another 1 quart per acre at least 30 days after the previous application, but not after April 15. However, do not make more than two applications of this product per year.

Precautions: On newly sprigged turfgrass and hybrid bermudagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury:

1. Use only on turfgrass reasonably free of infestations of insects, nematodes and diseases.
2. Do not use on golf greens.
3. Do not use north of NC (except may be used in VA Coastal Plains) or west of the high rainfall areas of eastern OK and eastern TX.
4. Do not use on muck or alkaline soils.
5. Do not apply over the rooting area of trees or ornamentals not listed on this label.
6. Do not overseed with desirable turfgrass within 4 months before or 6 months after treatment.
7. Do not apply this product to newly seeded bermudagrass until it has overwintered and has a well-developed rhizome system. Do not exceed 2 quarts of product per acre within 12 months of seeding bermudagrass.

Note: Do not graze or feed turf clippings to animals or illegal residues may result.

MACADAMIA NUTS

For preemergence control of many broadleaf and grass weeds, including crabgrass, foxtail, wiregrass, Flora's paintbrush, spanishneedles and fireweed, broadcast 4 to 8 pints per acre before harvest and before weeds emerge. Repeat as necessary. Do not spray when nuts are on ground during harvest period. Do not apply by air.

GUAVA

Use only on established plantings that are at least 18 months old. Apply as a directed spray at 4 to 8 pints per acre of this product in 20 to 50 gallons of spray mix preemergence or early postemergence to weeds. When applying postemergence the use of a surfactant and greater spray volume (80 to 100 gallons of spray mix per acre) may enhance weed control. This product controls many annual broadleaf and grass weeds, including fireweed, purslane, scarlet pimpernel, spanishneedles and sowthistle.

Notes: To avoid illegal residues,

1. Do not allow spray to contact foliage or fruit.
2. Do not apply more frequently than at 4-month intervals.
3. Do not apply more than 16 pints of this product per year.

CONIFERS

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established conifers (including Douglas fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, lodgepole pine (shore pine), monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce and Sitka spruce): Broadcast 4 to 8 pints in a minimum of 5 gallons of water per acre by air or 10 gallons by ground before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are dormant. For new transplants, apply

during or soon after transplanting. For applications prior to transplanting, allow sufficient precipitation to activate this product before transplanting. In areas where spring and summer rainfall is inadequate to activate this product, apply during fall prior to spring transplanting.

For the list of weeds controlled see *Atrazine 4L Applied Alone – Corn or Grain Sorghum – Preplant Surface-Applied, Preplant Incorporated or Preemergence*.

Quackgrass Control: Broadcast 8 pints in a minimum of 5 gallons of water per acre by air or 10 gallons by ground between fall and early spring while trees are dormant and before quackgrass is more than 1.5 inches tall.

Precautions:

1. In areas west of the Rocky Mountains (except the Great Basin) grazing may begin 7 months after a fall application or 3 months after a winter or spring application.
2. To prevent illegal residues do not graze treated areas of the Great Basin or areas east of the Rocky Mountains.
3. Temporary injury to trees may occur following use of this product on coarse-textured soil.
4. To avoid crop injury do not apply to seedbeds.
5. Apply only once per year.

Aerial Application: To assure that the spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 feet above vegetation using low-drift nozzles at a maximum pressure of 40 psi and restrict application to periods when the wind speed does not exceed 10 mph. To assure that the drift will not adversely affect adjacent sensitive nontarget plants apply this product by aircraft at a minimum upwind distance of 400 feet from sensitive plants.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal, or cleaning of equipment.

Storage: Ground water contamination may be reduced by diking and flooring or permanent liquid bulk storage sites with an impermeable material. **Pesticide Disposal:** Open dumping is prohibited. Wastes resulting from the use of this product are acutely toxic. Improper disposal of unused pesticide, spray mixture or rinsate is a violation of Federal law. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, state or local procedures. For guidance in proper disposal methods, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office. **Container Disposal (2.5 gallons):** Do not reuse empty container. Triple rinse (or equivalent), puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke. **Container Disposal (Bulk, 100 and 200 gallons):** Refer to label on container for disposal instructions.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency call (919) 292-7100 day or night.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Do not breathe vapors or spray mist. Avoid contact with eyes, skin or clothing.

First Aid

If Swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Note to Physician: There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous activated charcoal slurry may be considered.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. [You may also contact 1-800-xxx-xxxx for emergency medical treatment information.]

{Note: The first aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Personal Protective Equipment (PPE)

Applicators and other handlers (other than mixers and loaders) must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear plus socks.

Mixers and loaders must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells and sink holes. Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading sites.

States may have in effect additional requirements regarding well-head setbacks and operational area containment.

This product may not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be applied aerially or by ground within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66-foot buffer or setback from runoff entry points must be planted to crop, seeded with grass or other suitable crop.

Tile-Terraced Fields Containing Standpipes: To ensure protection of surface water from runoff through standpipes with tile-outlets in terraced fields, one of the following options may be used:

1. Do not apply this product within 66 feet of standpipes in tile-outletted terraced fields.
2. Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2 to 3 inches in the entire tile-outletted terraced field.
3. Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management practice is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Caliber 90[®], Concep[®], Dual[®] and Princep[®] trademarks of Syngenta Crop Protection
 Bexton[®] trademark of Dow AgroSciences
 Compex[®] trademark of KALO Agricultural Chemicals, Inc.
 Gramoxone[®] and Sutan+[®] trademarks of Aventis Environmental Science
 Lasso[®], Ramrod[®] and Roundup[®] trademarks of Monsanto Company
 X-77[®] trademark of Loveland Industries, Inc.
 Unite[®] trademark of HACO, Inc.

MARMAN USA, INC.
 P.O. Box 22829; Tampa, FL 33622-2829

[] Denotes alternate/optional language
 { } Denotes language that does not appear on market labeling

tc

{Container Label – For 2.5 gallons, Bulk, 100 gallons and 200 gallons containers}

RESTRICTED USE PESTICIDE
(Ground and Surface Water Concerns)
For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification. This product is a restricted-use herbicide due to ground and surface water concerns. Users must read and follow all precautionary statements and instructions for use in order to minimize potential for atrazine to reach ground and surface water.

ATRAZINE 4L HERBICIDE

For Season-Long Weed Control in Corn, Sorghum and Certain Other Crops

Active Ingredients:

Atrazine: 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	40.8%
Related compounds	2.2%
Other Ingredients	57.0%
Total	100.0%

Contains 4 pounds active ingredients per gallon.

Shake well before using.

See Directions for Use in Attached Booklet

Keep Out of Reach of Children
CAUTION
See additional precautionary statements below.

_____ Gallons
U.S. Standard Measure

Agricultural Use Requirements
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. Refer to supplemental labeling under *Agricultural Use Requirements* in the *Directions for Use* section for information about this standard.

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Do not breathe vapors or spray mist. Avoid contact with eyes, skin or clothing.

First Aid

If Swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Immediately rinse skin with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Note to Physician: There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous activated charcoal slurry may be considered.

Have the product container or label with you when call a poison control center or doctor, or going for treatment. [You may also call 1-800-xxx-xxxx for emergency medical treatment information.]

{Note: The first aid statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.}

Environmental Hazards

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells and sink holes*. This product may not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be applied aerially or by ground within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66-foot buffer or setback from runoff entry points must be planted to crop, seeded with grass or other suitable crop.

*For exceptions to this restriction see the *Environmental Hazards* section of the *Precautionary Statements* in the attached booklet.

Tile-Terraced Fields Containing Standpipes: To ensure protection of surface water from runoff through standpipes with tile-outlets in terraced fields, one of the following options may be used:

4. Do not apply this product within 66 feet of standpipes in tile-outletted terraced fields.
5. Apply this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2 to 3 inches in the entire tile-outletted terraced field.
6. Apply this product to the entire tile-outletted terraced field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management practice is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Chemigation Prohibition

Do not apply this product through any type of irrigation system.

Container Disposal

(To be used in accordance with Directions for Use in the Atrazine 4L label booklet.

See the Atrazine 4L label booklet for pesticide disposal instructions.)

2.5 Gallons: Do not reuse empty container. Triple rinse (or equivalent), puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke.

Bulk: Thoroughly clean container before reuse, unless this is a Marman refillable container. **If this is a Marman refillable container, do not rinse or introduce any pesticide other than Atrazine 4L.** Reseal and return the container to an authorized Marman refilling facility. If the Marman container is not to be refilled, triple rinse (or equivalent) and offer for recycling or reconditioning, or by open burning, if allowed by state and local authorities. If burned, keep out of smoke

100 and 200 Gallons: This is a refillable container. **If the container is to be refilled, do not rinse with any material or introduce any pesticide other than Atrazine 4L.** Reseal and return the container to an authorized Marman refilling facility. If the container is not to be refilled, triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or by open burning, if allowed by state and local authorities. If burned keep out of smoke.

For minor spills, leaks, etc. follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the even of a major spill, fire or other emergency call (919) 292-7100 day or night.

EPA Reg. No. 48273-11 EPA Est. _____

MARMAN USA, INC.
P.O. Box 22829; Tampa, FL 33622-2829

[] Denotes alternate/optional language

{ } Denotes language that does not appear on the market label