

BEST QUALITY AVAILABLE



MAJESTIC GREEN

ROTENONE plus COPPER DUST

A Combination Insecticide and Fungicide Dust For Home Garden Use

EARLY BLIGHT - LATE BLIGHT DOWNY MILDEW APHIDS, FLEA BEETLES, STRIPED AND SPOTTED CUCUMBER BEETLES

Table with 2 columns: Ingredient Name and Percentage. Includes Rotenone (75%), Other Cube Extractives (1.20%), Copper (Basic Copper Sulphate) expressed as metallic (5.00%), and Inert Ingredients (93.05%). Total is 100.00%.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash thoroughly with soap & water.

IF IN EYES: Wash out with plenty of water.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 48391-6
EPA Est. No. 10107 - NB 1

NET WEIGHT 5 POUNDS

Manufactured For
Majestic Green Products, Inc.
P.O. Box 1064
Stoughton, MA 02157

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
Irritant if inhaled or ingested. Do not get into eyes, on tender parts of body or clothing. In case of contact wash thoroughly with soap and water. If it gets into eyes wash out eyes at once with plenty of water. Avoid breathing dust.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment, or disposal of wastes.
PHYSICAL OR CHEMICAL HAZARDS
Keep this package away from excessive heat. Avoid storage near steam pipes and radiators.

DIRECTIONS FOR USE
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE AND DISPOSAL

STORAGE - Keep this package away from excessive heat. Avoid storage near steam pipes and radiators.
DISPOSAL - Do not reuse empty container. Bury in approved landfill area or incinerate. Do not burn. For more information on consumer packing, visit our website at www.majesticgreen.com.

DIRECTIONS (CONT'D)

Best results will be obtained if Majestic Rotenone plus Copper Dust is applied when first signs of disease or insects appear. Dust the plants thoroughly, applying to top as well as underside of leaves. Repeat applications every 4 to 6 days or as required.

In conventional areas apply at the rate of 20 to 25 lbs. per acre with hand or ground equipment or at the rate of 35 to 40 lbs. per acre with airplane equipment. For the home garden dust lightly by punching out the perforated top in the 1 lb. can or applying with a hand duster from larger containers.

For the control of early blight, late blight, spruce, flea beetle and Colorado potato beetle on potatoes, separate pests on asparagus, early blight, late blight, Septoria leaf spot and flea beetle on tomatoes; downy mildew, anthracnose, striped and spotted cucumber beetles on cucurbits (pumpkins, cucumbers, melons and squash); downy mildew, anthracnose, tyrus bug, Mexican bean beetle and bean leaf beetle on beans; downy mildew, leaf spot, cabbage looper, flea beetle and imported cabbage worm on cabbage; early blight, late blight and leaf spot on celery; flea beetle on beets and radishes.

ACCEPTED
NOV 10 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 48391-6

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NOTE: This product sold by weight. Package is full when packed, but due to natural settling of contents may be found to settle when shipped.