

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MAY IRRITATE EYES, NOSE, THROAT, AND SKIN. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. FIRST AID: In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions shift from areas treated.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire or sparks.



**MAJESTIC
GREEN**

BORDEAUX MIXTURE

For Disease Control on Peonies,
Evergreens, Shrubs, Flowers,
Fruit and Shade Trees

ACTIVE INGREDIENT:

Copper, expressed as metallic,
[in form of Bordeaux Mixture]..... 12.75%

INERT INGREDIENTS..... 87.25%

TOTAL 100.00%

STORAGE AND DISPOSAL

Do not reuse container; dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

DEC 1 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

48391-13

CAUTION:
Keep Out of Reach of Children.
SEE SIDE PANELS FOR ADDITIONAL CAUTIONS

EPA Reg. No. 48391-
EPA Est. No. 10187-NB-1

Mfg. For:
Majestic Green Products Inc.
P.O. Box 1064
Manhattan, KS 66502

Net Contents: 3 Lbs.

DIRECTIONS FOR USE

(Shake well before using)

Dry BORDEAUX MIXTURE is made from equal parts of Copper Sulfate and Lime. Each pound of this product mixed with 50 gallons of water is equivalent to a 4-4-10 BORDEAUX MIXTURE. The strength in percentages referred to is 4-4-10.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DIRECTIONS:

APPLES: To control late blight, black rot, bitter rot and scab, use 4 TBS/gal. spray 4 TBS of hydrated lime. May repeat certain varieties.
PEARS: For blight, use 4 TBS/gal. in full bloom. Do not add insecticide.
APRICOTS, PLUMS: Brown rot and leaf spot, use 4 TBS per gallon plus equal amount of lime. Do not use on Japanese plums. Use at about split stage and in first and second greening stages.
CHERRIES: To control Leaf Spot, use 3 TBS/gal. of water or 3 lbs. to 50 gal. water. On Sweet Cherries it is advisable to add 4 lbs. hydrated lime spray as soon as the petals fall and again one week later, plus two additional subsequent applications at 10-14 day intervals. The 4th application after fruit is harvested at 14 day intervals.
PEACH: Leaf curl, use 12 TBS/gal. in early spring. Repeat late fall after leaf drop.
PECAN: Scab, use 6 TBS/gal. when cutting above. Repeat 4 times at 3 week intervals. Spanish moss, use 12 TBS/gal. spray during winter.
GRAPES: Black rot and mildew, use 6 TBS/gal. when shoots are 18 inches long. Repeat after bloom and 10 days later, again in mid-August.
POTATOES: Early and late blight, use 6 TBS/gal. Apply when plants are 3 to 4 inches high, maintain a 10 to 14 day schedule for entire season.
STRAWBERRY: Leaf spot, use 6 TBS/gal. at 10 to 14 day intervals, except during harvest. Final spray in late summer.
CITRUS: Lemon berry rot, grapefruit scab, rosette, use 6 TBS/gal. Spray just before spring flush. Repeat 1 to 3 weeks after bloom.
TOMATO: Early and late blight, Septoria leaf spot, use 6 TBS/gal. Repeat at 7 to 10 day intervals. Alternate with Zinc 75W if Anthracnose is a problem.
CELERY: To control early and late blight, Leaf Spot use 4.5 TBS/gal. or 4.5 lbs. to 50 gal. water. Spraying or drenching plants in plant bed and at 10 day intervals in the field.
MELONS: To control Anthracnose, Scab, Leaf Spot, use 3 TBS/gal. or 3 lbs. to 50 gal. of water. Spray as soon as plants appear and repeat at 5 day intervals during rising.
SYCAMORES, LINDEN, MAPLE, OAK, TULIP TREE, ELM: Control of Anthracnose, Leaf Spot and Elm Leaf Curl, use 6 TBS/gal. as leaves appear in spring. Repeat in 3 to 3 weeks.
JUNIPER, CEDAR, CYPRESS, ARBORVITAE, SPRUCE, PINE, YEW: For Needle Cast, Sphaeria blight, Twig blight, Cytospora blight apply 1 to 2 TBS/gal. and spray using 1 TBS/gal. Spray Cedar and Juniper July-August for Cedar Apple Rust.
IVY: Leaf Spot, use 6 TBS/gal. in spring. Repeat in 14 days or at first sign of disease.
LAUREL, LILAC, RHODODENDRON: Leaf Spot and blight, use 4 TBS/gal. in early spring. Again after bloom.
DOGWOOD, BOXWOOD, CAMELLIAS, Daphne, Leaf Spot and Nectria Canker, use 6 TBS/gal. as growth starts in early spring.
DANGLIAS, TULIPS, GLADIOLUS, LILIES AND IRIS: For Leaf Spot and Nectria blight spray weekly, use 6 TBS/gal. in early spring. Repeat in 7 to 10 days.
PEONY: Leaf blight, Botrytis blight (gray mold), use 6 TBS/gal. Spray before shoots are one foot tall, repeat in two weeks. If any sign of bud blast, spray again. Remove any wilted shoots by cutting below ground. Place in paper bag and burn. Remove old masts and leaves in fall and burn - do not compost.
DELPHINIUM, GERANIUM: Leaf Spot, use 4 TBS/gal. Spray at first sign of disease.
PALMS: Anthracnose, Scab, Leaf Spot, use 6 TBS/gal. Spray at first sign of disease.
MIXING TABLE:
Use level spoons in measuring, 2 tablespoons equal 1 tablespoonful. Use higher dosage if one has been shown to kill the pest.

Amount of Spray Desired	Amount of Bordeaux Mixture Required 4-4-10	4-4-10
1 Pint	1/4 to 1/2 tablespoon	1 to 1 1/2 tablespoons
1 Quart	1/4 to 1/2 gallon, 200	2 to 3 tablespoons
1 Gallon	6 to 7 tablespoons	8 to 12 tablespoons
2 Gallons	1 1/2 to 1 3/4 cups	16 to 24 cups

NOTICE

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Plant injury, ineffectiveness, or other untoward consequences may result because of such factors as weather conditions, presence of other materials, or the use of application of the product contrary to label instructions, all of which are beyond the control of the manufacturer. All such risks shall be assumed by the user.

Seller warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to in this label.

Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages such as loss of profits or value or any other special or indirect damages.

Seller makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY.

46371-13

-2-

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure