

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

STATEMENT OF PRACTICAL TREATMENT

Danger Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. May be fatal if swallowed. Avoid breathing vapors. If in eyes flush with plenty of water and get medical attention. If on skin wash with plenty of soap and water. Get medical attention if irritation persists. If swallowed drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

Environmental Hazard

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by the cleaning of equipment or disposal of wastes. Apply this product only as directed on label.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Pesticide Disposal

Pesticide spray mixture or residue that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

Container Disposal

Metal - Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary land fill or by other approved State and local procedures.

Plastic - Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary land fill or by incineration if permitted by State and local authorities.

IMPORTANT

Do not apply on or near desirable trees or plants or in locations where chemicals may be washed or misted in contact with their roots.


Avoid drift of spray (coarse sprays are less likely to drift) onto vegetables, flowers, ornamental shrubs, trees and other desirable crop plants.

Do not contaminate domestic or irrigation waters.

Do not pour solution used to clean sprayer into desirable plants or where it may contaminate water.

Do not store in direct contact with fertilizers, insecticides or fungicides.

Do not use on centipede grass, St. Augustine, or buffalo grass on lawns or turf where bent grass or clovers are present. Newly seeded lawns may be treated after they have been mowed the second time. To reseed straw areas in treated lawns, wait at least 3 weeks after spraying. A good rain or irrigation should follow during this period.



**MAJESTIC
GREEN**

BANVEL® + 2, 4-D

(Contains 1.5 Pounds Dicamba and 2.9** Pounds 2, 4-D per gallon)

ACTIVE INGREDIENT

Dimethylamine salt of dicamba 1.5 lb.
Dimethylamine salt of 2, 4-D 2.9 lb.
Dimethylamine salt of 2, 4-D isomer (phenoxycetic acid) 0.2 lb.

INERT INGREDIENTS

TOTAL 5.6 lb.

*Equivalent to 1.5 lb. of dicamba per gallon of water.
**Equivalent to 2.9 lb. of 2, 4-D per gallon of water.

†Equivalent to 0.2 lb. of 2, 4-D isomer (phenoxycetic acid) per gallon of water.
Banvel is a Registered Trademark of Velsicol Chemical Corporation.

DANGER

KEEP OUT OF REACH OF CHILDREN

See back panel for additional caution statements.

EPA Reg. No: 43891-

EPA Est. No: 10167-NB-1

MANUFACTURED For
Majestic Green Products Inc.
P.O. Box 1064
Manhattan, KS 66502

IMPORTANT

Avoid drift of spray onto desirable flowers, ornamental shrubs, trees and other desirable crop plants. Coarse sprays are less likely to drift. Do not apply within the drip line of trees or shrubs. Do not contaminate domestic or irrigation waters. Do not pour solution used to clean sprayer into desirable trees or shrubs or where it may contaminate water. Do not store in direct contact with fertilizers, insecticides or fungicides. Do not use on centipede grass, St. Augustine grass or buffalo grass on lawns or turf where bent grass or clovers are present. Do not use on newly seeded lawns or turf. When container is empty immediately wash thoroughly and destroy. Never reuse the spray equipment for applying 2, 4-D herbicides on any turf.

Read LIMIT OF WARRANTY AND LIABILITY before buying or using. If terms are not acceptable, return at once unopened.

LIMIT OF WARRANTY AND LIABILITY

The exclusiveness of the user or buyer and the limit of the liability of this company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. There is no warranty and this company and any other seller disclaim all liability for all losses, personal injuries or damages arising from combining substances and from any use of this product not recommended in label directions or arising from misuse, improper handling or improper storage.

DIRECTIONS

For use on lawns, turf, hot pastures and fairways, aprons, tees and rough.

USE	WEEDS	DOSAGE	APPLICATION
On lawns, turf, hot pastures, aprons, tees and rough.	dandelion plantain curly dock common chickweed moss stachys dog fennel sheep sorrel clover crabgrass hardy pigweed knockweed ground ivy scarlet and onion hardtail and geranium poppy poppy	Apply 2 pints to 3 pints BANVEL + 2, 4-D per acre or 4 to 6 teaspoons per 1000 square feet. Equivalent to 34 lb. Dicamba + 68 lb. 2, 4-D to 51 lb. Dicamba + 102 lb. 2, 4-D per acre.	Apply as a foliar spray to actively growing weeds with 1 to 2 gallons of water per 1000 sq. ft. and 40 to 250 gallons of water per acre to give good coverage. Fast germinating weeds may be controlled more effectively when treated in the fall or early winter. Make 1 or 2 applications per year.
On golf courses, fairways, aprons, tees and rough.	dandelion plantain curly dock common chickweed moss stachys dog fennel sheep sorrel clover crabgrass hardy pigweed knockweed ground ivy scarlet and onion hardtail and geranium poppy poppy	Apply 2 pints to 3 pints BANVEL + 2, 4-D per acre or 4 to 6 teaspoons per 1000 square feet. Equivalent to 34 lb. Dicamba + 68 lb. 2, 4-D to 51 lb. Dicamba + 102 lb. 2, 4-D per acre.	Apply as a foliar spray to actively growing weeds with 1 to 2 gallons of water per 1000 sq. ft. and 40 to 250 gallons of water per acre to give good coverage. Fast germinating weeds may be controlled more effectively when treated in the fall or early winter. Make 1 or 2 applications per year.