

**Precautionary Statements  
Hazards to Humans and Domestic Animals**

**CAUTION**

Use adequate ventilation and avoid breathing dust. Wear a suitable dust mask, i.e. NIOSH/MSHA TC-21C-132, when using this product during prolonged exposure.

**INSECTS DIE FROM DEHYDRATION**

INSECTO is derived from diatomaceous earth refined to certain specifications. Insects that come in contact with INSECTO are lacerated by the microscopic particles and lose their body fluid (dehydration) which will cause their deaths. This is a physical control to which insects have a difficulty building up a resistance.

**STORAGE AND DISPOSAL:**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool dry area.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not reuse bag. Discard bag in trash.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**ACCEPTED**  
14 AUG 1995  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 48598-1

**INSECTO  
AN  
INSECTICIDE  
FOR CONTROL OF GRAIN INSECTS**

**AND  
HOUSE INSECTS**

**INGREDIENT STATEMENT:**

Active Ingredient	
Silicon Dioxide, from diatomaceous earth . . . . .	90.0%
Inert Ingredients . . . . .	10.0%
<b>TOTAL</b>	<b>100.0%</b>

JAPAN REG. NO. 17078  
CANADA P.C.P. NO. 22489  
SAUDI ARABIA 1379-63-63  
INDONESIA R.I. 910-1-903  
CHINA 19-04024  
NICARAGUA NA-171A-6-93  
MALAYSIA CRMP R/2850

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS  
RECOMMENDED FOR COMMERCIAL/INDUSTRIAL USE**

*Manufactured by:*

**NATURAL INSECTO PRODUCTS, INC.**  
630 North Eckhoff Street  
ORANGE, CA 92668

E.P.A. REG. NO. 48598-1-AA  
E.P.A. ESTABLISHMENT NO. 48598-CA-1

**NET WEIGHT 25 POUNDS (11.34 Kg.)**

**DIRECTIONS FOR USE:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

INSECTO is intended for application with hand or power duster, or other suitable means, to hiding and running areas and places insects are found. To apply INSECTO into cracks and crevices, use Bulbous Duster or other suitable equipment.

INSECTO is an effective insecticide for killing exposed insects Rice Weevils, Granary Weevils, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Red Flour Beetles, and Tribolium in Empty Grain Bins, Boxcars, Ships and Silos. Its use will aid in protecting Feed Grains, Seed, Stored Grains, Wheat, Barley, Buckwheat, Corn, Oats, Rice, Rye, Flax, Pea, Soybean, Peanuts, Popcorn, Cocoa Beans, and Sorghum (milo).

INSECTO should be dusted lightly, approximately one (1) pound per 1,000 square feet on all interior wall surfaces of empty storage areas. Where possible, special attention should be given to dusting cracks, crevices and other areas where insects hide or crawl. After dusting, there is no need to remove the dust or aerate the areas. It is safe to enter the area and can safely be filled with grains. It is recommended that the area be left undisturbed for two to three days to eliminate existing infestations before filling. It is desirable to redust each time the storage area is emptied.

To control accessible stages of Indian Meal Moth and Angoumois Moth. Apply to surface of binned grain at the rate of four (4) pounds per 1,000 square foot. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations break up webbing with a rake before dusting and make second application within two weeks.

**FOR IN-TRANSIT PROTECTION:** Clean and dust truck beds, boxcars, and ship holds before loading. Thoroughly treat walls and floors and dust lightly after container is loaded.

**TREATMENT OF STORED GRAIN:** INSECTO can safely be applied to the grain at the time of storage, at the rate of one (1) to two (2) pounds per ton of grain, to protect it from further damage. Treat grains and seed immediately after harvest.

**CAUTION:** Apply INSECTO at recommended rate. Excessive rate (application) may reduce grain to a lower grade, have reduced flowability and reduce test weight.

**PEST CONTROL AROUND THE HOUSE:** Lightly coat the area (16oz. per 1,000 Sq. Ft.) infested with Ants, Roaches, Fleas, Silver Fish, Lice, Water Bugs, and Moths.

**FOOD AND FEED AREAS:** In areas where food or feed is received, processed (including enclosed processing systems), packed (bottled, boxed, canned, wrapped), transported, stored (including waste), prepared, or served, application is limited to crack and crevice and spot treatments only. All food or feed should be removed or covered before treatment.

Apply sparingly into cracks and crevices using equipment designed to place powder in specific locations. Treat open spaces below floors, above ceilings, within walls, under stairs, underneath and behind equipment or appliances, around pipes, under sinks, in utility closets, around garbage cans, etc.

No individual spot will exceed two (2) square feet. Limit spot treatments to wall or floor surfaces where pests have been seen or where pests are suspected of hiding.

**NON-FOOD, FEED AREA:** Use crack and crevice and spot treatments as described above. Also apply to other interior spaces where pests hide, run or land, such as attics and spaces above suspended ceilings, garages, porches, utility rooms, storage rooms, basements and crawl spaces, cracks in baseboards, moldings and doorframes, corners in closets, cabinets and shelves, windowsills and frames, under and behind furniture and appliances, etc. Work into and apply under carpets, rugs, doormats, couches, and other furniture. Wash bedding for pest and apply dust to dry bedding and surrounding areas. Apply to mattresses and bed frames. Outdoors, apply to window and door frames, covered entrances and porches, and along foundations where dust will remain dry.