


PRECAUTIONARY STATEMENTS

HARMFUL TO HUMANS & DOMESTIC ANIMALS

POISON



Fatal if inhaled. Corrosive, causes irreversible eye damage. May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, or on skin. Do not breathe dust, vapor, or spray mist. When handling either products containing, EPTA or spray-diluted mixtures, wear protective clothing (long pants, long sleeve shirt, impermeable gloves, boots, and a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration and the National Institute for Occupational Safety and Health. When handling the concentrated products, wear a face shield. Wash thoroughly with soap and water after handling and before eating, or smoking. Remove contaminated clothing and wash before reuse. Do not enter treated areas for at least 24 hours after treatment.

The United States Environmental Protection Agency has determined that triphenyltin hydroxide causes birth defects in laboratory animals. Exposure to triphenyltin hydroxide during pregnancy should be avoided.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water or wetlands. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Cover or incorporate spills. Do not contaminate water by cleaning, or equipment or disposal of wastes.

Directions For Use

BEST AVAILABLE COPY

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

WASTES AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinse is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container disposal: Triple rinse (or equivalent). Then offer for recycling, or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

4-11a fungicide can be applied as a ground or aerial spray to control fungicidal infestations on listed crops. Application rates are for general use. The state agricultural extension or agricultural experiment station specialists should be consulted for specific applications and timing recommendations. With ground spray application thorough coverage is essential for good control.

Center Panel

"RESTRICTED USE PESTICIDE: Because of the high acute toxicity of triphenyltin hydroxide, and its potential for creating teratogenic effects, this product may be applied only by certified applicators or persons directly under their supervision."

4-TIN

Flowable Fungicide

Triphenyltin hydroxide Flowable Fungicide

ACTIVE INGREDIENT:

Triphenyltin hydroxide.....40.0%

INERT INGREDIENTS:.....60.0%

Total.....100.0%

This product contains 4 pounds of Triphenyltin hydroxide per gallon.

KEEP OUT OF REACH OF CHILDREN

DANGEROUS POISON

STATEMENT OF FIRST AID TREATMENT

If on skin: Wash with plenty of soap and water.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If in eyes: Flush with plenty of water. Call a physician.

If swallowed: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician or Poison Control Center.

See Side Panel For Additional Precautionary Statements.

Mark Corporation
Valdosta, Georgia 31601

EPA REG. NO. 49085-1
EPA EST. NO.

Net Contents: 2.7 gallons

BEST AVAILABLE COPY

Directions for Use:

Use this product only as specified on this label.

Application:

Use of surfactants, spreaders or stickers. Do not graze dairy or meat animals in treated areas.

Directions for Use: FOR APPLICATION:

4-TIN 3.13: Cercospora Leafspot

Ground application, apply 3.8-9.5 fl. ounces of 4-TIN per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. Aerial application, (helicopter or fixed wing aircraft), apply 3.8-9.5 fl. ounces of 4-TIN per acre in 5-10 gallons of water. Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infection periods. Application should begin when leafspot conditions appear or when the disease is in the area and repeated at 10 to 14 day intervals. Do not treat within 14 days of harvest. Do not graze or feed beet tops to livestock.

4-TIN 0.13: Early and Late Blight

Ground application, apply 4.7-9.5 fl. ounces of 4-TIN per acre in at least 15 gallons of water. Full coverage of the foliage is necessary for best results. A minimum rate of less than 200 pounds is recommended. Aerial application, apply 4.7-9.5 fl. ounces per acre in 3-10 gallons of water. For helicopter application, fly high enough so as not to whip the vines. Diluted spray should be directed uniformly to all parts of the plant and the gallonage increased according to the size of the plants. Application should begin with the appearance of blight or early conditions and continue on a 7 day schedule. Do not treat within 7 days of harvest when applied by ground or aerial application equipment. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area. IRRIGATION SYSTEM (Solid-set or center-pivot system only) - Apply 5.5-9.5 fl. ounces per acre. Do not apply through center-pivot (circle) system which uses water pressure for movement of system. Irrigation system must be equipped with safety valves or other devices to prevent backsiphoning of pesticide into water source. Irrigation water treated with this product should be held on the treated area until it is absorbed by the soil. It should not be turned into the tail water or fed back into the irrigation system. For complete directions regarding application, refer below. Application should begin with the appearance of blight or early conditions and continue on a 7 day schedule. Do not treat within 7 days of harvest when applied by ground or aerial application equipment and not within 33 days when applied by irrigation system. The lower rate of application should be used early in the season and the high rate mid to late season or when blight infection is in the area.

BEST AVAILABLE COPY

4-TIN 3.13: Cercospora Leafspot and Alternaria Blight

Apply 3.8-7. fl. ounces per acre. Use lower rate when diseases are light, increase rate when disease pressure increases. Apply in sufficient water for uniform, full coverage of foliage 25-100 gallons of water per acre. Use lower volumes in early season when plants are small and increase spray volume as plant size increases. Applications should begin 6 weeks after planting or as soon as first signs of disease appear, spray applications should continue at 7 day intervals or as needed for continued control. Do not apply within 14 days of harvest. Do not use cobs as feed for livestock.

4-TIN 3.13: Anthracnose, Brown Leafspot, Downy Spot, Powdery Mildew, Liver Spot and Leaf Blight. Aerial application, apply 11.4-22.8 fl. ounces per acre in a minimum of 10 gallons of water. Ground application, mix 4-9 fl. ounces of 4-TIN for each

100 gallons of water, 15.2-45.0 fl. ounces per 500 gallons tank and apply as a full coverage spray. Diluted spray should be directed to all parts of the tree. Applications should begin at pre-pollination stages when the young leaves are unfolding, and a second application made when the small nuts are forming. Repeat spray applications at 2-4 week intervals as needed to maintain control. Use the lower rate for the first two applications or until the disease becomes severe, or during dry weather. Use the higher rate during wet weather or during severe scab, powdery mildew, or other disease infections. Do not apply after the shucks have started to open.

WARRANTY STATEMENT

Seller makes no warranty, expressed or implied, concerning the use of the product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.