

Bristol Chemical, Inc.

HERBAN®

WETTABLE POWDER (WP)
SELECTIVE HERBICIDE

For Weed Control In Cotton, Soybeans, Sorghum,
Sugar Cane, Spinach and Commercial Nurseries.

ACTIVE INGREDIENT:	PERCENT BY WEIGHT
Norea [3-hexahydro-4,7-methano-1H-cyclohepta-a-1,4-dihydro-5-oxo-1,2-dihydro-4,7-dimethylurea]	76
Related Reaction Products	4
INERT INGREDIENTS:	20
TOTAL	100%

CAUTION!
KEEP OUT OF THE REACH OF CHILDREN

Read left panel for additional handling precautions

EPA REG. NO.

MANUFACTURED FOR

Bristol Chemical, Inc.

Virginia Beach, Virginia 23451

NET CONTENTS

10 POUNDS NET WT. WETTABLE

REST DOCUMENT AVAILABLE

EQUIPMENT

Herban-AMBER may be applied by hand or mechanical methods. Hand application may be made by hand spraying or by brush or roller application. Mechanical methods include ground boom sprayers, aerial sprayers, and self-propelled sprayers.

CAUTION: Do not apply by airblast sprayer or by ground boom sprayer.

POST-BLOOM POST-EMERGENCE WEED CONTROL IN COTTON

DIRECTED SPRAY APPLICATIONS Directed spray applications are recommended for post-bloom post-emergence weed control in cotton. This method involves spraying only the weeds in the field, leaving the cotton plants unharmed. It is important to ensure that the spray pattern covers the weeds effectively while avoiding the cotton plants.

PRE-EMERGENCE WEED CONTROL IN SORGHUM

Herban-AMBER may be used for pre-emergence weed control in sorghum. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to prevent weeds from growing and competing with the sorghum plants.

POST-EMERGENCE WEED CONTROL IN SORGHUM

Herban-AMBER may also be used for post-emergence weed control in sorghum. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to control weeds that have already emerged and are competing with the sorghum plants.

CAUTION:

Do not apply by airblast sprayer or by ground boom sprayer.

NOTICE:

Do not apply to sorghum until it has reached the V6 stage of growth.

CONTROL FOR COTTON

Herban-AMBER may be used for control of cotton. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to control weeds that are competing with the cotton plants. It is important to ensure that the spray pattern covers the weeds effectively while avoiding the cotton plants.

IRRIGATION

Herban-AMBER may be used for irrigation. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to control weeds that are competing with the cotton plants. It is important to ensure that the spray pattern covers the weeds effectively while avoiding the cotton plants.

REGISTRATION NUMBER

Herban-AMBER is registered under the name of "Herban-AMBER". It is a water-soluble herbicide used for weed control in various crops.

WEED CONTROL IN SUGAR CANE IN LOUISIANA AND PUERTO RICO

Herban-AMBER may be used for weed control in sugar cane in Louisiana and Puerto Rico. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to control weeds that are competing with the sugar cane plants.

PRE-EMERGENCE WEED CONTROL IN SOYBEANS

Herban-AMBER may be used for pre-emergence weed control in soybeans. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to prevent weeds from growing and competing with the soybean plants.

COMBINATION OF HERBAN-AMBER IN SOYBEANS

Herban-AMBER may be used in combination with other herbicides in soybeans. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to control weeds that are competing with the soybean plants.

CAUTION: Do not apply to soybeans until the seedlings have emerged. If any damage occurs, repeat application at a lower rate. Do not apply to soybeans during the reproductive stage. Do not apply to soybeans if there is a risk of frost damage.

PRE-EMERGENCE WEED CONTROL IN SPINACH

Herban-AMBER may be used for pre-emergence weed control in spinach. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to prevent weeds from growing and competing with the spinach plants.

WEED CONTROL IN COMMERCIAL NURSERIES

Herban-AMBER may be used for weed control in commercial nurseries. It is recommended to apply the herbicide at the rate of 1.5 to 2.0 pounds per acre. This will help to control weeds that are competing with the plants in the nursery.

CAUTION: Do not apply to young plants. Do not apply to plants that are sensitive to herbicides. Do not apply to plants that are in the reproductive stage. Do not apply to plants that are in the fruiting stage. Do not apply to plants that are in the seedling stage.

NOTICE: Do not apply to plants that are sensitive to herbicides. Do not apply to plants that are in the reproductive stage. Do not apply to plants that are in the fruiting stage. Do not apply to plants that are in the seedling stage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

GASIFICATION

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

AVOID INHALATION OF DUST OR SPRAY MIST
AVOID CONTAMINATION OF FEED AND FOODSTUFFS
KEEP CLEAN AVOID CONTACT WITH SKIN
DO NOT GET IN EYES WASH UP WHEN JOB IS FINISHED

FIRST AID

If swallowed: induce vomiting by giving one (1) glass of 1 oz. of Syrup of Ipecac (for patient one year of age and older) followed by one 8 ounce glass of water. If vomiting does not occur in 15 minutes, the dose may be repeated. If syrup of Ipecac is not available, vomit spontaneously by holding the back of the hand against the chin and pushing the thumb down the patient's throat. Call a physician. In case of eye contact immediately flush eyes with plenty of water for at least 15 minutes holding the eyelids open. Get medical attention if irritation persists.

In case of skin contact wash skin with soap and plenty of water.

ENVIRONMENTAL HAZARDS

DO NOT CONTAMINATE DOMESTIC OR IRRIGATION WATER SUPPLIES BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES KEEP OUT OF LAKES, STREAMS, OR PONDS

USE PRECAUTIONS

FOLLOWING CROPS

Herban-AMBER is quickly destroyed in the soil. However, if more than 100 days elapse between application and the next crop, it is recommended that the field be plowed or tilled to bury the residue after application. Do not cultivate the field for 30 days after application.

ACCURATE DOSAGE

Use the exact amount and do not exceed the maximum amount specified for each application. Do not apply more than the maximum amount specified for each application.

PREVENT INJURY TO OTHER CROPS OR PLANTS

Do not apply Herban-AMBER directly to trees, shrubs, and other plants except as directed for specific use. Do not apply to fruit trees or fruiting crops. Do not apply to seedbeds or seedlings. Do not apply to lawns or ornamental grasses. Do not apply to aquatic plants. Do not apply to trees, shrubs, and other plants except as directed for specific use.

CLEANING SPRAYER

Do not clean sprayer with organic solvents such as gasoline, kerosene, or mineral spirits. Do not use strong acids or alkalis. Do not use sharp metal objects to clean the sprayer.