

PM 25

49517-3

177

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input checked="" type="checkbox"/> REREGISTRATION <i>(Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)</i>	EPA REGISTRATION NO. 49517-3	DATE OF ISSUANCE 1994
	TERM OF ISSUANCE Conditional	
	NAME OF PESTICIDE PRODUCT Moore Ag Brand Poly-Defoliant-Desiccant	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code) <div style="border: 1px solid black; padding: 10px; width: fit-content; margin: 0 auto;"> <p>Moore Agricultural Products 11521 Excelsior Avenue Hanford, CA 93230</p> </div>		
<p>NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.</p> <p>A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.</p> <p>Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4." 2. Add the phrase, "EPA Registration No. 49517-3" and submit five (5) copies of your final printed labeling before you release the product for shipment. 3. In order for your label to be substantially similar to the cited label you must adhere to the following labeling amendments: <ol style="list-style-type: none"> a. "Harmful or fatal if swallowed" must be added to the "Precautionary Statements". b. Under the "PPE" you must add.... <ol style="list-style-type: none"> 1. Chemical-resistant footwear plus socks 2. Chemical-resistant apron when cleaning equipment, mixing, or loading. 		
<input type="checkbox"/> ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL		DATE

75050
HICKS
3-11-94

2 of 11

- c. Update the "Environmental Hazard" statement by adding "For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark" and "Clean and wash spray equipment before and after use and do not allow washwater to contact desirable plants."
- d. Under the "Physical and Chemical Hazards" statement add "To avoid fire or explosion do not mix with insecticides".
- e. The following PPE is required under "Agricultural Use Requirements":
 - 1. Coveralls over short-sleeved shirt and short pants
 - 2. Waterproof gloves
 - 3. Chemical-resistant footwear plus socks
 - 4. Protective eyewear
 - 5. Chemical-resistant headgear for overhead exposure
- f. The title "General Information" should proceed the listing of the crops.
- g. The following statement should be included under "Soybeans":
 - 1. For ground application, apply a minimum of 20 gallons of spray solution per acre.

This acceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR Part 158, subpart K, section 156.200(c) (3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

M.A.P.CO.

**5 LB. SODIUM CHLORATE
DEFOLIANT-DESICCANT**

Active Ingredient:

Sodium Chlorate	45.0%
Inert Ingredients	55.0%
Total	100.0%

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

) Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES, Immediately flush with water for 15 minutes. Get medical attention.

IF ON SKIN, Immediately wash with plenty of soap and water. Get medical attention if irritation persists.

IF SWALLOWED, Immediately drink a large quantity of either milk, egg whites or gelatin solution. If these are not available, drink a large quantity of water. Avoid alcohol. Get medical attention.

NOTE TO PHYSICIAN:

Probable mucosal damage. Use of gastric lavage may be contraindicated.

Net contents: _____

EPA Reg. No. 49517-

EPA Est. No. 49517-TX-001
49517-CA-001

Approved with Conditions to EPA Letter Dated

10/22/93

Manufactured By:
Moore Agricultural Product Company, Inc.
Hanford CA Munday TX

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

49517-3

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes eye damage and skin irritation. Harmful if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin or on clothing.

Applications and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Sodium chlorate is a strong oxidizing agent. Do not mix with insecticides.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about person protective equipment (PPE) and restricted-entry notification. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standards and that involves contact with anything that has been treated such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks and protective eyewear.

Weather conditions and plant conditions affect defoliation. Weather conditions that favor good defoliation are high temperature, high humidity, low wind velocity, high to adequate soil moisture. Plant conditions which favor good defoliation are ample fertility and moisture, complete insect control. Nutrients should be ample but not too much nitrogen. LEAVES SHOULD BE GREEN AND TURGID (NOT WILTED, TOUGHENED OR STUNNED BY LACK OF FERTILIZERS).

MATURITY - Boll load heavy, most bolls mature, vegetative growth ceased, little or no second growth.

BEANS, DRY - To facilitate harvest and reduce bean seed moisture, apply 1 gallon of 5 lb. Sodium Chlorate in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 7-10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5-10 gallons of total solution per acre; for ground application, use a minimum of 20 gallons of total solution per acre. Do not graze treated fields or feed treated fodder or forage to livestock.

CHILI PEPPERS - Apply 1/2 to 1-1/4 gallons per acre in 5 to 15 gallons of water by aircraft and 20 to 40 gallons of water by ground rig. To be used on peppers for processing only. Apply 10 days before harvest. When grown for processing, consult a processor before applying.

GUAR BEANS - To facilitate harvest and reduce bean seed moisture, apply 1 gallon of 5 lb. Sodium Chlorate in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 7-10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5-10 gallons

of total solution per acre; for ground application use a minimum of 20 gallons of total solution per acre. Do not graze treated fields or feed treated fodder or forage to livestock.

COTTON - Defoliation - do not treat later than 7 to 10 days prior to harvest. Thorough coverage of the plant is essential.

AIR APPLICATION: Apply 2 or 3 quarts of 5 lb. Sodium Chlorate per acre in sufficient water to make 4 to 10 gallons total solution per acre. Use the lower rate when plants and weather conditions are ideal for defoliation. Swath width should not exceed wing spread; Altitude and speed of aircraft should be regulated to give sufficient spray coverage. Do not apply during periods of excessive wind as spray may drift and cause defoliation of adjoining cotton or injury to other crops. Do not apply during very high temperature found in some areas at mid-day.

GROUND MACHINE APPLICATION: Apply 2 to 3 quarts of 5 lb. Sodium Chlorate per acre in sufficient water to make 15 to 30 gallons total solution per acre. Cotton defoliant should be applied when 70% of the bolls are open. Poor defoliation often occurs under unfavorable weather, soil mixture, fertility and plant conditions. Uniform defoliation within a field may sometimes be difficult to obtain because of irregular plant conditions. Satisfactory defoliation within a field may not be obtained when cotton plants are suffering from lack of moisture or when plant growth is retarded by cold weather. Do not graze treated areas or feed gin waste to livestock. Flush out equipment with water after each day's work.

FLAX - For use as a harvest aid: to desiccate trashy weeds and desiccate the mature crop, apply 1 gallon of 5 lb. Sodium Chlorate per acre in enough water to give thorough coverage. Make application on bright, sunny days (preferable when temperature is above 60 degrees F.) 7-10 days before anticipated harvest date. For air application, use minimum 5 gallons ready-to-apply material per acre. Activity of the spray will be enhanced by the addition of a good nonionic surfactant at the rate of 1 quart per 100 gallons of spray.

GRAIN SORGHUM - As a desiccant, apply 3 quarts to 1 gallon of 5 lb. Sodium Chlorate per acre 7-10 days before harvest. Application should be made in enough water for thorough coverage. Apply a total of 4 to 10 gallons of spray per acre by air and 10 to 20 gallons per acre by ground equipment.

CORN - 5 lb. Sodium Chlorate may be used as a harvest aid for desiccation of trashy weeds in early maturing corn. Use 1 gallon of 5 lb. Sodium Chlorate in 5 to 7 gallons of water per acre by air. Apply at least 14 days before anticipated harvest date on clear days when temperature is expected to go above 70 degrees F. Use the 5 gallon rate when weeds are small and the crop canopy is fairly open. DO NOT graze treated fields or feed fodder, forage or residual seeds within 14 days of application.

RICE - As a pre-harvest desiccant, apply 3 quarts to 1 gallon of 5 lb. Sodium Chlorate in enough water for thorough coverage. By air, use a minimum of 5 gallons per acre. Application should be made 7 to 10 days before harvest. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SAFFLOWERS - As a pre-harvest desiccant, apply 1 gallon of 5 lb. Sodium Chlorate per acre in enough water for thorough coverage. By air, apply 5 to 10 total gallons of spray per acre. By ground, apply 20 to 30 gallons of total spray per acre. Application should be made 7 to 10 days before harvest when safflower heads are fully mature. Do not apply later than 7 days before harvest. Do not feed treated forage or graze treated fields. Not for use in California.

SOYBEANS - As a pre-harvest desiccant, apply 1 gallon of 5 lb. Sodium Chlorate per acre in enough water for thorough coverage. By air, apply a minimum of 5 gallons of spray solution per acre. Application should be made 7 to 10 days prior to normal anticipated harvest. Do not feed treated foliage or graze treated fields. Not for use in California.

7 of 11

SUNFLOWER - As a pre-harvest desiccant, apply 1 gallon of 5 lb. Sodium Chlorate per acre to fully mature seedheads in enough water for thorough coverage. By air, apply 5 to 10 gallons of total spray solution per acre. By ground equipment, apply 20 to 30 gallons of total spray solution per acre. Application should be made 7 days prior to normal harvest. Do not feed treated forage or graze treated fields. Not for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be in a cool, dry area away from any heat or ignition source.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of use according to label instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Moore Agricultural Products Co., Inc. (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning this product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damage and in no event shall damages or any other recovery of any kind against the Company exceed the price of the product which causes the alleged loss, damage, injury, or other claim. The Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income.

The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

11/2/83

M.A.P.C.O.

5 LB. SODIUM CHLORATE DEFOLIANT - DESSICANT

ACTIVE INGREDIENT:

Sodium Chlorate 45.0%
Inert ingredients 55.0%
TOTAL 100.0%

**KEEP OUT OF REACH
OF CHILDREN**

DANGER PELIGRO

Si usted no entiende la etiqueta,
busque a alguien para que se la
explique a usted en detalle. (If you
do not understand the label, find
someone to explain it to you in
detail.)

Net contents:

NOT REVIEWED

In Accordance with IR Notice 82-

EPA Reg. No. 49517-3
EPA Est. No. 49517-TX-001
49517-CA-001

JUN 1994
JUN 3 1994



Manufactured by:
Moore Agricultural Products Company, Inc.
Hanford CA Munday TX

STATEMENT OF PRACTICE

IF SPLASHED IN EYES, flush with water for 15 minutes.
IF SPILLED ON SKIN, immediately wash with plenty of water.
If irritation persists.
IF SWALLOWED, immediately drink a large quantity of water.
If these are not available, drink a large quantity of water.
NOTE TO PHYSICIAN: Probable mucosal damage. First aid measures contraindicated.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

DANGER

Corrosive. Causes eye damage and skin irritation. Harmful through the skin or inhaled. Do not get in eyes on or on clothing.
Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear
- Protective eyewear
- Chemical-resistant apron when loading



Follow manufacturer's instructions for use. If no such instructions for washing, wash PPE separately from other laundry.

1185

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Clean and wash spray equipment before and after use and do not allow washwater to contact desirable plants.

PHYSICAL AND CHEMICAL HAZARDS

Sodium chlorate is a strong oxidizing agent. Do not mix with insecticides. To avoid fire or explosion, do not mix with insecticides.

CHEMICAL PROHIBITION

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact other workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about Personal Protective Equipment (PPE) and restricted-entry notification. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for entry to treated areas that is permitted under the Worker Protection Standards and that involves contact with anything that has been treated such as plants, soil, or water is:



- 1) Coveralls over short-sleeved shirt and short pants
- 2) Waterproof gloves
- 3) Chemical-resistant footwear plus socks
- 4) Protective eyewear
- 5) Chemical-resistant headgear for overhead exposure

GENERAL INFORMATION

Weather conditions and plant conditions affect defoliation. Weather conditions that favor good defoliation are high temperature, high humidity, low wind velocity, high to adequate soil moisture. Plant conditions which favor good defoliation are ample fertility and moisture, complete insect control. Nutrients should be ample but not too much nitrogen. LEAVES SHOULD BE GREEN AND TURGID (NOT WILTED, TOUGHENED OR STUNNED BY LACK OF FERTILIZERS).

11801

MATURITY - Boll load heavy, most bolls mature, vegetative growth ceased, little or no second growth.

BEAN, DRY - To facilitate harvest and reduce bean seed moisture, apply 1 gallon of 5 lb. Sodium Chlorate in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 7-10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5-10 gallons of total solution per acre; for ground application, use a minimum of 20 gallons of total solution per acre. Do not graze treated fields or feed treated fodder or forage to livestock.

CHILI PEPPERS - Apply 1/2 to 3/4 gallons per acre in 5 to 15 gallons of water by aircraft and 20 to 40 gallons of water by ground rig. To be used on peppers for processing only. Apply 10 days before harvest. When grown for processing, consult a processor before applying.

GUAR BEANS - To facilitate harvest and reduce bean seed moisture, apply 1 gallon of 5 lb. Sodium Chlorate in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 7-10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5-10 gallons of total solution per acre; for ground application, use a minimum of 20 gallons of total solution per acre. Do not graze treated fields or feed treated fodder or forage to livestock.

COTTON - Defoliation - do not treat later than 7 to 10 days prior to harvest. Thorough coverage of the plant is essential.

AIR APPLICATION: Apply 2 or 3 quarts of 5 lb. Sodium Chlorate per acre in sufficient water to make 4 to 10 gallons total solution per acre. Use the lower rate when plants and weather conditions are ideal for defoliation. Swath width should not exceed wing spread; Altitude and speed of aircraft should be regulated to give sufficient spray coverage. Do not apply during periods of excessive winds as spray may drift and cause defoliation of adjoining cotton or injury to other crops. Do not apply during very high temperatures found in some areas at mid-day.

GROUND MACHINE APPLICATION: Apply 2 to 3 quarts of 5 lb. Sodium Chlorate per acre in sufficient water to make 15 to 30 gallons total solution per acre. Cotton defoliant should be applied when 70% of the bolls are open. Poor defoliation often occurs under unfavorable weather, soil moisture, fertility and plant conditions. Uniform defoliation within a field may sometimes be difficult to obtain because of irregular plant maturity. Satisfactory defoliation

within a field may not be obtained when cotton plants are suffering from lack of moisture or when plant growth is retarded by cold weather. Do not graze treated areas or feed gin waste to livestock. Flush out equipment with water after each day's work.

FLAX - For use as a harvest aid: to desiccate trashy weeds and desiccate the mature crop, apply 1 gallon of 5 lb. Sodium Chlorate per acre in enough water to give thorough coverage. Make application on bright, sunny days (preferably when temperature is above 60 degrees F.) 7-10 days before anticipated harvest date. For air application, use a minimum of 5 gallons ready-to-apply per acre. Activity of the spray will be enhanced by the addition of a good non-ionic surfactant at the rate of 1 quart per 100 gallons of spray.

GRAIN SOYBEAN - As a desiccant, apply 3 quarts to 1 gallon of 5 lb. Sodium Chlorate per acre 7-10 days before harvest. Application should be made in enough water for thorough coverage. Apply a total of 4 to 10 gallons of spray per acre by air and 10 to 20 gallons per acre by ground equipment.

CORN - 5 lb. Sodium Chlorate may be used as a harvest aid for desiccation of trashy weeds in early maturing corn. Use 1 gallon of 5 lb. Sodium Chlorate in 5 to 7 gallons of water per acre by air. Apply at least 14 days before anticipated harvest date on clear days when temperature is expected to go above 70 degrees F. Use the 5 gallon rate when weeds are small and the crop canopy is fairly open. DO NOT graze treated fields or feed fodder, forage or residual seeds within 14 days of application.

RICE - As a pre-harvest desiccant, apply 3 quarts to 1 gallon of 5 lb. Sodium Chlorate in enough water for thorough coverage. By air, use a minimum of 5 gallons per acre. Applications should be made 7 to 10 days before harvest. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SAFFLOWER - As a pre-harvest desiccant, apply 1 gallon 5 lb. Sodium Chlorate per acre in enough water for thorough coverage. By air, apply 5 to 10 total gallons of spray per acre. By ground, apply 20 to 30 gallons of total spray per acre. Application should be made 7 to 10 days before harvest when safflower heads are fully mature. Do not apply later than 7 days before harvest. Do not feed treated forage or graze treated fields. Not for use in California.

11
1/11

SOYBEANS - As a pre-harvest desiccant, apply 1 gallon of 5 lb. Sodium Chlorate per acre in enough water for thorough coverage. By air, apply a minimum of 5 gallons of spray solution per acre. For ground application, apply a minimum of 20 gallons of spray solution per acre. Application should be made 7 to 10 days prior to normal anticipated harvest period. Do not feed treated foliate or graze treated fields. Not for use in California.

SUNFLOWER - As a pre-harvest desiccant, apply 1 gallon of 5 lb. Sodium Chlorate per acre to fully mature seedheads in enough water for thorough coverage. By air, apply 5 to 10 gallons of total spray solution per acre. By ground equipment, apply 20 to 30 gallons of total spray solution per acre. Application should be made 7 days prior to normal harvest. Do not feed treated forage or graze treated fields. Not for use in California.



STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be in a cool, dry area away from any heat or ignition source.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture or

rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.



CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Moore Agricultural Products Co., Inc. (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning this product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damage and in no event shall damages or any other recovery of any kind against the Company exceed the price of the product which causes the alleged loss, damage, injury, or other claim. The Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income.

The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.