

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes eye damage and skin irritation. Harmful or fatal if swallowed, absorbed through the skin or inhaled. Do not get in eyes, on skin or on clothing.

Applicators and other handlers must wear:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant apron when cleaning equipment, mixing, or loading.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Clean and wash spray equipment before and after use and do not allow washwater to contact desirable plants. Do not contaminate water by disposal of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Sodium chlorate is a strong oxidizing agent. To avoid fire or explosion do not mix with insecticides.

IRRIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

M.A.P.C

MOORE AG BRAND P DEFOLIANT-DES

Active Ingredient:

Sodium Chlorate

Inert Ingredients

Total

KEEP OUT OF REACH OF
DANGER

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICES

IF IN EYES, Immediately flush with water for at least 15 minutes. Get medical attention.

IF ON SKIN, Immediately wash with soap and water for at least 15 minutes. Get medical attention if irritation persists.

IF SWALLOWED, Immediately drink milk, egg whites or gelatin solution. Do not induce vomiting. Get medical attention.

IF SWALLOWED, Immediately drink milk, egg whites or gelatin solution. Do not induce vomiting. Get medical attention.

IF SWALLOWED, Immediately drink milk, egg whites or gelatin solution. Do not induce vomiting. Get medical attention.

NOTE TO PHYSICIAN

Probable mucosal damage. Use contraindicated.

Net contents:

EPA Reg. No. 49517-3 EPA Es

Manufactured By
Moore Agriculture
Hanford CA

194

P.CO.

**ND POLY-FOLIANT V
T-DESICCANT**

..... 45.0%
..... 55.0%
..... 100.0%

**EACH OF CHILDREN
PELIGRO**

a, busque a alguien para que se la
you do not understand the label, find
detail.)

PRACTICAL TREATMENT

sh with water for 15 minutes. Get

sh with plenty of soap and water.
on persists.

ly drink a large quantity of either
olution. If these are not available,
er. Avoid alcohol. Get medical

PHYSICIAN:

Use of gastric lavage may be

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE INSTRUCTIONS: Storage should be in a cool, dry area away from any heat or ignition source.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.


CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

**CONDITIONS OF SALE - LIMITED WARRANTY
AND LIMITATIONS OF LIABILITY AND REMEDIES**

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions, the failure to follow the label directions, or good application practices, all of which are beyond the control of Moore Agricultural Products Co., Inc. (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. The Company makes no other warranties or representations of any kind, express or implied, concerning this product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damage and in no event shall damages or any other recovery of any kind against the Company exceed the price of the product which causes the alleged loss, damage, injury, or other claim. The Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental or consequential damages or expense of any nature, including, but not limited to, loss of profits or income.

The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.



ments: _____

PA Est. No. 49517-TX-001
49517-CA-001

ed By:
ultural Products Company, Inc.
Munday TX

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For air, requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about Personal Protective Equipment (PPE) and restricted-entry notification. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standards and that involves contact with anything that has been treated such as plants, soil, or water is: 1) coveralls over long-sleeved shirt and long pants, 2) waterproof gloves, 3) chemical-resistant footwear plus socks, 4) protective eyewear and 5) chemical-resistant headgear for overhead exposure.

Weather conditions and plant conditions affect defoliation. Weather conditions that favor good defoliation are high temperature, high humidity, low wind velocity, high to adequate soil moisture. Plant conditions which favor good defoliation are ample fertility and moisture, complete insect control. Nutrients should be ample but not too much nitrogen. LEAVES SHOULD BE GREEN AND TURGID (NOT WILTED, TOUGHENED OR STUNNED BY LACK OF FERTILIZERS).

MATURITY - Boll load heavy, most bolls mature, vegetative growth ceased, little or no second growth.

GENERAL INFO

BEANS, DRY - To facilitate harvest and reduce defoliation of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant thorough coverage either by aircraft or ground application. Do not apply during the middle of the day. For air application, use a minimum of 10 gallons per acre; for ground application, use a minimum of 20 to 40 gallons of water by ground processing only. Apply 10 days before harvest or before processing.

CHILI PEPPERS - Apply 1/2 to 1-1/4 gallons of Poly-Foliant V Defoliant-Desiccant per acre by aircraft and 20 to 40 gallons of water by ground processing only. Apply 10 days before harvest or before processing. Do not apply during the middle of the day. For air application, use a minimum of 10 gallons per acre; for ground application use a minimum of 20 to 40 gallons of water by ground processing only. Apply 10 days before harvest or before processing. Do not graze treated fields or feed treated forage to livestock. Thorough coverage of the plant is essential.

AIR APPLICATION: Apply 2 or 3 gallons of Poly-Foliant V Defoliant-Desiccant per acre in sufficient water to make a 2% solution per acre. Use the lower rate when applied for defoliation. Swath width should not exceed 100 feet. Aircraft should be regulated to give sufficient coverage during periods of excessive wind as spray may drift and cause injury to other crops. Do not apply during areas at mid-day.

GROUND MACHINE APPLICATION: Apply 2 or 3 gallons of Poly-Foliant V Defoliant-Desiccant per acre in sufficient water to make a 2% solution per acre. Cotton defoliation is best when the bolls are open. Poor defoliation often occurs with low moisture, fertility and plant conditions. Uniform defoliation within a field may not be obtained because of uneven moisture or when plant growth is retarded in treated areas or feed gin waste to livestock. Do not graze treated areas or feed gin waste to livestock. Do not apply during areas at mid-day.

AL INFORMATION

and reduce bean seed moisture, apply 1 gallon Defoliant-Desiccant in enough water to provide it or ground equipment. Make application 7-10 days before harvest. Do not apply under conditions of extreme heat. For application, use 5-10 gallons of total solution per acre. Do not use on treated fodder or forage to livestock.

Apply 4 gallons per acre in 5 to 15 gallons of water by air or ground equipment. To be used on peppers for processing. When grown for processing, consult manufacturer.

and reduce bean seed moisture, apply 1 gallon Defoliant-Desiccant in enough water to provide it or ground equipment. Make application 7-10 days before harvest. Do not apply under conditions of extreme heat. For application, use 5-10 gallons of total solution per acre. Do not use on treated fodder or forage to livestock. Do not treat later than 7 to 10 days prior to harvest. Essential.

Apply 2 or 3 quarts of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant in sufficient water to make 4 to 10 gallons total per acre when plants and weather conditions are ideal. Do not exceed wing spread; Altitude and speed of application should provide sufficient spray coverage. Do not apply during high wind may drift and cause defoliation of adjoining cotton fields. Only during very high temperature found in some areas.

APPLICATION: Apply 2 to 3 quarts of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant per acre in sufficient water to make 4 to 10 gallons total per acre. Cotton defoliants should be applied when 70% of cotton has opened. Cotton defoliation often occurs under unfavorable weather, soil conditions. Uniform defoliation within a field may be caused by irregular plant conditions. Satisfactory defoliation is obtained when cotton plants are suffering from stress which is retarded by cold weather. Do not graze treated fields or feed treated forage to livestock. Flush out equipment with water after application.

FLAX - For use as a harvest aid: to desiccate trashy weeds and desiccate the mature crop, apply 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant per acre in enough water to give thorough coverage. Make application on bright, sunny days (preferable when temperature is above 60 degrees F.) 7-10 days before anticipated harvest date. For air application, use minimum 5 gallons ready-to-apply material per acre. Activity of the spray will be enhanced by the addition of a good nonionic surfactant at the rate of 1 quart per 100 gallons of spray.

GRAIN SORGHUM - As a desiccant, apply 3 quarts to 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant per acre 7-10 days before harvest. Application should be made in enough water for thorough coverage. Apply a total of 4 to 10 gallons of spray per acre by air and 10 to 20 gallons per acre by ground equipment.

CORN - Moore Ag Brand Poly-Foliant V Defoliant-Desiccant may be used as a harvest aid for desiccation of trashy weeds in early maturing corn. Use 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant in 5 to 7 gallons of water per acre by air. Apply at least 14 days before anticipated harvest date on clear days when temperature is expected to go above 70 degrees F. Use the 5 gallon rate when weeds are small and the crop canopy is fairly open. DO NOT graze treated fields or feed fodder, forage or residual seeds within 14 days of application.

RICE - As a pre-harvest desiccant, apply 3 quarts to 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant in enough water for thorough coverage. By air, use a minimum of 5 gallons per acre. Application should be made 7 to 10 days before harvest. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SAFFLOWERS - As a pre-harvest desiccant, apply 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant per acre in enough water for thorough coverage. By air, apply 5 to 10 total gallons of spray per acre. By ground, apply 20 to 30 gallons of total spray per acre. Application should be made 7 to 10 days before harvest when safflower heads are fully mature. Do not apply later than 7 days before harvest. Do not feed treated forage or graze treated fields.

SOYBEANS - As a pre-harvest desiccant, apply 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant per acre in enough water for thorough coverage. By air, apply a minimum of 5 gallons of spray solution per acre. For ground application, apply a minimum of 20 gallons of spray solution per acre. Application should be made 7 to 10 days prior to normal anticipated harvest. Do not feed treated foliage or graze treated fields.

SUNFLOWER - As a pre-harvest desiccant, apply 1 gallon of Moore Ag Brand Poly-Foliant V Defoliant-Desiccant per acre to fully mature seedheads in enough water for thorough coverage. By air, apply 5 to 10 gallons of total spray solution per acre. By ground equipment, apply 20 to 30 gallons of total spray solution per acre. Application should be made 7 days prior to normal harvest. Do not feed treated forage or graze treated fields.

SOUTHERN PEAS - Moore Ag Brand Poly-Foliant V Defoliant-Desiccant at a rate of 1 gallon per acre in 5 to 10 gallons of water by air or 10 to 20 gallons by ground equipment. Thorough coverage is essential. Make application 7 to 10 days before anticipated harvest, longer if temperatures are cool. DO NOT graze treated fields or feed treated fodder or forage to livestock.