

Sevin®

brand carbaryl insecticide

Garden Dust

[Kills: Armyworms, tomato Hornworms, Leafhoppers, and many other listed pests.]

Optional Art Work of Target Pests:

[Caterpillar]	[Armyworm]
[Leafhopper]	[Beetle]

Active Ingredients:
 *Carbaryl: 1-naphthyl N-methylcarbamate 5.00%
 Inert Ingredients: 95.00%
 [TOTAL: 100.00%]
 (*Sevin® is a registered trademark of Rhône-Poulenc for carbaryl insecticides.)

Keep Out Of Reach Of Children
CAUTION See back panel for additional precautionary statements.

NET WT. [MASS] [1] LB. [(453 g)]

[Read entire label.] [Stop. Read entire label before use.]
 [IMPORTANT: To avoid damage or poor results, carefully read and follow all directions and precautionary statements.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use around the home only.

OPTION A - [For Bag]

(PARAGRAPH FORMAT)

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can or other appropriate garden duster. Treat when insects or their damage first appear. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust. Lightly cover both upper and lower leaf surfaces with a thin, even film of dust. Foliage should be dry before dust is applied. Allow dust to settle in treated areas before returning. For best results, treat every 7 - 10 days. Do not apply more than once a week. For plants above chest height, consider using a liquid sevin (carbaryl) product.

(NUMBERED FORMAT)

HOW TO USE

1. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can or other appropriate garden duster.
2. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust.
3. Lightly cover both upper and lower leaf surfaces with a thin, even film of dust. Foliage should be dry before dust is applied.
4. Allow dust to settle in treated areas before returning.

USE TIPS

1. Treat when insects or their damage first appear.
2. For best results, treat every 7 - 10 days. Do not apply more than once a week.
3. For plants above chest height, consider using a liquid sevin (carbaryl) product.

OPTION B - [For Shaker Canister]

(PARAGRAPH FORMAT)

Shake canister to loosen any settled dust. Treat when insects or their damage first appear. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust. Wearing household latex or

rubber gloves, open top, invert canister, and lightly cover both upper and lower leaf surfaces with a thin even film of dust. Foliage should be dry before dust is applied. Allow dust to settle in treated areas before returning. For best results, treat every 7 - 10 days. Do not apply more than once a week. For plants above chest height, consider using a liquid sevin (carbaryl) product.

(NUMBERED FORMAT)

HOW TO USE

1. Shake canister to loosen any settled dust.
2. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust.
3. Wearing household latex or rubber gloves, open top, invert canister, and lightly cover both upper and lower leaf surfaces with a thin even film of dust. Foliage should be dry before dust is applied.
4. Allow dust to settle in treated areas before returning.

USE TIPS

1. Treat when insects or their damage first appear.
2. For best results, treat every 7 - 10 days. Do not apply more than once a week.
3. For plants above chest height, consider using a liquid sevin (carbaryl) product.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of the gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow people or pets to contact treated areas until dust has settled.

[FLEAS & TICKS]

To control fleas and ticks in home lawns, apply evenly at the rate of one [1] pound per 250 sq. ft.

ACCEPTED

MAR 5 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 49585-4

[Bracketed Text] = Optional Language

BEST AVAILABLE COPY

FN: A4-REG

PN 19 49585-4

1/

[FRUITS & VEGETABLES]

Number of Days to Wait Between Last Application and Harvest			
FRUIT & VEGETABLE	PHI*	FRUIT & VEGETABLE	PHI*
Beans	0	Lettuce	14
Broccoli	3	Melons	0
Brussels sprouts	3	Mustard greens	14
Cabbage	3	Peppers	0
Carrots	0	Potatoes	0
Cauliflower	3	Pumpkins	0
Collards	14	Spinach	14
Corn	0	Squash	0
Cucumbers	0	Strawberries	1
Eggplants	0	Swiss chard	14
Grapes	7	Tomatoes	0

*PHI = Pre-Harvest Interval.

FRUITS OR VEGETABLES	[GARDEN PESTS] INSECTS
Tomatoes	Armyworm (Fall), Beetles (Flea), European corn borer, Leafhopper, Tomato fruitworm, Tomato hornworm
Beans	Armyworm, Beetles (Beanleaf, Cucumber, Flea, Japanese, Mexican bean), Cutworm (Western bean), Corn earworm, Velvetbean caterpillar, Stink bug
Cabbage, Broccoli, Brussels Sprouts, Cauliflower	Armyworm, Beetles (Flea), Cabbage caterpillar, Corn earworm, Harlequin bug

FRUITS OR VEGETABLES	[GARDEN PESTS] INSECTS
Corn	Armyworm (Fall), Beetles (Flea, Japanese, Sap), European corn borer, Leafhopper
Cucumbers, Melons, Pumpkin, Squash	Beetles (Cucumber, Flea), Leafhopper, Melonworm, Pickleworm, Squash bug Do not use on watermelons in Florida. Avoid excessive applications
Lettuce, Carrots	Armyworm, Beetles (Flea), Cabbage caterpillar, Corn earworm, Harlequin bug, Leafhopper, Lygus bug, Spittlebug, Stink bug, Tarnished plant bug
Peppers, Potato, Eggplant	Armyworm (Fall), Beetles (Colorado potato, Flea), European corn borer, Lace bug, Leafhopper, Lygus bug, Stink bug, Tarnished plant bug, Tomato fruitworm, Tomato hornworm
Strawberries	Beetles (Flea), Leafhopper, Lygus bug, Spittlebug, Strawberry leafroller
Collards, Mustard Greens, Spinach, Swiss Chard	Armyworm, Beetles (Flea), Corn earworm, Leafhopper, Stink bug
Grapes	Beetles (Japanese, June), Grape berry moth, Grape leaf (Folder, Hopper, Skeletonizer)

STORAGE AND DISPOSAL

Storage: Keep in original [container] [bag] [canister]. Store in a cool, dry place, preferably in a locked storage area inaccessible to children.
Disposal: Do not reuse empty [container] [bag] [canister]. Securely wrap [container] [bag] [canister] in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Keep people and pets out of treated areas.

Statement of Practical Treatment

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention immediately. If on Skin: Wash thoroughly with soap and water. Get medical attention if irritation develops. If in Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists. Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Environmental Hazards

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call [800-645-4957] [800-332-5553]

[Made in the USA for KMART Corporation, Troy, Michigan 48084]

Distributed by Alljack, Division of United Industries Corporation, P. O. Box 15482, St. Louis, MO 63114-0842

EPA Reg. No.: 49585-4

EPA Est. Nos.: 49585-MI-1 A, 769-GA-1 C, 42056-IL-1 E, 44616-MO-1 H, 42761-MS-1 N, 9591-OH-1 O

Superscript used is first letter of lot number.

© 1996

272