

PM-19

Reg # 49585-24

193

FRONT PANEL

Sevin® Plus Multi-Purpose Garden Dust

Controls Powdery Mildew, Tomato Hornworm, Leafhoppers and many other listed pests

ACTIVE INGREDIENTS:

*Carbaryl: 1-naphthyl N-methylcarbamate	5.00%
**Piperonyl butoxide, Technical	0.45%
Pyrethrins	0.03%
Sulphur	30.00%
INERT INGREDIENTS:	64.52%

**Equivalent to 0.36% of (butylcarbityl)(6-propylpiperonyl)ether and to 0.09% of related compounds.

*Sevin® is a registered trademark of Rhône-Poulenc for carbaryl insecticides.

Keep Out Of Reach Of Children

CAUTION

See back panel for additional precautionary statements.

NET WT. 10 OZ. (283 g)

ACCEPTED
OCT 18 1996
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 49585-24

96 JUL 19 P4:47

REC'D EPA/REG/P/DPD02

BACK PANEL

IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ AND FOLLOW ALL DIRECTIONS AND PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. Wash thoroughly after handling and before eating or smoking.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison Control Center immediately. Give victim 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If in Eyes: Flush eyes with plenty of water. If irritation persists get medical attention.

If on Skin: Remove contaminated clothing and wash skin with plenty of soap and water.

Note to Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water when disposing of rinsate or equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STOP. Read Entire Label Before Use.

BACK PANEL (CONTINUED)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. FOR USE AROUND THE HOME ONLY.

APPLICATION: Before using, shake bottle and tap bottom lightly to loosen any settled product.

Hold this container so that the discharge nozzle is pointed away from the face and body, then squeeze. Avoid standing in the path of dust. For successful control of insects and disease, apply dust to all exposed plant parts, including upper and lower leaf surfaces, covering thoroughly. For best results, apply dust in early morning or late evening when the temperature is low and the air is calm. Avoid excessive application. Apply when insects or their damage first appear and repeat treatment every 7-10 days unless shorter interval is specified below. To protect against diseases such as Rust, Black Spot and Powdery Mildew on plants, apply dust as foliage appears and before disease is apparent. Repeat application at weekly intervals. Subsequent planting of food or feed crops in treated soil must be made with crops listed on this or other carbaryl labels.

CROPS	Days between last application and harvest	GARDEN PEST
Tomatoes	0	Tomato Hornworm, Tomato Fruitworm, Tarnished Plant Bug, Stink Bug, Lacebug, Flea Beetle, Fall Armyworm, Aphids.
Corn	0	Corn Earworm, Fall Armyworm, Leafhopper, Flea Beetle.
For Corn Earworm control, apply at 3 day intervals starting when silks appear and continuing until silks begin to dry. Three or more applications may be necessary for severe infestations.		
Beans	0	Black Bean Aphid, Mexican Bean Beetle, Red Spider Mite, Flea Beetle, Tarnished Plant Bug, Leafhopper, Climbing Cutworm.
Lettuce, Leaf	14	Leaf-eating Caterpillars, Climbing Cutworms,
Head	3	Leafhoppers, Flea Beetle.
Peas	3	Powdery Mildew, Aphids, Leafhopper, Colorado Potato Beetle, Climbing Cutworm, Grasshopper.
Cabbage	3	Cabbage Caterpillar, Armyworm, Flea Beetle, Corn Earworm, Harlequin Bug.
Radishes, Beet	3	Flea Beetles, Armyworm, Harlequin Bug.
Roots, Turnip Roots		
Beet Tops, Turnip Tops	14	Flea Beetles, Armyworm, Harlequin Bug.
Strawberries	1	Strawberry Weevil, Spittlebug, Leafroller, Aphids
Blackberries, Loganberries, Boysenberries, Dewberries	7	Red Spider Mite, Cankerworm, Blackberry Mite, Exposed Thrips, Orange Tortrix, Powdery Mildew
Grapes	7	Powdery Mildew, Leafhopper, Grape Leaf Skeletonizer, Climbing Cutworm, Grapeberry Moth, June Beetle, Grape Leaf Folder.
For Grape Leaf Folder control, apply just before first brood of Leaf folder larvae emerge from rolls.		
Carrots	0	Flea Beetle, Armyworm, Harlequin Bug, Six-spotted Leafhopper.
To control Six-spotted Leafhopper, apply in a 5 to 7 day schedule.		

393

BACK PANEL (CONTINUED)

Roses and other ornamental Flowers, Shrubs and Shade Trees: Rose Aphid, Red Spider Mite, Rose Slug, Scale Crawlers, Tent Caterpillars, Plant bugs, Leafhoppers, Elm Leaf Beetle, Leaf-eating Caterpillars, Climbing cutworms, Thrips (exposed), Flea Beetles, Oak and Birch Leafminers, Orange Tortrix, Mealybugs, Grasshoppers, Powdery Mildew, Rust and Black Spot on Roses.

NOTE: Do not apply on Boston Ivy, Virginia Creeper and Maidenhair Fern as injury will result.

Outdoors: Earwigs, fleas, ticks, grasshoppers, ants, sowbugs. Dust freely on soil surfaces, turf and foliage of infested areas, along fences, walls, curbs and foundations, in places that provide shelter and shade, in cracks and crevices. Treat when air is still, to avoid drift. Apply at rate of 1 pound to each 250 sq.ft. of surface area.

STORAGE AND DISPOSAL

Store in a cool dry area away from heat or open flame. Do not reuse container. Discard container in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Made in the USA for KMart Corporation, Troy, Michigan 48084

Questions or comments? Call 800-645-4957.

Alljack, Division of United Industries Corp., P.O. Box 15842, St. Louis, MO 63114-0842

EPA Reg. No.: 49585-24

EPA Est. No.: 49585-MI-1 A, 769-GA-1 C, 42056-IL-1 E, 44616-MO-1 H, 42761-MS-1 N, 9591-OH-1 O

Superscript used is first letter of lot number.