

AUG 09 1995
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 49585-26

Pm 19 49585-26 page 1 of 3

FRONT PANEL

K GRO® SEVIN® brand carbaryl insecticide 10 DUST Formula II

[To Control Leafhoppers, Japanese Beetles, Bagworms, Tent Caterpillars and other insects listed on this label.]

ACTIVE INGREDIENT:

*Carbaryl (1-Naphthyl N-methylcarbamate)	10.0%
INERT INGREDIENTS:	90.0%
[TOTAL:	100.0%]

[*SEVIN® is the registered trademark of Rhône-Poulenc for carbaryl insecticides.]

Keep Out Of Reach Of Children

CAUTION

**See back panel for additional precautionary statements.
 Read Entire Label.**

NET WT. 4 lbs. (1.81 kg)

BACK PANEL

IMPORTANT: TO AVOID DAMAGE OR POOR RESULTS, CAREFULLY READ ALL PRECAUTIONARY STATEMENTS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled: Remove victim to fresh air.

If in eyes: Flush eyes with plenty of water.

If on skin: Wash with plenty of soap and water.

Note to physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine by injection is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **FOR HOUSEHOLD USE ONLY.**

APPLICATION: Apply uniformly to foliage. Cover all exposed parts (both upper and lower leaf surfaces) with a thin, even film of dust using any garden type duster. Apply preferably in early

1 of 3

morning or late evening when the air is quiet. Apply when insects or their damage appear. Best results are secured by dusting regularly at 7-10 day intervals throughout the growing season. Foliage should be dry before dust is applied. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Crops	Insects Controlled	Last Application Allowed Before Harvest
Asparagus (On Spears)	Asparagus Beetle, Grasshoppers.	1 day
Asparagus (On Ferns & Brush)	Do not repeat application within 3 days. Asparagus Beetle, Apache Cicada	Post-Harvest Application
Beans	Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Flea Beetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms.	0 days
Cucumbers, Squash, Cantaloupes, Watermelons	Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm, Pickleworm, Squash Bug, Leafhoppers, Flea Beetles.	0 days
Potatoes, Tomatoes, Eggplants, Peppers	Flea Beetles, Armyworms (Fall), Tomato Fruit Worm, Corn Earworm, Tomato Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Lace Bug, Lygus Bugs.	0 days
Lettuce	Flea Beetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebugs, Lygus Bugs.	3 Days (head) 14 Days (Leaf)
Carrots	Flea Beetles, Leafhoppers, Six-spotted Leafhopper, (Aster Yellows Vector), Armyworms, Spittlebugs, Lygus & Stink Bugs.	0 days
Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi	Flea Beetles, Harlequin Bugs, Corn Earworms, Armyworms.	3 days
Okra	Corn Earworm, Stink Bugs	0 days
Strawberries	Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil.	1 day
Grapes	Grape Leafhopper, Grape Leaf Folder, Grape Berry Moth, European Fruit Lacanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-Banded Leaf Roller.	7 days

ORNAMENTALS: To control Apple Aphid, Bagworm, Birch Leafminer, Blister Beetles, Boxelder Bug, Boxwood Leafminer, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetles, Japanese Beetle, Lacebugs, Leafhopper, Leaf Rollers, Oak Leafminer, Orange Tortrix, periodical Cicada, Plant Bugs, Psyllids, Rose Slug, Rose Aphid, scale insects in the crawler stage, Tent Caterpillars, Willow Leaf Beetle and exposed Thrips, apply thoroughly to the infested plants at the first sign of damage and repeat at 7 day intervals or as necessary for good insect control. To obtain scale control, apply in spring and early summer when scale crawlers are present.

LAWNS: To control Leafhoppers, Fall Armyworm, and Sod Webworm, apply at the rate of 2 pounds per 1000 square feet. For best results, apply after rain or watering. Do not water for at least 2 days after application. Repeat in 2 or 3 weeks if necessary.

EARWIGS: This material is effective in controlling earwigs and when properly used will reduce the number gaining entrance to homes and buildings. Apply heavily in a band 3 to 4 inches wide using a shaker-type can. Apply at the rate of 1 lb. for 150 linear feet. Earwigs breed over wide areas around yards and are driven by cold weather to seek warm shelter. As they move along the foundation wall seeking entrance, they come in contact with the treated barrier and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation wall. Treat also around refuse piles, lumber, mulch and other areas in the yard offering areas of concealment to earwigs. Repeat application in 10 days or when earwigs become troublesome. If the dust barrier is washed away by rain, it should be replaced with a new application if earwigs are still a problem.

PEST INSECT CONTROL

DOGS: For ticks and fleas, dust over the entire animal, avoiding eyes, and rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of animals less than four weeks of age. Do not use on pregnant dogs.

DOG QUARTERS: For ticks and fleas, dust in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary.

STORAGE AND DISPOSAL

Storage: Store product in original container preferably in a locked storage area inaccessible to children. Do not store near fertilizer, seed or food products.

Disposal: Do not reuse empty container. Securely wrap container in several layers of newspaper and put in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

[Made in the USA for K Mart Corporation, Troy, Michigan 48064]

Registered and Manufactured by Alljack & Company, [377 Amelia,] Plymouth, Michigan 48170

EPA Reg. No. 49585-26

EPA Est. No. 49585-MI-1