

PM 19 49585-26

Page 1 of 3

KGRO Sevin brand carbaryl insecticide 10 Dust Formula II

EPA Reg. No. 49585-26

page 1 of 3

KGRO

SEVIN®

brand carbaryl insecticide

10 DUST Formula II

[To Control Leafhoppers, Japanese Beetles, Bagworms, Tent Caterpillars and other insects listed on this label.]

ACCEPTED
MAR 27 1997
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 49585-26

Active ingredient: Carbaryl (1-Naphthyl N-methylcarbamate) 10.0%
Inert ingredients: 90.0%
[TOTAL: 100.0%]
[*SEVIN® is the registered trademark of Rhône-Poulenc for carbaryl insecticides.]

Keep Out Of Reach Of Children

CAUTION See back panel for additional precautionary statements.

NET WT. 4 lbs. (1.81 kg)

[Read entire label.] [Stop. Read entire label before use.]
[IMPORTANT: To avoid damage or poor results, carefully read and follow all directions and precautionary statements.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use around the home only.

OPTION A - [For Bag]

(PARAGRAPH FORMAT)

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can or other appropriate garden duster. Treat when insects or their damage first appear. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust. Lightly cover both upper and lower leaf surfaces with a thin, even film of dust. Foliage should be dry before dust is applied. Allow dust to settle in treated areas before returning. For best results, treat every 7 - 10 days. Do not apply more than once a week. For plants above chest height, consider using a liquid sevin (carbaryl) product.

(NUMBERED FORMAT)

HOW TO USE

1. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can or other appropriate garden duster.
2. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust.
3. Lightly cover both upper and lower leaf surfaces with a thin, even film of dust. Foliage should be dry before dust is applied.
4. Allow dust to settle in treated areas before returning.

USE TIPS

1. Treat when insects or their damage first appear.
2. For best results, treat every 7 - 10 days. Do not apply more than once a week.
3. For plants above chest height, consider using a liquid sevin (carbaryl) product.

OPTION B - [For Shaker Canister]

(PARAGRAPH FORMAT)

Shake canister to loosen any settled dust. Treat when insects or their damage first appear. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust. Wearing household latex or rubber gloves, open top, invert canister, and lightly cover both upper and lower leaf surfaces with a thin even film of dust. Foliage should be dry before dust is

applied. Allow dust to settle in treated areas before returning. For best results, treat every 7 - 10 days. Do not apply more than once a week. For plants above chest height, consider using a liquid sevin (carbaryl) product.

(NUMBERED FORMAT)

HOW TO USE

1. Shake canister to loosen any settled dust.
2. Apply when air is calm to avoid drift and contact with eyes and skin. Start at the farthest corner of area to be treated and work backward to avoid contact with dust.
3. Wearing household latex or rubber gloves, open top, invert canister, and lightly cover both upper and lower leaf surfaces with a thin even film of dust. Foliage should be dry before dust is applied.
4. Allow dust to settle in treated areas before returning.

USE TIPS

1. Treat when insects or their damage first appear.
2. For best results, treat every 7 - 10 days. Do not apply more than once a week.
3. For plants above chest height, consider using a liquid sevin (carbaryl) product.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use.
- Wash the outside of the gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow people or pets to contact treated areas until dust has settled.

LAWNS

INSECT	HOW TO USE
Armyworm (Fall), Leafhoppers, Sod webworm,	Apply at the rate of 2 pounds per ,1000 sq. ft. For best results, apply after rain or watering. Do not water for at least 2 days after application. Repeat in 2 or 3 weeks if necessary.

FRUITS & VEGETABLES

Number of Days to Wait Between Last Application and Harvest			
FRUIT & VEGETABLE	PHI*	FRUIT & VEGETABLE	PHI*
Asparagus (Brush)	**	Eggplants	0
Asparagus (Ferns)	**	Grapes	7
Asparagus (Spears)	1	Kohlrabi	3
Beans	0	Lettuce	14
Broccoli	3	Okra	0
Brussels Sprouts	3	Peppers	0
Cabbage	3	Potatoes	0
Cantaloupes	0	Squash	0
Carrots	0	Strawberries	1
Cauliflower	3	Tomatoes	0
Cucumbers	0	Watermelons	0

*PHI = Pre-Harvest Interval. **Post-Harvest Application.

FRUITS OR VEGETABLES	[GARDEN PESTS] INSECTS
Asparagus (Brush, Ferns, Spears)	Apache cicada, Asparagus beetle, Grasshoppers
Beans	Amyworms, Bean leaf roller, Beetles (Bean leaf, Flea, Japanese, Mexican bean), Corn earworm, Cutworm (Western Bean), Leafhoppers, Lygus bugs, Stink bugs, Tarnished plant bug, Velvetbean caterpillar
Cucumbers, Squash, Cantaloupes, Watermelons	Beetles (Flea, Spotted cucumber, Striped cucumber), Leafhoppers, Melonworm, Pickleworm, Squash bug

FRUITS OR VEGETABLES	[GARDEN PESTS] INSECTS
Potatoes, Tomatoes, Eggplants, Peppers	Amyworms (Fall), Beetles (Flea, Colorado potato), Corn earworm, European corn borer, Lace bug, Leafhoppers, Lygus bugs, Tomato fruit worm, Tomato hornworms, Stink bugs
Lettuce	Amyworms, Beetles (Flea), Corn earworm, Harlequin bug, Leafhoppers, Lygus bugs, Spittlebugs, Stink bugs
Carrots	Amyworms, Beetles (Flea), Leafhoppers (Six-spotted [Aster Yellow Vector]), Lygus bugs, Spittlebugs, Stink bugs
Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi	Amyworms, Beetles (Flea), Corn earworms, Harlequin bug
Okra	Corn earworm, Stink bugs
Strawberries	Spittlebugs, Strawberry leaf roller, Strawberry weevil
Grapes	Beetles (Japanese, June), European fruit lecanium, Grape berry moth, Grape leaf (Folder, Hopper, Skeletonizer), Red-banded leaf roller

ORNAMENTALS

INSECTS	HOW TO USE
Aphid (Apple, Elm leaf, Rose), Bagworm, Beetle (Blister, Elm leaf, Flea, Japanese), Boxelder bug, Lacebugs, Leafhopper, Leafminers (Birch, Boxwood, Oak), Leaf rollers, Orange loirix, periodical Cicada, Plant bugs, Psyllids, Rose slug, Scale insects (In the crawler stage, Tent caterpillars, Willow leaf beetle), Thrips (Exposed)	Apply thoroughly to the infested plants at the first sign of damage and treat every 7 day or as necessary for good insect control. To obtain scale control, apply in spring and early summer when scale crawlers first appear.

EARWIGS

THINGS TO KNOW	HOW TO USE
Use as a barrier treatment around buildings. Earwigs breed over wide areas around yards and are driven by cold weather to seek warm shelter. As they move along the foundation wall seeking entrance, they come in contact with the treated area and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation. Also treat around refuse piles, lumber, mulch and other areas in the yard offering areas of concealment to earwigs.	Using a shaker-type can, around the foundation apply a 3 to 4 inches wide band at the rate of 1 lb. for 150 linear feet. Repeat application in 10 days. If the dust is washed away by rain, it should be replaced with a new application if earwigs are still a problem. Do not repeat application more than once a week.

STORAGE AND DISPOSAL

Storage: Store product in original [container] [bag] [canister] preferably in a locked storage area inaccessible to children. Do not store near fertilizer, seed or food products. **Disposal:** Do not reuse empty [container] [bag] [canister]. Securely wrap [container] [bag] [canister] in several layers of newspaper and put in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

Statement of Practical Treatment

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **If Inhaled:** Remove victim to fresh air. **If In Eyes:** Flush eyes with plenty of water. **If on Skin:** Wash with plenty of soap and water. Note to Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine by injection is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

Environmental Hazards

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct

3 7 3

treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or comments? Call [800-845-4957] [800-332-5553]

[Made in the USA for KMART Corporation, Troy, Michigan 48064]

Distributed by Alljack, Division of United Industries Corporation,
P. O. Box 15482, St. Louis, MO 63114-0842

EPA Reg. No.: 49585-26

EPA Est. Nos.: 49585-MI-1 A , 789-GA-1 C , 42056-IL-1 E ,
44616-MO-1 H , 42761-MS-1 N , 9591-OH-1 O

Superscript used is first letter of lot number.

© 1996