

49668-1 Pinal 10/72

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**CAUTION**


Hazard if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff conditions are likely to occur. Do not apply where weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL AND CHEMICAL HAZARDS**

Sulfur dust suspended in air irritates eyes. Keep away from heat, sparks and flame. Do not smoke while applying this product.



**SUPERFINE**  
**DUSTING SULPHUR**

**INSTRUCTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**RE-ENTRY STATEMENT**

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Notice of oral warnings must be given to workers who are expected to be on a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**CAUTION:** Area treated with **SUPERFINE DUSTING SULFUR** (See label for details). Do not enter without appropriate protective clothing for 24 hours following application. In case of accidental exposure, see the Statement for Practical Treatment on this label.

For Spraying  
 Vegetables & Ornamentals

**ACTIVE INGREDIENT:**  
 Sulphur..... 93.0 %  
**INERT INGREDIENTS .. 7.0 %**

**TOTAL..... 100.0 %**

**FINESS: 90% passing 325 screen**

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store unopened product in original containers only in cool, dry area out of reach of children and animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

If on skin, wash immediately with soap and water. Get medical attention.

If in eyes, immediately flush eyes with plenty of clear water for at least 15 minutes. Get medical attention if irritation persists.

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

See Side Panel For Additional Precautionary Statements.

**GENERAL INFORMATION**

Apply this product only as specified on this label. **DO NOT** use this product for any uses other than those specified on this label.

Superfine Dusting Sulphur is formulated for use on vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined with most of the commonly used fungicides & insecticides in the dry form. Apply to all plant surfaces to thoroughly cover.

EPA REG. NO. 99665- EPA EST. NO. 99669-TX-1

**INTERNATIONAL CHEMICALS, INC.**  
 P.O. Box 611 Mt. Pleasant, TX 77455

**NET CONTENTS 50 LBS**

**CAUTION: DO NOT** use sulphur in combination with oil or other three weeks of an oil spray. **DO NOT** use sulphur on sensitive crops when shade temperatures exceed, or likely to exceed 90°F. **DO NOT APPLY OR ALLOW** sulphur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tang. To do so may result in severe fruit and leaf injury.

When growing crops for processing, consult processors before applying sulfur.

**WARRANTY STATEMENT**

**INTERNATIONAL CHEMICALS, DC.** - Warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purpose set forth on the label when used according to directions under normal use conditions on the crops specified.

There are no other warranties, expressed or implied concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

**BEST AVAILABLE COPY**

RECOMMENDATIONS

Consult your State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

ROSES, SHRUBS, FLOWERS: (as a Fungicide & Insecticide)

Black Spot, Brown Canker, Leaf Spot, Powdery Mildew, Thrips & Nites on Roses. Begin dusting when new growth starts in the spring. Repeat at 7 day intervals and respray after rainfall. For disease and insect control on shrubs, flowers and ornamentals, begin application when diseases appear and repeat at 5-10 day intervals and after rains. For control of Mildew, Thrips and Nites on Azalea, Alder, Button bush, Cotoneaster, Crape Myrtle, Halia, Daisy, Delphinium, Aster, Columbine, Dogwood, Eucalyptus, Gladiolus, Golden Fleece, Hibiscus, Holly, Hydrangeas, Lady's Mantel, Ligustrum, Lilac, Petunia, Poplars, Silver Vine, Spirea, Sunflower, Sweet Pea, Verbena, Violets, Willows, Carnations, Cedar, Cherry Laurel, Chrysanthemums, Hollyhock, Juniper, Smilax & Spruce. For control of Leaf Spot, Thrips, & Nites on Calceola, Cedar, Cherry Laurel, Chrysanthemums, Clematis, Columbine, Fonglove, Hollyhock, English Ivy, Juniper, Laurel, Sage, Smilax, Snapdragon and Spruce. For control of Rust, Thrips, and Nites on Asters, Carnations, Chrysanthemums, Hollyhock and Snapdragon.

VEGETABLES: (as Fungicide)


For control of Powdery Mildew on Beans, Peas, Broccoli, Brussel Sprouts, Cabbage, Cauliflower. For control of Rust on Beans & Peas. Begin dusting at first sign of disease. Repeat applications at 7-10 day intervals. NOTE: Sulphur may injure certain varieties of beans. For control of Powdery and Downy Mildew and Botrytis Blight on Onions. Begin application before diseases are expected to appear. Repeat as necessary.

CHIGGER CONTROL: For control of chiggers in yard, dust over entire lawn, flower garden, shrubs and any place chiggers may bite. This dusting should be done a day or two before protection is desired and should be repeated at weekly intervals. The content of this package will cover from 20,000 to 25,000 sq. ft.

IMPORTANT: Use only according to state recommendations. When used as directed, this product will provide effective and economical control of listed diseases and insects and promote healthy bush growth.

IMPORTANT: Use the higher dosage during periods of frequent and heavy rainfall. Dust often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall.

[Insert]:  
Use 6 to 8 lbs. per acre.  
Keep new growth protected.



[Insert]:  
Use 16 lbs. per acre for thorough coverage.

[Insert]:  
Use approximately 100 lbs. per acre.

BEST AVAILABLE COPY