

RECOMMENDATIONS

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

All dosages given on this label are for dilute sprays (1% concentrations). For use in concentrate sprays, increase the amount of product by the concentrate factor. For example, if normal use is 5 lbs. per 100 gallons applied to one acre, the 5% concentrate rate will be 25 lbs. per 100 gallons applied to 5 acres. Spray often enough to keep new growth protected. **IMPORTANT:** Use the higher dosage during periods of frequent and heavy rainfall. Spray often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Superfine Wettable Sulphur, always add the same amount of Spray Line, or other safener, as would be used when applying Lead Arsenate without sulphur. When using Zinc Sulfate and Spray Line as a safener on peaches, always add Zinc Sulfate to the tank first, followed by Spray Line, Lead Arsenate and sulphur in order given.

APPLE/PEAR - Powdery Mildew, Scab: Use 6 to 8 lbs. per 100 gallons of water in the pre-bloom, bloom and petal fall sprays. Use 4 to 6 lbs. in the cover sprays.

CHERRY/PLUM/PRUNE - Brown Rot, Leaf Spot: Use 6 to 8 lbs per 100 gallons of water in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

CITRUS - Rust Mite: Use 5 lbs. per 100 gallons of water. Begin application at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russeting of fruit and greasy spot on leaves.

SCALE CRAWLERS: Add 5 to 8 lbs. to each 100 gallons of tank strength lime sulphur solution.

PEACH - Brown Rot, Scab: Use 6 to 8 lbs. per 100 gallons of water in pink, bloom, shuck, cover and pre-harvest sprays.

BEANS, PEAS - Powdery Mildew, Rust: Use 16 lbs per acre in sufficient water for thorough coverage. Repeat application in 7 to 14 days intervals as necessary.

ROSES - Black Spot, Powdery Mildew: Use 6 to 8 lbs per 100 gallons of water. (3 to 4 Tablespoons per 1 gallon of water). Make first application prior to first signs of disease and maintain preventative schedule through out the season repeating application every 7 to 14 days. Keep new growth protected.

GENERAL PURPOSES - Powdery Mildew, Rust: Use 6 to 8 lbs per 100 gallons of water. (3 to 4 Tablespoons per 1 gallon of water). Make first application prior to first sign of disease and maintain preventative schedule through out the season, repeating applications every 7 to 14 days. Keep new growth protected.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION
 Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS
 This product is toxic to fish. Do not apply directly to water. Do not apply where runoff conditions are likely to occur. Do not apply where weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS
 Sulfur dust suspended in air ignites easily. Keep away from heat, sparks and flame. Do not smoke while applying this product.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


RE-ENTRY STATEMENT
 Do not enter treated area without protective clothing and respiratory protection for 24 hours following application. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:
CAUTION: Area treated with **SUPERFINE WETTABLE SULPHUR**
 Do not enter without appropriate protective clothing for 24 hours following application. In case of accidental exposure, see the Statement for Practical Treatment on this label.

STORAGE AND DISPOSAL
 Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store unopened product in original container only in cool, dry area out of reach of children and animals.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loose clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

GENERAL INFORMATION
 Apply this product only as specified on this label. Do not use this product for any uses other than those specified on this label. Superfine Wettable Sulphur is formulated for use in water as a spray on fruit, vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined in the spray tank with most of the commonly used fungicides and insecticides. Pour the specific amount of this product on surface of water in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants.

CAUTION: Do not use sulphur in combination with oil or within three feet of an oil spray. Do not use sulphur on apples or other sensitive crops when shade temperatures exceed, or likely to exceed 90°F. Do not apply or allow sulphur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tung. To do so may result in severe fruit and leaf injury.

BEST AVAILABLE COPY



For Spraying Fruits
 Vegetables & Ornamentals

ACTIVE INGREDIENT:
 Sulphur 94.0 %
INERT INGREDIENTS .. 6.0 %

TOTAL..... 100.0 %

FINENESS: 95% passing 325 screen

KEEP OUT OF REACH OF CHILDREN
CAUTION
STATEMENT OF PRACTICAL TREATMENT
 If on skin, wash immediately with soap and water. Get medical attention.
 If in eyes, immediately flush eyes with plenty of clear water for at least 15 minutes. Get medical attention if irritation persists.
 If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
 See Side Panel For Additional Precautionary Statements.

EPA REG. NO. 49668- EPA EST. NO. 49668-TX-1
INTERNATIONAL CHEMICALS, INC.
 P.O. Box 6118 ML Pleasant, TX 75466
NET CONTENTS 50 LBS

WARRANTY STATEMENT
INTERNATIONAL CHEMICALS, INC. warrants that this product in its unopened package conforms to the chemical description on the label and is reasonable fit for the purposes set forth on the label when used according to directions under normal conditions on the crops specified. There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instruction or under abnormal conditions or conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

[Insert] - When growing crops for processing, consult the processor before applying sulfur.

ACCEPTED
 JUN 22 1987
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 49668-2