

COPY

International Chemicals, Inc
c/o Peggy M. Galloway
Landis Associates, Inc.
3025 Madison Highway
P.O. Box 5126
Valdosta, GA 31603-5126

253282	300
3	18

Gentlemen

Subject: Add Homeowner Use Pattern
Redball Superfine Wettable Sulfur
EPA Registration No. 49668-2
Your Submission Dated October 18, 1989

The amendment referred to above, submitted in connection with registration under PIFRA is acceptable, provided that you:

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

SL
Susan Lewis
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (H7505C)

SYMBOL	SURNAME	DATE	CONCURRENCES						
55185-1	Stone	R-13	KENCO	03/29/90	04/29/90	ka	vo	dd	dg



**SUPERFINE
WETTABLE
SULPHUR**

For
Spraying
Fruits Vegetables
& Ornamentals

ACTIVE INGREDIENTS

Sulphur	94.0%
INERT INGREDIENT	6.0%
TOTAL	100.0%

FINENESS: 95% passing 325 screen

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

Wash skin with soap and water.
In case of chemical contact with eyes, flush eyes with plenty of clear water for at least 15 minutes. Call a physician if irritation persists.
If swallowed, drink 1 or 2 glasses of water and induce vomiting by pumping bag. Do not treat with ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
See Back Panel For Additional Precautionary Statements.

EPA REG. NO. 13668-2 EPA EST. NO. 49668-TX-1
INTERNATIONAL CHEMICALS, INC.
P. O. Box 611 • MI. Pleasant, TX • 75455

NET CONTENTS 2 LBS.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 03 1990

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide
registered under EPA Reg. No.

49668-2

GENERAL INFORMATION

Apply this product only as specified on this label. Do not use this product for any uses other than those specified on this label. Superfine Wettable Sulphur is formulated for use in water as a spray on fruit, vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined in the spray tank with most of the commonly used fungicides and insecticides. Pour the specific amount of this product on surface of water in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants.
CAUTION: Do not use sulphur in combination with oil or within three weeks of an oil spray. Do not use sulphur on apples or other sensitive crops with shade temperatures exceed, or likely to exceed 90° F. Do not apply or allow sulphur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tung. To do so may result in severe fruit and leaf injury. When growing crops for processing, consult the processor before applying sulphur.

RECOMMENDATIONS

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

All dosages given on this label are for dilute sprays (1x concentrations). For use in concentrate sprays increase the amount of product by the concentrate factor. For example if normal users 1 lb. per 20 gallons applied to 9,000 sq. ft. the 5x concentrate rate will be 5 lbs. per 20 gallons applied to 1 acre. Spray often enough to keep new growth protected.

IMPORTANT: Use the higher dosage during periods of frequent and heavy rainfall. Spray often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Superfine Wettable Sulphur, always add the same amount of Spray Lime or other safener as would be used when applying Lead Arsenate without sulphur. When using Zinc Sulfate and Spray Lime as a safener on peaches, always add Zinc Sulfate to the

BEST AVAILABLE COPY

INTERNATIONAL CHEMICALS, INC.
P.O. Box 611 • ML Pleasant, TX • 75455

NET CONTENTS 2 LBS.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly in water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. *When disposing of equipment washwaters.*

PHYSICAL AND CHEMICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks and flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION: Area treated with SUPERFINE WETTABLE SULPHUR. Do not enter without appropriate protective clothing for 24 hours following application. In case of accidental exposure, see the Statement for Practical Treatment on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store unused product in original container, tightly closed, and out of reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Empty containers may be recycled. Do not reuse. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.



Do not reuse empty container. Rinse thoroughly before discarding in trash.

~~weekly or bi weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Superfine Wettable Sulphur, always add the same amount of Spray Lime, or other satener, as would be used when applying Lead Arsenate without sulphur. When using Zinc Sulfate and Spray Lime as a satener on peaches, always add Zinc Sulfate to the~~

tank first, followed by Spray Lime, Lead Arsenate and sulphur in order given. **APPLES/PEAR:** Powdery Mildew, Scab Use 1/3 to 1/2 lb per 6 gallons of water in pink bloom, petal fall, shuck, cover and post harvest sprays.

Citrus-Rust Mite: Use 1 lb per 20 gallons of water. Begin application at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russetting of fruit and greasy spot on leaves. **Scale Crawlers:** Add 1 to 1 1/2 lbs per 20 gallons of tank strength lime sulphur solution.

PEACH-Brown Rot, Scab: Use 1/3 to 1 1/2 lbs per 20 gallons of water in pink bloom, shuck, cover and pre harvest sprays.

BEAN, PEAS-Powdery Mildew, Rust: Use 2 lbs per 5,500 sq ft in sufficient water for thorough coverage. Repeat application in 7 to 14 day intervals as necessary.

ROSES-Black Spot, Powdery Mildew: Use 1/3 to 1 1/2 lbs per 6 gallons of water (3 to 4 tablespoons per 1 gallon of water). Make first application prior to first signs of disease and maintain preventative schedule through out the season repeating application every 7 to 14 days. Keep new growth protected.

PERENNIALS, SHRUBS-Powdery Mildew, Rust: Use 1/3 to 1 1/2 lbs per 6 gallons of water (3 to 4 teaspoons per 1 gallon water). Make first application prior to first sign of disease and maintain preventative schedule through out the season, repeating applications every 7 to 14 days. Keep new growth protected.

WARRANTY STATEMENT

INTERNATIONAL CHEMICALS, INC. - Warrants that this product in its unopened package conforms to the chemical description on the label and is reasonable fit for the purpose set forth on the label when used according to directions, under normal use conditions on the crops specified. There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instruction or under abnormal conditions or conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

NOT AVAILABLE ANY