

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Use of this product in a manner not in accordance with the information on this label may injure domestic animals, wildlife, aquatic life or produce other undesirable results.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

(see attached statement)

STORAGE AND DISPOSAL

Product may be stored and if frozen may generally be restored after thawing and mixing. Repeated freezing/thawing may impair utility. Do not reuse container. Dispose by burying in area away from water supply.

PRODUCT INFORMATION

Sunbelt Flowable Sulfur is a flowable sulfur formulation that may be applied as a ground or aerial application. Consult your State Agricultural Experiment Station or Extension Specialist for advice in selecting treatments from this label to best fit local conditions.

USE ONLY ON CROPS LISTED AND ONLY AS SPECIFIED ON THIS LABEL. EXCEPT ON COTTON, do not apply when temperatures exceed or are likely to exceed 90°F. Do not use sulfur with oil or within 14 days of an oil spray (for citrus 21 days). Do not combine with emulsifiable liquids.

NOTE: Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply or allow to drift to apricots, (Anjou and Comice pears, cranberries, cucurbits (cucumber, cantaloupe, melons, squash, I fibers, spinach, lung trees, walnuts or other sensitive plants. Sulfur may burn foliage and fruit during periods of high moisture and under certain climate conditions.

Sunbelt Flowable Sulfur can be mixed and applied with liquid fertilizer or water.

DO NOT STOP AGITATION WITH SULFUR IN SPRAY EQUIPMENT. SULFUR IS HIGHLY CORROSIVE, AND EQUIPMENT SHOULD BE CLEANED THOROUGHLY AFTER EACH DAY'S SPRAYING.

RECOMMENDATIONS:

FRUIT TREE CROPS: Unless otherwise specified for specific crops, dosage rates are given as pints of Sunbelt Flowable Sulfur per 100 gallons water for use in a thorough coverage spray. The total amount of Sunbelt Flowable Sulfur per acre should be 1 to 4 gallons depending on the size of the trees. Because of variations in the types of spray equipment used, a range of low and high rates is listed. For high volume sprayers (about 800 to 1000 gallons spray per acre) use the low rate for low volume sprayers (about 20 to 300 gallons spray per acre) use the high rate per 100 gallons.

Apply in sufficient water for thorough coverage. For aerial application, use a minimum of 5 gallons of finished spray per acre.

ACCEPTED

SEP 30 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 49832-5

see attached replacement

Sunbelt

FLOWABLE SULFUR

ACTIVE INGREDIENT:

Sulfur as S 53.0%

INERT INGREDIENTS: 47.0%

TOTAL: 100.0%

This product contains 8 lbs. of Sulfur per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Statement of Practical Treatment

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If on skin, wash thoroughly with soap and water. Get medical attention.

If in eyes, wash eyes with plenty of clear water for at least 15 minutes.

See side panel for additional precautionary statements.

E.P.A. Est. No. 49832-AL-1

E.P.A. Reg. No. 49832-

NET
CONTENTS:

MANUFACTURED BY:

SUNBELT CHEMICALS, INC.

P. O. BOX 5
ATMORE, AL 36504

APPLE, PEAR:

Scab: Pre-bloom through calyx sprays 1 1/2 to 3 1/2 pints
Cover Sprays 1/2 to 2 pints

Powdery Mildew:

Pre-bloom through calyx sprays 3 pints
Cover Sprays 2 pints

CHERRY, PLUM, PRUNE:

Brown Rot, leafspot: 1 1/2 to 3 1/2 pints
Pink and Bloom Sprays 1/2 to 1 1/2 pints
Petiole fall, shuck & cover sprays

CITRUS:

Rust Mite, clove mite: 1/2 to 4 pints

Apply sprays November thru May

THRIPS (Early spring treatment for the reduction of Thrips population). Apply 1 to 3 gallons per acre in sufficient water for thorough coverage. Make application after spring flush has made 3 to 4 inch growth. Repeat if necessary. Do not apply within 21 days of an oil application.

PEACH, NECTARINE:

Powdery Mildew, brown rot, scab: 1 1/2 to 3 1/2 pints

Pink and Bloom Sprays: 1/2 to 1 1/2 pints

Petal fall, shuck and cover sprays

Application to mature nectarines may cause discoloration.

FIELD AND VINE CROPS:

Unless otherwise specified for specific crops, dosage rates are given as pints of Sunbelt Flowable Sulfur per acre. This product should be applied by ground equipment or aircraft in sufficient water to provide thorough coverage.

For aerial application, use a minimum of 5 gallons of finished spray per acre.

IMPORTANT: Use the higher dosage during periods of frequent and heavy rainfall. Apply often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall.

COTTON: Atlantic Mite. 2 pints for early season control and up to 1 gallon for mid to late season control.

GRAPES: Powdery Mildew. Use 1 to 2 pints per 100 gallons water, maximum of 8 pints Sunbelt Flowable Sulfur per acre in spring and summer. Make first application when shoots are 6 to 8 inches long, second when shoots are 12 to 15 inches long, and third at about bloom stage. The timing, rate and number of applications will vary with local conditions. Consult State Agricultural Extension Service or State Agricultural Experiment Stations in your region for specific information. Follow the USE PRECAUTIONS given on this label.

PEANUTS: For rust use 2 to 4 pints per acre of Sunbelt Flowable Sulfur in enough water for good coverage. Begin 40 to 60 days after planting. Use 7 to 14 day intervals as long as needed to control rust.

POTATOES: Powdery Mildew. Use 3 to 4 pints in 8 gallons water per acre by air. Apply at the first sign of infection and repeat at 3 week intervals.

SUGAR BEETS: For control of powdery mildew apply the recommended rate in 10 gallons of water per acre by air at first sign of infection. Use 8 pints per acre in Arizona and 4 pints per acre in other states. Repeat applications should be made at 10 to 30 day intervals or as necessary throughout the season. Under most disease conditions only one or two applications will be needed.

STRAWBERRIES: Powdery Mildew. Use 2 pints. Apply at first sign of infection and repeat at 3 week intervals. For Strawberries that will be canned, discontinue application of sulfur well before harvest in accordance with local canners' recommendations.

TOMATOES: Tomato russet mite. Use 1/2 to 1 gallon per acre depending upon size of plants. Use full coverage sprays.

WARRANTY STATEMENT

Sunbelt Chemicals, Inc. warrants that this product in its unopened package conforms to the chemical description on the label and is reasonably fit for the purposes set forth on the label when used according to directions under normal use conditions.

There are no other warranties, expressed or implied, concerning the use of this product other than indicated on the label. This warranty does not extend to the handling or use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes all risk of any such use.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS
& DOMESTIC ANIMALS

CAUTION


Harmful if swallowed, inhaled, or absorbed through skin. May cause irritation of nose, throat, or skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff conditions are likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, and flame. Do not smoke while applying this product.



RE-ENTRY STATEMENT

Do not enter treated area without protective clothing and respiratory protection for 24 hours following application.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION: Area treated with SUNBELT FLOWABLE
SULFUR on (date of application).

Do not enter without appropriate protective clothing for 24 hours following application.
In case of accidental exposure, see the Statement for Practical Treatment on this label.

