

U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7511C)
1200 Pennsylvania Avenue NW
Washington, DC 20460

NOTICE OF PESTICIDE:

X Registration

___ Reregistration

(under FUFRA, as amended)

| EPA | Reg. | Num | ber: |
|-----|------|-----|------|
| | | | |

Date of Issuance:

49911-3

6/20/0/

Term of Issuance:

Conditional

Name of Pesticide Product:

CARPOVIRUSINE

Name and Address of Registrant (include ZIP Code):

Sumitomo Corporation of America

500 Third Avenue

New York, New York 19016

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above hamed pesticide is hereby registered/reregistered under the Federal Insecticide. Fundicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement in recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The appearance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA Sec. 3(c)(7)(A) provided you:

- 1. Submit and/or cite all data required for registration, reregistration of your product under FIFRA section 3(c)(5) and section 4 when the Agency requires all registrants of similar products to submit such data.
- 2. Make the following label changes listed below before you release the product for shipment: a. Revise the label to read EPA Reg. No. 49911-3
- 3. Submit five(5) copies of the revised final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.
- 4. Submit a Cell Culture Study within one year from the date of registration for this product.

Signature of Approving Official:

(See second page for signature)

Date

6/20/01

EPA Form \$570-6

7571C L.Hollis 16-14-01 Lutton 6/15

Page 2

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

Janet L. Andersen, Director

Kamt L. andwer

Biopesticides and Pollution Prevention Division

M-I-D-D-L-E P-A-N-E-L OF D-R-A-F-T L-A-B-E-L

CARPOVIRUSINE A selective larvicide against the Codling moth

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.

| FIRST AID | | |
|--------------------------------------|--|--|
| If in eyes | ■In case of contact with eyes, hold eyelids open and rinse eyes with a steady gentle stream of water for at least 15 minutes. ■Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. ■Get medical attention if irritation persists. | |
| If on skin or clothing | If on skin or clothing, remove contaminated clothing. Wash hands and exposed skin thoroughly with soap and water. Get medical attention if irritation occurs. | |
| Have the production doctor, or going | ct container or label with you when you are calling a poison control center or for treatment. | |

SUMITOMO CORPORATION OF AMERICA

600 Third Avenue New York, NY 10016

Manufactured by
Natural Plant Protection
Route d'Artix
64150 Noguères
France

EPA Registration No. 49911-EPA Establishment No. 070057-FRA-001 with COMMENTS In EPA Letter Dated
JUN 2 0 2001

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Net volume: 1 liter (1.0567 quart)

R-I-G-H-T P-A-N-E-L OF D-R-A-F-T L-A-B-E-L

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Agricultural Use Requirements

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours. Workers may be allowed early entry to treated areas if the following required PPE is worn:

- ■Protective eyewear such as goggles or full face mask
- Long sleeve shirt, long pants
- Shoes and socks

For treatment against the Codling moth in apple, pear and walnut orchards.

Dilute one liter of Carpovirusine in 1000 liters of water with agitation (1 quart in 250-264 gallons of water). Apply diluted Carpovirusine with any conventional sprayer at the beginning of egg hatch of the first generation (to be determined by the use of sex pheromone traps) or follow warnings given by Extension Offices. Repeat application every 10 to 12 days as long as oviposition is evident, and while adults are captured in sex pheromone traps. 1000 liters (264 gallons) of diluted product can treat one hectare (2.47-2.5 acres).

Carpovirusine is best used in an Integrated Pest Management Program.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal

Store in original container in a cool, dry place. Store at 15-20°C (59-68°F) for use during the current year; 4°C (39°F) for storage periods greater than one year.

Pesticide disposal:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Plastic container disposal:

Triple rinse (or equivalent). Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning.

L-E-F-T P-A-N-E-L OF D-R-A-F-T L-A-B-E-L

Precautionary Statements:

Hazards to humans and domestic animals

- Avoid contact with eyes as product may cause irritation.
- Avoid contact with skin and clothing, and avoid breathing spray mist.
- ■Do not eat, drink or smoke during application.
- •Keep away from food and animal feedstuffs.

Personal Protective Equipment:

Applicators and other handlers must wear:

- ■Protective eyewear such as goggles or protective face shield
- ■Long-sleeved shirt and long pants
- ■Shoes and socks

Wash PPE clothing with detergent and hot water, separately from other laundry.

User safety recommendation:

Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Environmental Hazards:

Do not dispose of equipment washwaters or rinsates in ponds, rivers or where surface water is present.