

49991-1

11/13/2002

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

NOV 13 2002

Dr. H. R. McLane
H. R. MCLANE, Inc.
Agent for Bonus Crop Fertilizer, Inc.
7210 RED ROAD, Suite 206
Miami, Florida 33143-5321

Dear Dr. McLane:

Subject: Turf Fertilizer with Barricade® 20/75
EPA Registration Number 49991-1
Applications and Letters Dated October 23, 2002
Request To Amend Register To Comply with PR Notices
2001-1 and 2001-6

Your request to amend the subject labeling for this pesticide product has been reviewed and found acceptable under the Federal Insecticide, Fungicide and Rodenticide Act, (FIFRA) as amended, provided that you:

1. Submit one label and respective Confidential Statement of Formula that reflects each percent active ingredient (a.i.) between 0.20 and 0.75% a.i. for each product shipped under this registration. As this amendment is limited to the 5 formula stated on your application, you may only submit amended labeling for each of those products prior to shipment under this amendment of registration.
2. Submit one (1) copy of the final printed labeling submitted before you release these products for shipment under the subject labeling.

If these conditions are not complied with, the registration

-2-

will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of these products under this labeling constitutes acceptance of these conditions.

A stamped copy of the proposed labeling is enclosed for your records.

Sincerely yours,

for Eugene M. Wilson
Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

Turf Fertilizer with Barricade® 20 / 75

Label copy
unit 1 of 2

Contains (0.20%-0.75%) Barricade®

20 Pounds Treats (up to 2,680 sq.ft.) (up to 10,000 sq.ft.)

3/4

ACTIVE INGREDIENT:

Prodiamine [N,N-Di-*n*-propyl-2,4-dinitro-6-(trifluoromethyl)-*m*-phenylenediamine] 0.20% - 0.75%
INERT INGREDIENTS: - - - 99.80% - 99.25%

Total 100.00% - 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.
Barricade® is a Registered Trademark of Syngenta Corp..

EPA Reg. No. 49991-1 EPA Est. No. 49991-TX-1

FIRST AID

IF SWALLOWED: - Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: - Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: - Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: - Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice. Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment advice. See side panel for additional Precautionary Statements.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal. **PRODUCT DISPOSAL:** As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. **IF EMPTY:** Do not reuse this bag. Place in trash or offer for recycling if available. **IF PARTLY FILLED:** Call your local waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **SPILL:** In case of spill, sweep up and dispose of material according to PRODUCT DISPOSAL directions above.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, * (landscape ornamentals, established perennials and wildflower plantings). It controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application. 10-23-02

Not for use on plants being grown for: (1) Sale or other commercial use; (2) for commercial seed production or; (3) for research. For use on plants intended for esthetic purposes or climatic modification and being grown in ornamental gardens, or parks, or on golf courses, or lawns and grounds. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.


Application Directions

Apply uniformly with suitable, calibrated application equipment. (continued on next column)

For Selective Preemergence Control of Grass and Broadleaf Weeds in Established Turf Grasses, (excluding golf course putting greens), and Lawns - ** For around Landscape Ornamentals Including Established Perennials and Wildflower Plantings.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.



USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns.

(1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

(2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

(3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

(4) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as diking may result in reduced weed control. (6) Do not apply this product to putting greens, or areas where dikandra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. It will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

Maximum Annual Rates

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate of active ingredient (a.i.) per acre as given in the following table.

Maximum Application Rate per Calendar Year of Product by Turf Grass Species¹

Turf Species	Lbs. Product per Acre	Lbs. a.i. / Acre
	0.20% a.i.	0.75% a.i.
Creeping Bentgrass	325 lbs.	86 lbs.
Creeping Red Fescue	375 lbs.	100 lbs.
Buffalograss	500 lbs.	133 lbs.
Kentucky Bluegrass		
Perennial Ryegrass		
Bermudagrass ²	750 lbs.	200 lbs.
Bahiagrass		
Centipede grass		
Seashore Paspalum		
St. Augustinegrass		
Tall Fescue		
Including turf type		
Zoysia		

¹These are the maximum rates per calendar year by species limitations. ²May be used on newly-sprigged or plugged Bermudagrass at rates not to exceed 0.50 lbs. a.i./acre (equal to xx lbs./acre of this product). Newly-sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations. Do not apply more than 1.50 lbs. of a.i. per calendar year (equal to xxx lbs. of this product). Use higher rates of product to achieve longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates section.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds.

Barnyardgrass	Goosegrass ¹	Pigweed
Bluegrass, Annual	Herbit	Purslane, common
(<i>Poa Annua</i>)	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass ²
Chickweed, Common	Jungleice	Shepherd's Purse ²
Chickweed, Mouse ear	Knockweed	Signalgrass, Broadleaf
(from seed)	Kochia	Speedwell, Persian
Crabgrass (Large, Smooth)	Lambsquarters, common	Sprangleop
Crowfootgrass	Lovegrass	Spurge, Prostrate
Cupgrass, Woolly	Panicum (Texas, Fall, Browntop)	Witchgrass
Foxtails, Annual		Woodsorrel, Yellow,
For weed reference numbers ³ refer to the following sentences.		from seed.

(1) In many areas a single application of 0.65 lb. to 1.5 lbs. active ingredient per acre (equal to xx lbs. to xxx lbs. of this product per acre) will control Goosegrass. However under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lbs. to 1.0 lb. a.i. per acre (equal to xx lbs. to xxx lbs. of this product per acre) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rates section. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate section.

(2) Application for this weed should be made in late summer, fall, or winter prior to germination (3) Suppression Only. Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Product Rate lbs./Acre			Months Before Overseeding		
0.20%	0.75%	a.i./A	North	Transition	South
250	67	0.50	4 mos.	4 mos.	4 mos.
325	86	0.65	5 mos.	4 mos.	4 mos.
375	100	0.75	6 mos.	5 mos.	5 mos.
400	106	0.80	-	6 mos.	6 mos.
500	133	1.00	-	7 mos.	7 mos.
570	152	1.14	-	-	9 mos.
650	173	1.30	-	-	10 mos.
750	200	1.50	-	-	12 mos.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	variable %
Available Phosphoric Acid (P ₂ O ₅)	variable %
Soluble Potash (K ₂ O)	variable %
Ingredients: Ureaformaldehyde, Urea, Sulfur Coated Urea, Ammonium Phosphate, Muriate of Potash	

SPREADER SETTINGS FOR 20 LBS.	2,680 SQ.FT.	10,000 SQ.FT.
Scott's Evergreen 18"	# XX	# XX
Scott's Speedy Brcdst	# XX	# XX
Red Devil Broadcast	# XX	# XX
Republic EZ Drop 18"	# XX	# XX
Republic EZ Broadcast	# XX	# XX
Republic EZ Trctr. Brcdst	# XX	# XX

**For use of this product on landscape ornamental plants, including established perennials and Wildflower Plantings, refer to this product Supplemental label for complete instructions and listings of the tolerant ornamental plant species.

ACCEPTED
with COMMENTS
In EPA Letter Dated
NOV 13 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
49991-1

TURF FERTILIZER WITH BARRICADE® 20/75

Common Name

Big Blue Lily; Turf; Silver Mound; Evergreen Giant Liriope, Variegated

Liriope, Creeping
Cardinal Flower, Indian Pink
Japanese Honeysuckle
Tatarian Honeysuckle
Burgundy
Loosestrife; Modern Pink
Magnolia species**
Ice Plant
Crabapple
Yaku Jima, Silberfeder
Heavenly Bamboo
Narcissus species**
Oleander
Olive
Mondo Grass
Osmanthus (False Holly):
Gulf Tide
Trailing African Daisy
Sourwood
Tree Peony
Fountain Grass (Dwarf)

Avocado*
Fraser's Photinia (Redtip)
Dragonhead, False; Vivid Spruce species** (Colorado Blue, Norway, etc.)
Lily-of-the-Valley Bush
Calabrian Pine
Canary Island Pine
Slash Pine
Aleppo Pine
Austrian Pine
Longleaf Pine
Monterey Pine
Japanese Black Pine
Eastern White Pine
Scotch Pine
Loblolly Pine
Virginia Pine
Pistachio*
Queensland Pittosporum
Japanese Pittosporum
Japanese Yew
English Laurel
Almond, Apricot, Nectarine, Peach, Plum and Prune*
Douglas Fir**
Firethorn (Scarlet)
Firethorn
Firethorn
Bradford Pear sp.
Oak, Shumard's Red
Oak species
Indian Hawthorn
Yedda Hawthorn
Coral Bells
Delaware Valley White
Flame Creeper
Formosa
Girard Crimson
George L. Tabor
Hino-crimson
PJM
Roseum Elegans
Wakelibus
White Gumbo
Lady Bank's Rose
Rosemary
Black-Eyed Susan
Goldstrum Santolina
Saxifrage; Purple Dome
Pincushion Flower
Stonecrop
Stonecrop; Lidakense
Stonecrop
Stonecrop; Dragon's Blood
Spiraea
Anthony Waterer
Australian Brushcherry
Japanese Yew
Yew
Germander
Meadow Rue
American Arborvitae
Star Jasmine
Canada Hemlock
Tulip species
Veronica, Speedwell; Sunny

Japanese Viburnum
Sweet Viburnum
Japanese Snowball
Cane Island Viburnum
Arrowood Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Greater Periwinkle, Vinca
Common Periwinkle, Vinca
Crape
Old Fashioned Weigela
Saxifrage Bayonet, Yucca
Adam's Needle, Yucca

Scientific Name

Liriope muscari

Liriope Muscari var. variegata

Liriope spicata
Lobelia Cardinalis
Lonicera japonica
L. tatarica
Loropetalum chinense
Lythrum sp.
Magnolia spp.
Maleophora luteola
Malus sp.
Miscanthus sinensis
Nandina domestica
Narcissus spp.
Nerium oleander
Olea europaea
Ophiopogon japonicus
Osmanthus heterophyllus

Osteospermum fruticosum
Oxydendrum arboreum
Paeonia suffruticosa
Pennisetum setaceum
Perovskia atriplicifolia
Persea americana
Photinia fraseri
Physotegia virginiana
Picea spp.

Pieris japonica
Pinus brutia
P. canariensis
P. elliotii
P. halepensis
P. nigra
P. palustris
P. radiata
P. thunbergiana
P. strobus
P. sylvestris
P. taeda
P. virginiana
Pistachio sp.
Pittosporum rhombifolium
P. tobira
Podocarpus macrophyllus
Prunus laurocerasus
Prunus sp.

Pseudotsuga menziesii
Pyracantha coccinea
P. fortuneana
P. koidzumii
Pyrus sp.
Quercus shumardii
Quercus spp.
Raphiolepis indica
Raphiolepis umbellata
Rhododendron
(including Azalea)

Rosa banksiae
Rosmarinus officinalis
Rudbeckia sp.
Santolina, virens
Saxifraga sp.
Scabiosa sp.
Sedum album
Sedum caudicicola
Sedum dasyphyllum
Sedum spumum
Spiraea Bumalda
Syzygium paniculatum
Taxus cuspidata
T. media
Tauricum sp.
Thalictrum dipterocarpum
Thuja occidentalis
Trachelospermum asiaticum
Tsuga canadensis
Tulipa spp.
Veronica sp.

Viburnum japonicum
V. odoratissimum
V. plicatum
V. rigidum
V. suspensum
V. tinus
V. trilobum
V. wrightii
Vinca major
Vinca minor
Vitis sp.
Weigela florida
Yucca aloifolia
Y. filamentosa

Contains 0.20% - 0.75% Barricade® For Selective Preemergence Control of Grass and Broadleaf Weeds in Landscape Ornamentals Including Established Perennials and Wildflower Plantings.

EPA Reg. No. 49991-1 EPA Est. No. 49991-TX-1

LANDSCAPE ORNAMENTALS Including Established Perennials and Wildflower Plantings

This product may be applied for residual preemergence weed control in ornamentals.

Use Rates and When to Apply

Apply in fall and/or spring, 0.65 lb. to 1.5 lbs. active ingredient per acre, which is equal to x.xx to x.x lbs. of this product per 1,000 sq. ft. of ornamental plantings; therefore 20 lbs. of this product will treat xx.xxx to x.xxx sq. ft. Use the higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of (xxx lbs. of this product/acre), equal to 1.5 lb. of a.i./acre.

Application Timing and Information

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific Name

Common Name

Abies spp. Fir species (Balsam, Fraser, Noble, etc.)

Abelia grandiflora
Acer palmatum
A. platanoides
Achillea, sp.
Actinidia chinensis
Agapanthus africanus
Agapanthus orientalis
Akebia quinata
Allium cernuum
Anemone hybrida
Aquilegia sp.
Arctostaphylos densiflora
Arctostaphylos uva-ursi
Artemisia sp.
Aster sp.
Aster X Frikartii
Athyrium filix-femina
Aucuba japonica
Begonia sp.
Berberis gladiolifolia
B. julianae
B. mentorensis
B. thunbergii
B. verruculosa
Bergenia cordifolia
Boehmeria asteroides
Bougainvillea spp.
Buddleia davidii

Abelia: Sherwood
Japanese Maple
Norway Maple
Yarrow: King Edward
Kiwi
Lily-of-the-Nile (African Lily)
Five-Leaf or Chocolate Vine
Lady's Leek, Nodding Onion
Japanese Anemone
Aquilegia: Red and Gold
Theophrasti Manzanita
Cape Weed
Wormwood; Silver Mound, Castile
Aster: Bonny Blue, Purple Dome
Lady Fern; Fern Lady
Japanese Aucuba
Fibrous Begonia: Hardy Grandis
Barberry
Wintergreen Barberry
Mentor Barberry
Japanese Barberry
Warty Barberry
Snowbank
Bougainvillea
Butterfly-Bush (Dwarf Blue);
Royal Red

Scientific Name
Buxus microphylla
Callistemon viminalis
Calluna vulgaris
Campanula carpatica
Campis X Tagliabuana
Carpobrotus edulis
Cassia artemisoides
Ceanothus rigidus
Ceratostigma
plumbagonoides
Chamaecyparis pisifera
Chrysanthemum nipponicum
Cleyera japonica
Citrus spp.
Coreopsis sp.

Comus florida
C. stolonifera
Cortaderia selloana
Cotoneaster apiculatus
C. buxifolius
C. dammeri
C. microphyllus
Cretaeus spp.
Crocus spp.
Cupressus sempervirens
Delosperma alba
Delosperma sp.
Delphinium sp.
Dianthus deltoideus
D. gratianopolitanus
Dodonea viscosa
Echinacea purpurea
Elaeagnus pungens
Euonymus fortunei
E. japonica

E. kiautschovica
Fatsia japonica
Forsythia intermedia
F. suspensa
F. viridissima
Gaillardia sp.

Gardenia jasminoides
Gaura sp.
Gentiana dahurica
Geranium cinereum
Gladiolus spp.
Gypsophila repens
Hedera helix
Helianthemum sp.
Hemerocallis sp.

Hibiscus
Hibiscus sp.
Hibiscus rosa-sinensis
Hosta plantaginea
H. sieboldiana
Houttuynia cordata var. variegata
Hydrangea macrophylla
Ilex cornuta
I. crenata
I. opaca
I. pernyi
I. vomitoria
Inula ensifolia
Iris ensata
Iris sibirica
Iris spp.
Jasminum nudiflorum
Juniperus chinensis

J. conferta
J. davurica
J. horizontalis
Juglans sp.
Justicia brandegeana
Lagerstromia indica
Lagerstromia indica X
Lantana montevidensis
Lavender sp.
Leontopodium alpinum
Legustrum sinense
L. amurense
L. japonicum
L. lucidum
Lilium sp.

Snowbank
Bougainvillea
Butterfly-Bush (Dwarf Blue);
Royal Red

Common Name
Japanese Boxwood
Weeping Bottlebrush
Scotch Heather
Tussock Bellflower (White Clips)
Trumpet Creeper, Trumpet Flower, Madame Galen
Hottentot Fig (Ice Plant)
Wild Lilac

False Cypress

Cleyera
Citrus species*
Coreopsis (Calliopsis): Early Sunrise, Moonbeam
Flowering Dogwood
American Dogwood
Pampas Grass
Cranberry Cotoneaster
Cotoneaster
Bearberry Cotoneaster
Rockspray Cotoneaster
Hawthorne
Lucifer
Italian Cypress
White Trailing Ice Plant
Cooperi Pink
Larkspur: Blue Elf
Dianthus: Maiden Pinks "Zing"
Cheddar Pink
Hop Bush
Coneflower, Purple; Magnus
Silverberry
Wintercreeper
Japanese Spindle Tree (Evergreen Euonymus)

Spreading Euonymus
Japanese Araiia
Border Forsythia
Weeping Forsythia
Greenstem Forsythia
Gaillardia, Blanket Flower: "Goblin"
Gardenia, Cape-jasmine
Radicans

Gentian
Cranesbill
Gladiolus species**
Baby's Breath
English Ivy
Sunrose
Daylily: Aztec Gold, Stella De Oro, Tender Love
Rose of Sharon
Mallow: Disco Belle White
Chinese Hibiscus
Hosta, Plantain Lily (Fragrant)
Hosta, "Searsucker"

Bigleaf Hydrangea
Chinese Holly
Japanese Holly; Helleri
American Holly
Holly
Yaupon Holly; Schillings

Sword-Leaved Iris; Jodlesong
Siberian Iris; Cabernet
Iris species**
Winter Jasmine
Chinese Juniper; Nick's
Compact, Parsonii
Shore Juniper; Blue Pacific
Parsoni
Creeping Juniper
Walnut
Shrimp Plant
Crape Myrtle
Crape Myrtle; Tuscarora

Weeping Lantana
Lavender; Munstead
Edelweiss
Chinese Privet; Variegata
Amur Privet
Japanese Privet
Glossy Privet (Waxleaf)
Lily: Jazz

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Bonus Crop Fertilizer or the Seller. All such risks shall be assumed by the Buyer.

Bonus Crop Fertilizer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Direction for Use, subject to the inherent risks referred to above. Bonus Crop Fertilizer makes no other express or implied Warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Bonus Crop Fertilizer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Bonus Crop Fertilizer and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Bonus Crop Fertilizer.

Bonus Crop Fertilizer, Inc.
P.O. Box 1725 - Bay City, TX 77404