U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (##-467) AASHINGTON, C.C. 2-44	EPA REGISTRATION NO.	AUG 22 983
	TERM OF ISSUANCE	TERM OF ISSUANCE
NOTICE OF PESTICIDE: THE SECTION TO SERVICE	NAME OF PESTICIDE PRODUC	(T
(Under the Federal Insectivide Function and Rodentis (de Act. as anouded)		
AME AND ADDRESS OF REGISTRANT (Include ZIP cod)		
r	Т	
L_	7	
NOTE: Changes in labeling formula differing in substance submitted to and accepted by the Registration Division priproduct always refer to the above U.S. EPA registration nu	or to use of the label in commerc	n with this registration must be se. In any correspondence on th
On the basis of information furnished by the registrant, the he Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is hereb	by Registered/Reregistered und
A copy of the labeling accepted in connection with this R	egistration/Reregistration is reti	urned herewith.
health and the environment, the Administrator, on his moti icide in accordance with the Act. The acceptance of any n Act is not to be construed as giving the registrant a right by others.	same in connection with the regis	stration of a product under this
ATTACHMENT IS APPLICABLE		
MATURE OF APPROVING OFFICIAL	<del></del>	DATE

BP-220 BIOCIDE

ACCEPTED with COMMENTS in EPA Letter Dates

AUG 19 1983

Pril r the Petric to a netter Personne region : is unerded, a single con petric tune a life way for SCOSC - 2

ACTIVE INGREDIENT:	
Poly[oxyethylene (dimethyliminio) ethylene-	
(dimethyliminio) ethylene dichloride)	20.0.%
INERT INGREDIENT	80.0 %
This product contains 1.74 lb. of active ingredie 8.69 lb. per gallon.	ent per gallon and weighs

## **KEEP OUT OF REACH OF CHILDREN**

## **CAUTION**

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician in mediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds, or public water. For guidance, contact the regional office of the Environmental Protection Agency.

GERTAL CLASSIFICATION (				
It is a violation of Federal law to use this product in a manner in-				
consistent with its labeling.				
BP-220 BIOCIDE _	is used to control algae, bacteria,			
and fungi in recirculating commercial and industrial water cooling towers. Prior to				
its use, systems must be cleaned to remove algal growth, microbiological sline,				
and other deposits. An initial slug addition of $0.9$ , to $7.3$ fluid ounces				
ofBP=220_BTQCTDE per 1000 gallons of water				
to provide a concentration of 24 BP=220 BIOCIDE	, to , 60 parts per million of , , based on the total weight of water .			
in the system, is recommended. Repeat initial dosage until control is evident.				
Subsequent slug additions of $0.73$ to $7.3$ fluid ounces of				
	per 1000 gallons of water (.6			
to 60 parts per million of BP-2				
be employed every 2 to 5 days, or as needed. The frequency of addition depends				
upon the relative amount of bleedoff and the severity of the microbiological prob-				
lem. Slug additions should be made in the sump of water cooling towers.  BP=220_BIOCIDE is used to control bacteria in industrial.				
air-washing systems that maintain effective mist eliminating components. Prior				
to its use, systems should be cleaned to remove bacterial slime and other deposits. An initial slug dose of $11.1$ to $18.3$ fluid ounces of				
BP-220 Biocide per 1000 gallons of water is recom-				
mended. Repeat initial dosage until control is evident. Subsequent slug additions				
mended. Repeat initial dosage until control is evident. Subsequent slug additions of 7.5 to 18.3 fluid ounces of BP-220 BIOCIDE				
per 1000 gallons of water should be employed each 1 to 5 days, or as needed. The				
frequency of addition depends upon the relative amount of bleedoff and severity				
of the bucterial problem. Slug additions may be made to the sump or to the water				
collection trays of the airwash system.				
STORAGE & DISPOSAL: Koop con	tainer alexad when not in use. Do not			
STORAGE & DISPOSAL: Keep container closed when not in use. Do not				
contaminate water, food, or feed by storage, disposal or cleaning of equipment.  Rinsate that cannot be used or reprocessed should be disposed of in a landfill ap-				
proved for pesticides or buried in a safe place away from water supplies. Open				
dumping is prohibited.	sales and the sales and sales are			
·	☐ PLASTIC CONTAINERS: Do not			
rinse and offer for recycling,	reuse empty container. Triple rinse			
reconditioning, or disposal in an	and incinerate or dispose of in an			
approved landfill or bury in a safe	approved landfill or bury in safe in place.			
place.	piace.			
Manufactured by:				
Tricon Chemical Corporation				
8020 Fernham Lane				
Forustville, MD 20747				
EPA REG. NO EPA EST. NO				
NET CONTENTS				