

APR 3 1992

Mr. Brad Peirce
Wilson Laboratories, Inc.
36 Head Street
Dundas, Ontario L9H 3H3

Dear Mr. Peirce:

Subject: Revised Labeling
Wilson Lucky Sevin Dust
EPA Reg. No. 50383-16
Your resubmission of March 20, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling change given below before you release the product for shipment bearing the amended labeling.

Revise the Statement of Practical Treatment for oral exposure to read as follows: "IF SWALLOWED: Call a physician immediately. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Never give anything by mouth to an unconscious or convulsing person."

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide Rodenticide Branch
Registration Division (H7505C)

292

ACCEPTED
with COMMENTS
in EPA Label Label

APR 1992



LUCKY SEVIN* DUST

**USE ON: Fruit, Flowers,
Vegetables and Ornamentials**

**KILLS: Mexican Bean Beetles,
Bag Worm, Tent Caterpillar,
Chinch Bug, Aphid & 50 Other
Varieties of Insects**

ACTIVE INGREDIENTS:

Carbaryl 1-naphthyl N-methyl-
carbamate..... 5.00%
Inert ingredients 95.00%

SEVIN* Reg. TM of Rhorer, Poulenc Nederland, B.V. used
under licence by Wilson Laboratories Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION:
SEE SIDE PANELS FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

E.P.A. REG. No. 50383-16
E.P.A. EST. NO. 50383-CN-1

1 LB. Net Wt.



LUCKY SEVIN*

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling. Avoid contamination of food, feed, feeding troughs and water receptacles.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, call a physician immediately. Induce vomiting. If medical advice cannot be obtained, then take person and product container to the nearest medical emergency treatment centre or hospital. In case of eye contact wash eyes immediately with fresh water for at least 15 minutes. Get medical attention if irritation persists. For skin, wash with plenty of soap and water. If inhaled, remove person from exposure area.

NOTE TO PHYSICIANS: Emergency Information-Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

LUCKY SEVIN dust kills a wide range of insects both by contact and through the stomach. Apply a good duster as insects appear. Treat both upper and lower parts of foliage. Repeat at 7 to 10 day intervals or as needed. In general 1 lb. will treat a garden approx. 25' X 25'.

RAPES-To control Grape Leaf Folder, Grape Leafhopper, Grape Berry Moth, Japanese Beetles. Apply when insects appear, then as needed. Apply before first brood larvae emerge from rolls. Do not apply within 7 days of harvest.

STRAWBERRIES-To control Meadow spittlebug and Strawberry leaf roller. Apply in post bloom and summer sprays, repeat application as insects appear. Maybe applied up to one day of harvest: 1lb per 2000 sq. ft.

BEANS-To control Mexican Bean Beetles-1lb. per 1700 sq. ft. Leafhopper, Flea Beetle, Bean leaf beetle-1lb. per 1000 sq. ft.

Apply when insects appear including day of harvest. Do not use on meat animals.

TOMATOES, EGGPLANT, SQUASH-To control Flea beetle, melon worm, corn borer, cutworm, bug, leafhoppers- 1lb. per 2000 sq. ft. Apply when insects first appear and including day of harvest.

POTATOES-To control Colorado potato beetle, leafhoppers, flea beetles, etc. Apply when insects first appear and including the day of harvest.

CORN-To control Corn earworm, fall army worm-1lb. per 2000 sq. ft. Apply at 2 to 3 day intervals until silk begins to form. Timing and Coverage essential. Bees should be applied to day of harvest and including day of harvest.

LAWN INSECT CONTROL- Lawn moths, (sod webworm), etc. Apply after rain or watering and watering. Repeat in 2 or 3 weeks.

ORNAMENTALS AND FLOWERS- etc. - To control Japanese Beetle, Boxelder bug, Certain Leafhopper, Leaf rollers, etc. Apply when insects first appear.

STORAGE

STORAGE: Keep pesticide in original container. Avoid contact with food or drink containers. Avoid contact with children. Store in a cool, dry place.

DISPOSAL: PRODUCT- Dispose of by securely wrapping in newspaper and discard in trash.

CONTAINER:-Do not reuse.

NOTICE: Buyer assumes all responsibility for use in accordance with directions.

Manufactured by
WILSON LABORATORIES INC

Dundas, Ontario L9H 3H3

Code 7084