

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Remove pets, birds, and cover fish aquaria before spraying. Do not apply as a space spray while food processing is underway. Foods should be removed or covered before treatments. Thoroughly wash all food processing surfaces before reuse. In other food processing plants, if washing of equipment, benches, shelving, etc. where exposed food will be handled, is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as directed on the label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection. For rapid control of flies, fruit flies, gnats, mosquitoes, slipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths.

This use of the product in food processing or food handling establishments (bakeries, beverage plants, canneries, food processing plants, hospitals, hotels, industrial installations, meat packing plants, restaurants, schools, supermarkets and warehouses) should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before reusing.

Hospital patients should be removed for 2 hours and rooms aired before re-entry.

Close doors and windows and shut off ventilating systems. Use a mechanical aerosol generator or a fogger adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas to provide maximum penetration and impingement on the insects. Then treat as a space spray, directing the spray towards the ceiling and upper corners of the area and behind any obstructions, apply at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least one hour. Vacate treated area and ventilate before reoccupying. Repeat treat-

**PYRENONE FOOD FLANT
FOGGING INSECTICIDE**

Apply with Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

Designed for use in all areas of official establishments operating under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs.

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, Technical	5.0%
Petroleum Distillate	94.5%
	100.0%

*Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and 1.0% related compounds.

PYRENON - Registered trademark of Fairfield American Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panel for Additional Precautions

Manufactured By:

PAC-TEC, Inc.

P.O. Box 332

Cheshire, CT 06410

EPA Reg. No. 50429-3
EPA Est. No. 50429-CI-

ACCEPTED

FEB 25 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 50429-3

fish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetle, mealworm, grain mites and cadellies: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets around containers of store foods, around the base of machinery, behind shelves and drawers. For silverfish, spray book-cases, for afts, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and rugs. Make a localized application of floor and baseboards. Spray directly into cracks, closet, and infested areas of shelving. Repeat treatment as necessary.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Treat baseboards, moldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, molding, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally and empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining. Store clothes in moth tight containers or repeat treatments monthly.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed five miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy areas, stagnant pools or ponds and shore. Apply at a rate of 2 fluid ounces per acre. (0.2 fluid ounces per 1000 square feet.)

Using truck-mounted conventional misting equipment traveling at 5 m.p.h. to treat a 300 feet swath, calibrate equipment to deliver 0.23 fluid ounces per minute.

Buyer assumes all risk of use, storage or handling this material not in strict accordance with directions given herewith.

BATCH NO.