PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. When using this product in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when the facility is not in operation. Remove pets, birds and cover fish aquariums before spraying. Do not remain in treated areas and ventilate the area after treatment is completed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not spray near or toward an open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill. Consult Federal, state or disposal authorities for approved alternative procedures.

PYRENONE FOOD

PLANT SPRAY

A READY TO USE INSECTICIDE FOR FORGING AND SURFACE SPRAY

For Use in Institutions, Hospitals, Hotels, Restaurants, Stores, Theaters, Food Processing Plants, Bakeries, Breweries, Blotting Plants, and Warehouses.

The insecticidal components piperanyl butoxide and pyrethrins are classified as food additives and when this product is used as directed allowable tolerance levels will not be exceeded.

ACTIVE INGREDIENTS:

	Pyrethrins			0.20%
	Piperoryl	Butoxide	Technicalte	1.00%
				98.80%
				100.00%

* Equivalent to 0.8% (butylcarbityl) (6-propylipiperony) ether and 0.2% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

Harmful if swallowed. Do not spray near or toward an open flame. Avoid contamination of feed and foodstuffs.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PYRENONE - Fairfield American Corporation Trademark Registered.

Manufactured By:
PAC-TEC, Inc.
P.O. Box 332
Cheshire, CT 06410

DIRECTIONS FOR USE

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

SPACE SPRAY: To control houseflies, fruit flies, mosquitoes, small flying moths and gnats: Close doors and windows and shut off ventilating systems. Apply using a good sprayer adjusted to deliver a fine mist. Direct spray toward upper portions of the enclosure filling the room with mist. Use ½ to 1 ounce per 1000 cubic faspace.

To kill exposed stages of roaches, ants, spiders, clover mites, cheese mites, confused flour beetles, saw-toothed grain beetles, red flour beetles, rice weevils, granary weevils and grain mites: Treat as above using 2 to 3 ounces per 1000 cubic feet of space. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY: To control roaches, ants, spiders clover mites, confused flour beetles, saw-toothed grain beetles, rice weevils, granary weevils and grain mites: apply as a coarse mist directed toward insects, into cracks and crevices, along baseboards, around the base of machinery, containers of stored products and other where insects may hide. Spray any trails, wests and points of entry. Spray badly infested areas daily. In other areas, repeat treatment twice weekly or when reinfestation occurs.

