Net Wt. 20 L's. 9,1 kg

5,000 Sq. Ft.

# WEED&FED 10-6-4 with 2,4-D PLUS IRON

This product is designed for the control of Dandelion and many other Broadleal weeds, in Ornamental Turf.  $\S$ 

ACTIVE INGREDIEN 'S: Isooctyl ester of 2.4-Dichlorophenoxyacelic acid* INERT INGREDIENTS		
*Equivalent to 66% 2 4-Dichtorophenoxyacetic acid Isomer Specific by AOAC Method 6 D01~5	TOTAL 100.00%	
GUARANTEED ANALYSIS Total Nitrogen (N) Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) Soluble Potash (K <sub>2</sub> O)	6.00%	

KEEP OUT OF REACH OF CHILDREN

# CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA ESTABLISHMENT NO 50507-OH-02 EPA REGISTRATION NO

PSC Inc.
P.O.Box 96
Spencerville, Oh 100/2753

FRONT SIDE

### PRECAUTIONARY STATEMENTS

## HAZARDS TO HUHANS & DOMESTIC ANIMALS

# KEEP OUT OF REACH OF CHILDREN

### CAUTION

HARMFUL IF SWALLOWED Avoid contact with skin, eyes, or clothing Avoid breathing dust Wash thoroughly after handling

### STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with soap and water; for eyes flush with water for 15 minutes and get prompt medical attention.

# ENVIRONMENTAL HAZARDS

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from target area. Avoid application to susceptible plants (flowers, fruits, shrubs, etc.). Avoid drift of dust or vapors which may injure susceptible plants in immediate vicinity. Do not apply on windy days.

### STORAGE AND DISPOSAL

Store in original container only. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash. Do not burn.

### DIRECTIONS FOR USE

NOTE: It is a violation of federal law to use this product in a manner inconsistant with its labeling

Do not use on Lippia, St. Augustine, Bentgrass, or Dichondra. Do not re-seed for at least two weeks after application. Do not apply to a newly-seeded lawn until after 3 mowings. Do apply when weeds are young and active for best results. Applications when the grass is wet from dew, rain or sprinkling will be more effective. Do not water for at least 24 hours after application. If rain occurs during 24 hours after application, a second application may be necessary.

This product will control many broadleaf weeds such as:

Bahia	Panweed	Lambsquarters	Pigweed
Bindweed	Goldenrod	Hallow	Plantain
Buckhorn	Ground Ivy	Morning Glory	Purslane
Burdock	Heal All	Mustard	Ragweed
Chickweed	Henbit	Nettles	Smartweed
Chicory	Jewelweed	Oxalis	Speedwell
Cocklebur	Jimson Weed	Pennywort	Thistle
Dandelion	Kochia	Pepper Grass	Yarrow

# APPROXIMATE SPREADER SETTINGS

This bag will cover 5,000 (50 ft. x 100 ft.) sq. ft. Apply 20 lb. per 5,000 eq. ft. or 4 lb. per 1,000 of ornamental turf.

# RECOMMENDED TWO-STEP SETTINGS

HAKE	TWICE OVER	ONE-STEP SETTINGS	
Central	6 1/2 · 7	7 1/2 - 8	
Cyclone	3 1/2 - 3 3/4	4 - 4 1/2	
Greenview	6 1/2 - 7	7 1/2 - 8	
Ortho 3000	6 1/2 - 7 1/4	8 - 8 3/4	
Scotts Drop	4 1/2 - 5	5 1/2 - 6	
Sears	7 - 8	9 - 10	

Resubmitted 10/28/85

. . .