

CUPROQUIM
SUPPLEMENTAL LABELING
FOR DISTRIBUTION AND USE ONLY WITHIN
FLORIDA

BASIC COPPER 53
 EPA Reg. No. 45002-8

FOR "SHOT BAG" MIXES ON CITRUS

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Citrus

Basic Copper 53 may be mixed with dry foliar nutritionals (micronutrients) to create "Shot Bag" mixes to meet the various nutritional requirements of citrus and provide disease protection as described on this label. Basic Copper 53 per acre rates in these mixes must not exceed the maximum recommended labeled rates for disease control.

IMPORTANT

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. This labeling contains additional precautionary language which may not appear on the package label. Follow instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.



Cuproquim Corporation
P. O. Box 171357
Memphis, TN 38187-1357

17m 22 50534-2 Page 1 3 9

Net Contents: 30 Gallons/114 Liters

Net Contents: 55 Gallons/208 Liters

Daconate®

**MSMA Plus Surfactant For Postemergence Weed Control
in Cotton, Turf and Non-Crop Areas**

Product Contains 4.0 Pounds MSMA Per Gallon

Active Ingredient: Monosodium Acid Methanearsonate 35.2%

Inert Ingredients: 64.8%

Total: 100.0%

Total Arsenic, all in water-soluble form, expressed as elemental 16.3%

Keep Out of Reach of Children

CAUTION:

See side panel for additional precautionary statements.

ISK Biosciences Corporation

5966 Heisley Road

P.O. Box 8000

Mentor, Ohio 44061-8000

EPA Reg. No. 50534-2

EPA Est. No. 50534-TX-01

3197-0000-9053-9308

DACONATE® is a registered trademark of ISK Biosciences Corporation

APR 4 1985
EPA Reg. No. 50534-2
EPA Est. No. 50534-TX-01
50534-2

1 of 5

Precautionary Statements

Hazards to Humans and Domestic Animals

Caution

Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing vapors or spray mist.

If swallowed, induce vomiting immediately by giving two glasses of water and sticking finger down throat. NEVER give anything by mouth to an unconscious person. Seek medical attention.

In case of contact, flush eyes with plenty of water for at least 15 minutes. Seek medical attention.

Wash exposed skin gently with plenty of soap and water.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- * Long-sleeved shirt and long pants
- * Waterproof gloves
- * Shoes plus socks
- * Protective eyewear
- * Chemical-resistant apron when cleaning equipment, mixing or loading

For exposures in enclosed areas:

- * A respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)

For exposure outdoors:

- * Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Follow manufacturer's instructions for cleaning maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- * Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT feed treated foliage to livestock or graze treated areas.

DO NOT spray or allow drift onto vegetables, ornamentals or other desirable plants.

DO NOT apply when weather conditions favor drift from target area.

DO NOT contaminate water by cleaning of equipment or disposal of wastes.

Daconate®

Read entire label before using this product.

General Information

DACONATE is a postemergence herbicide for use in cotton, turf and non-crop areas.

DACONATE contains a surfactant (wetting agent) in proper ratio for best results. It is not necessary to add additional surfactant to the spray tank. Local conditions and recommendations vary; consult local agricultural experiment station or extension service weed specialists for recommendations in your area.

Mixing Instructions: DACONATE herbicide must be thoroughly dissolved. Fill the spray tank with half the required amount of water and, with pump or agitator operating, add the recommended amount of DACONATE herbicide and continue filling the tank with balance of water.

DO NOT store spray solution in tank overnight. Clean application equipment thoroughly after use by flushing with water in a safe place.

Application Methods: DACONATE should be applied with a low-volume, properly calibrated sprayer having satisfactory pumping and bypass action. **ADJUST NOZZLES TO KEEP SPRAY OFF COTTON FOLIAGE AND IN A MANNER TO ALLOW MAXIMUM COVERAGE OF WEEDS.** Adequate coverage is very important for effective weed control. Apply during warm weather when weeds are in an active stage of growth.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and green-houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- * Coveralls
- * Waterproof gloves
- * Shoes plus socks
- * Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and domestic animals off treated areas until this material has been washed into the soil.

DO NOT apply this product through any type of irrigation equipment.

Cotton: DACONATE is useful as a directed postemergence spray for controlling weeds in cotton such as barnyardgrass, cocklebur, smooth and large crabgrass, dallisgrass, goosegrass, johnsongrass and nutsedge. Apply as a DIRECTED broadcast spray at rate of 40 gallons per acre of a solution containing 4 pints (2 quarts) of DACONATE in 40 gallons of water when weeds are small. For band applications, apply 1 gallon of above diluted spray per acre for each 1 inch band width to be treated of cotton grown on 40 inch row spacing. A second or repeat application, if needed, should be timed about 1 to 3 weeks after the first application. Keep spray off cotton foliage.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. DO NOT APPLY AFTER FIRST BLOOM.

Slight burning and reddish discoloration of cotton foliage may occasionally be seen following recommended treatment; however, cotton plants will develop normally and yield will not be affected.