

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty of fitness or of merchantability, and no agent or seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Storage And Disposal

Do not contaminate water, food or feed. Storage or disposal. Open dumping is prohibited.

Storage: Store in a dry place.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Net Contents: 2 Pounds/0.9 Kilograms



Daconil 2787 Fungicide

Turf Care[®]
75% Wettable Powder

Active Ingredient	75.0%
Inert Ingredients	25.0%
Total	100%

Keep Out of Reach of Children
DANGER

Only for Sale to, Use, and Storage by Lawn-care and Landscaping Personnel or Commercial Applicators

SDS Biotech Corporation
 10000 West 10th Avenue, Suite 100
 Denver, Colorado 80231
 (303) 751-1000

3202-4020-1004-8503

Precautionary Statements
Hazards To Humans And Domestic Animals
Danger

May cause severe eye damage. May be a potent skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling. Do not get in or on face. Do not get in hair. Wash face and hair immediately with plenty of water. Seek medical attention if in eyes, on face, or in hair.

Always start with skin of face. Wash exposed skin thoroughly with plenty of water after handling.

Do not take internally.

First Aid

Note to physician: Persons having an allergic reaction respond to treatment with anti-stimulants, or steroid creams and/or systemic steroids.

Environmental Hazards

This product should be kept away from aquatic life and other sensitive organisms. Runoff from treated areas may be hazardous to aquatic organisms in the receiving water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to treated areas.

Precautionary Statements
Hazards To Humans And Domestic Animals
Danger

Corrosive causes severe eye damage. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.

Do not take internally.

Avoid breathing dust or spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

First Aid:

Note to physician: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine organisms. Runoff from treated areas may be hazardous to organisms in neighboring areas. Do not apply directly to water bodies. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ATTACH INSERT HERE

**SEE ENCLOSED FOLDER FOR
DIRECTIONS FOR USE.**

BEST AVAILABLE COPY

ACCEPTED
JUL 22 1985

U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF PESTICIDE REGULATION
WASHINGTON, D.C. 20250
50334/4



Daconil 2787

Fungicide

Turf Care®

75% Wettable Powder

Active ingredient: Chlorothalonil (tetrachlorosophthalonitrile)*	75.0%
Inert ingredients:	25.0%
Total	100.0%

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

A Broad-Spectrum Fungicide For Use on Golf Course Tees, Greens and Fairways, Ornamental Turfgrass and Ornamentals.

Read entire label carefully and use only as directed.

3302-4020-8007-8503

A-275-XXX-L 1083

General Information

Daconil 2787 Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

Add the required amount of Daconil 2787 to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations.

Do not combine Daconil 2787 in the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown that combination physically compatible, effective and noninjurious under your conditions of use.

Precautionary Statements

Hazards to Humans and Domestic Animals

Danger

Corrosive, causes severe eye damage. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.

Do not take internally.

Avoid breathing dust or spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

First Aid:

Note to physician: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

Directions For Use

Only for sale to, use, and storage by lawncare and landscaping personnel or commercial applicators.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

Turf: Do not mow or water after treatment until spray deposit on turfgrass is thoroughly dry. Daconil 2787 should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply Daconil 2787 Fungicide in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease condition, use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

Disease	Application Interval	Application Rate Per Acre
Sclerotinia dollar spot	7 - 10 Days 14 - 21 Days	2 3/4 to 5 1/2 lb 5 1/2 to 10 lb
Helminthosporium leafspot	7 - 10 Days 14 - 21 Days	5-1/2 lb 5 1/2 to 10 lb
Rhizoctonia brown patch	7 - 14 Days	5 1/2 to 10 lb
Anthracoze	7 - 14 Days	8 to 16 lbs

Golf Course Tees and Fairways: Apply Daconil 2787 Fungicide at adequate amount of coverage. This amount is 1,000 square feet and less. Under severe conditions, use higher application rates and spray.

Disease	Application Rate Per Acre
Anthracoze	8 to 16 lbs
Copper spot	
Curvularia leaf spot	
Dollar spot	
Gray leaf spot	
Helminthosporium leaf spot and melting out	
Large brown patch	
Red thread	
Stem rust of bluegrass	
DICHOCHNEA Alternaria leaf spot (Callitriche only)	

* Recommended rates use lower rates when higher indicated rates.

** Rates for use on a 7-1/2 inch mowed turfgrass.

BEST AVAILABLE COPY

Ornamentals:

Species	Diseases Controlled	Suggested First Application
BROADLEAF SHRUBS AND TREES		
Ash (Fraxinus)	Cercospora Cercosporium Cylindrosporium leafspots	Spring bud break
Azalea* Rhododendron*	Phytophthora die back Diploma Powder blight	New leaf emergence Early bloom
Buckeye Horsechestnut	Leaf blotch Anthraxnose	Spring bud break
Cherry Laurel	Cercospora leafspot	Petal fall
Crabapple	Scab Cedar apple rust Sphaerotheca leafspot	Spring bud break
Dogwood	Sclerotia leafspot	Early bloom
Eucalyptus	Anthraxnose	Spring bud break
Firethorn	Scab	Spring bud break
Flowering Almond Dance Sand Cherry	Monilia blossom/ branch blight	Early bloom
Hawthorn	Rust Fabrea leafspot	Pre bloom
Holly	Rhizoctonia web blight	Warm, moist conditions
Mountain Laurel	Cercospora leafspot	Spring bud break
Oak (red group only)	Taphrina blister Achnopete leafspot Anthraxnose	Dormant buds well
Oregon Grape (Mahonia)	Rust	Spring bud break
Philirea	Fabrea (Entomosporium) leafspot	Spring bud break
Pieris (Andromeda)	Phytophthora die back	New leaf emergence
Poplar	Marschneria leafspot	Spring bud break
Pine	Cercospora leafspot	Prolonged wet conditions
Sycamore Pine	Anthraxnose	Spring bud break
Viburnum	Powdery mildew	Mid summer

Species	Diseases Controlled	Suggested First Application
BULBS AND FLOWERING PLANTS		
Carnation	Alternaria leafspot/ branch rot Botrytis flower blight	Transplant of cuttings Cool, moist conditions
Chrysanthemum/ Daisy	Mycosphaerella ray blight Sclerotia leafspot Botrytis flower blight (gray mold)	Transplant of cuttings Pre bloom
Geranium	Botrytis blight/rust	Cool, moist conditions
Gladiolus	Carnetaria leaf/ flower spot Botrytis leaf/ flower spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea* (leaves only)	Cercospora and Sclerotia leafspots Rust	Early propagation
Iris	Botrytis blossom blight Dichymella leafspot	Cool, moist conditions
Lily	Botrytis gray mold	Pre bloom
Petunia*	Phytophthora blight (folar phase) Botrytis blight	Pre bloom
Rose (Use 1 lb per 100 gallons)	Black spot Botrytis blight	Spring bud break
Statice	Anthraxnose Cercospora Alternaria Botrytis leaf blights	Spring bud break
Zinnia	Powdery mildew	First sign of disease

*Discoloration of blooms has been noted on certain varieties when applications are made during flowering.

Conifers:

Species	Diseases Controlled	Suggested First Application
FOLIAGE PLANTS		
Dracaena	Fusarium leafspot	Pre transplant
Pachysandra (Use 3 lb per 100 gallons)	Volutella leaf blight	Spring bud break
Leatherleaf fern	Ascochyta blight Cercospora leafspot Cylindrocladium leafspot Rhizorhiza blight	Spring bud break
Parlor palm (Chamaedorea)	Botrytis (Helmintho sporium) leafspot	Cool, moist conditions
Prayer plant (Maranta)	Helmintho sporium leafspot	Early propagation
Oyster plant (Rhoeo)	Tan leafspot	Early propagation
Synonymum	Cephalosporium leafspot	Warm, moist conditions
Philodendron	Phytophthora blight Dactylaria leafspot	Moist conditions

Diseases Controlled	Recent W-75 Rate/Acre	Application Directions
Sclerotia blight Canker (penic), Sclerotia blight, an. etc. etc. Need-leaf (Douglas fir)	1 1/2 to 2 3/4 lb	Make first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
Sirococcus Tip Blight	2 to 3 1/2 lb	
Rhizosphaera Needleaf (spruces) Sclerotia brown spot (pines)	5 1/2 lb	
Lophodermium Needleaf	1 1/2 to 2 3/4 lb	North Central and Northeastern States: Begin applications in mid July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 2 3/4 lb per acre on a 3 week schedule.
Botrytis seedling blight Phoma leaf blight	1 1/2 to 2 3/4 lb	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a dry place.

Postlabel Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

SDS Motech Corporation
Agricultural Chemicals Business
Painesville, Ohio 44077

EPA Reg. No. 50534-4
EPA Est. No. 50534-TX-01

Printed in U.S.A.

AG3-SL8 2/84



BEST AVAILABLE COPY

BEST AVAILABLE COPY