

**Precautionary Statements
Hazards to Humans and Domestic
Animals**

Danger

Corrosive, causes severe eye damage. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.

Do not take internally.

Avoid breathing dust or spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

First Aid

Note to physician: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SEE ENCLOSED FOLDER FOR
DIRECTIONS FOR USE.**

Net Contents: 2 Pounds/0.9 Kilograms



Daconil 2787 Fungicide

**Turf Care[®]
75% Wettable Powder**

Active Ingredient: Chlorothalonil (Tetrachloroisophthalonitrile)*	75.0%
Inert Ingredients:	25.0%
Total:	100.0%

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

**Keep Out of Reach of Children
DANGER**

See side panel for additional precautionary statements

Fermenta Plant Protection Company
5866 Hensley Road, P.O. Box 8000
Mentor, Ohio 44061-9000

EPA Reg. No. 50534-4
EPA Est. No. 35982-TX-1

3202-4020-1004-8705

10F3
pma
50534-4

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a dry place.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty of Fitness or of Merchantability, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

This package is sold by weight, not by volume. Settling of contents may have occurred during shipping and handling.



OCT 26 1987

50534-4



Daconil 2787

Fungicide

Turf Care[®] 75% Wettable Powder

Active ingredient: Chlorothalonil (tetrachlorosophthalonitrile)*	75.0%
Inert ingredients:	25.0%
Total	100.0%

*Covered under U.S. Patent No. 3,290,353
and No. 3,331,735

A Broad-Spectrum Fungicide For Use on Golf Course Tees, Greens and Fairways, Ornamental Turfgrass and Ornamentals.

Read entire label carefully and use only
as directed.

3302-4020-8007-8705

General Information

Daconil 2787 Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

Add the required amount of Daconil 2787 to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations.

Do not combine Daconil 2787 in the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use.

Precautionary Statements

Hazards to Humans and Domestic Animals

Danger

Corrosive, causes severe eye damage. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.

Do not take internally.

Avoid breathing dust or spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

First Aid:

Note to physician: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

Use: Do not mow or water after treatment until spray deposit on turfgrass is thoroughly dry; Daconil 2787 should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply Daconil 2787 Fungicide in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

Disease	Application Interval	Application Rate Per Acre
Sclerotinia dollar spot	7 - 10 Days 14 - 21 Days	2-3/4 to 5-1/2 lb 5-1/2 to 10 lb
Helmintosporium leafspot	7 - 10 Days 14 - 21 Days	5-1/2 lb 5-1/2 to 10 lb
Rhizoctonia brown patch	7 - 14 Days	5-1/2 to 10 lb
Anthraxnose	7 - 14 Days	8 to 16 lbs

Golf Course Tees and Greens and Ornamental Turfgrass: Apply Daconil 2787 Fungicide in an adequate amount of water to provide complete coverage. This amount may vary from 2 to 10 gallons per 1,000 square feet. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

Disease	Interval of Application	RATE Gallons Per 1,000 sq. ft.	
		Preventive*	Curative**
Anthraxnose	7-14 Days	3-6	—
Copper spot	7-10	4-6	6-8
Curculion leaf spot	7-10	2-4	4-8
Dollar spot	7-14	2-4	4-8
Gray leaf spot	7-10	2-4	4-8
Helmintosporium leaf spot and melting out	7-10	2-4	4-8
Large brown patch	7-10	2-4	4-8
Red thread	7-10	2-6	6-8
Stem rot of bluegrass	7-14	4-6	6-8
DICHONDRA Alternaria leaf spot (California only)	7-14	4-6	6-8

*Recommended rates for preventing disease establishment, use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe.

**Rates for use on a 7-day schedule when disease is present. Higher indicated rate should be applied under severe conditions.

Turfgrasses: Gray Snow Mold caused by *Typhula* spp. - Apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1,000 square feet). Apply 5-1/2 to 11 ounces of Daconil 2787 Fungicide per 1,000 square feet of turf area. Application must be made before snow cover in autumn. Use the higher rate if turf layer remains frozen prior to snow cover. If snow cover is intermittent or lacking during the winter, re-apply Daconil 2787 at 5-1/2 ounces per 1,000 square feet at monthly intervals until gray snow mold conditions no longer prevail. In areas where pink snow mold (Geriacha or Fusarium patch) is likely to occur, apply Daconil 2787 at 5-1/2 ounces per 1,000 square feet in combination with either Tercan[®] 1991 50WP at 2 ounces per 1,000 square feet or Chipco[®] 26019 50WP at 4 ounces per 1,000 square feet of turf area.

Fusarium (Geriacha) Patch For control of Fusarium patch only in areas where snow cover is intermittent or lacking during the winter, apply 5-1/2 to 10 ounces of Daconil 2787 Fungicide per 1,000 sq. ft. of turf area. Begin applications in late autumn and re-apply at 21 to 28 day intervals until conditions favorable for Fusarium patch no longer prevail.

*Tercan is a registered trademark of E I DuPont de Nemours & Company, Inc.

**Chipco is a registered trademark of Filtow Products, Inc.

Ornamentals and Conifers: Apply Daconil 2787 Fungicide at rate of 1-1/2 pounds per 100 gallons of water unless other directions are given in the tables below. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, use the higher rate specified and the shortest indicated interval between applications.

Daconil 2787 may be used in greenhouses. Applicators and attending personnel should wear protective clothing including long sleeves, gloves, goggles or eye shield plus a face-fitting respirator specifically designed to remove organic pesticide vapors and particles. Do not use mistblowers or high pressure spray equipment when making applications of Daconil 2787 in greenhouses.

Use of Daconil 2787 is recommended for disease control on the ornamentals and conifers listed in the tables below. Use of Daconil 2787 is not recommended for species that are not listed.

Ornamentals:

Species	Diseases Controlled	Suggested First Application
BROADLEAF SHRUBS AND TREES:		
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporum leafspots	Spring bud break
Azalea,* Rhododendron*	Phytophthora die-back, Ovary/ovule blight	New leaf emergence, Early bloom
Buckeye, Horsechestnut	Leaf blotch, Anthracnose	Spring bud break
Cherry - Laurel	Cercospora leafspot	Petal fall
Crabapple	Scab, Cedar-apple rust, Sphaeropegia leafspot	Spring bud break
Dogwood	Septoria leafspot	Early bloom
Eucalyptus	Anthracnose	Spring bud break
Fraxinus	Scab	Spring bud break
Flowering Almond, Quince, Sand Cherry	Mamma blossom/branch blight	Early bloom
Hazelnut	Rust, Fabraea leafspot	Pre-bloom
Holly	Rhizoctonia web blight	Warm, moist conditions
Mountain Laurel	Cercospora leafspot	Spring bud break
Oak (red group only)	Taphrina blister, Actinopeltis leafspot, Anthracnose	Dormant budswell
Oregon - Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea (Entomosporium) leafspot	Spring bud break
Pieris (Andromeda)	Phytophthora die-back	New leaf emergence
Poplar	Marsipora leafspot	Spring bud break
Privet	Cercospora leafspot	Prolonged wet conditions
Sycamore, Plane tree	Anthracnose	Spring bud break
Viburnum	Powdery mildew	Mid-summer

Species	Diseases Controlled	Suggested First Application
BULBS AND FLOWERING PLANTS:		
Camellia	Alternaria leafspot, bract rot, Botrytis flower blight	Transplant of cuttings, Cool, moist conditions
Chrysanthemum/ Daisy	Mycosphaerella ray blight, Septoria leafspot, Botrytis flower blight (gray mold)	Transplant of cuttings, Pre-bloom
Geranium	Botrytis blight, rust	Cool, moist conditions
Gladiolus	Curvatura leaf/flower spot, Botrytis leaf/flower spot	Early propagation
Hollyhock	Rust	Early seedling stage
Hydrangea* (foliage only)	Cercospora and Septoria leafspots, Rust	Early propagation
Iris	Botrytis blossom blight, Doidymella leafspot	Cool, moist conditions
Lily	Botrytis gray mold	Pre-bloom
Peonies*	Phytophthora blight (later phase), Botrytis blight	Pre-bloom
Rose (Use 1 lb per 100 gallons)	Black spot, Botrytis blight	Spring bud break
Stachys	Anthracnose, Cercospora, Alternaria, Botrytis leaf blights	Spring bud break
Zinnia	Powdery mildew	First sign of disease

* Discoloration of blooms has been noted on certain varieties when applications are made during blooming

Species	Diseases Controlled	Suggested First Application
FOLIAGE PLANTS:		
Dracaena	Fusarium leafspot	Pre-transplant
Pachysandra (Use 3 lb per 100 gallons)	Volvetella leaf blight	Spring bud break
Leatherleaf fern	Ascochyta blight, Cercospora leafspot, Cylindrocladium leafspot, Rhizoctonia blight	Spring bud break
Parlor palm (Chamaedorea)	Bipolaris (Helminthosporium) leafspot	Cool, moist conditions
Prayer plant (Maranta)	Helminthosporium leafspot	Early propagation
Hyacinth plant (Ritona)	Tan leafspot	Early propagation
Synonym	Cephalosporium leafspot	Warm, moist conditions
Philodendron	Phytophthora blight, Lactyaria leafspot	Moist conditions

Conifers:

Diseases Controlled	Amount W-75 Rate/Acre	Application Directions
Sclerotinia Canker (pines), Sema Needlecast and Rhizoctonia Needlecast (Douglas-fir)	1-1/2 to 2-3/4 lb	Make first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
Sarcococcus Tip Blight	2 to 3-1/2 lb	
Rhizosphaera Needlecast (spruces), Scirrhia brown spot (pines)	5-1/2 lb	
Lepidodermium Needlecast	1-1/2 to 2-3/4 lb	North Central and Northeastern States. Begin applications in mid-July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 2-3/4 lb per acre on a 3 week schedule.
Botrytis seedling blight, Phoma twig blight	1-1/2 to 2-3/4 lb	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a dry place.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinse is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

This package is sold by weight, not by volume. Some settling of contents may have occurred during shipping and handling.

Formosa Plant Protection Company
5966 Hershey Road, P.O. Box 8000
Mentor, Ohio 44061-8000

EPA Reg. No. 50534-4
EPA Est. No. 35982-TX-1

Printed in U.S.A.

