

**Proposed Labeling****ISK BIOSCIENCES™****SUPPLEMENTAL DIRECTIONS FOR THE USE OF****BRAVO® 500**

EPA Reg. No. 50534-8

**TO CONTROL DISEASES OF MUSHROOMS****DIRECTIONS FOR USE:**


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**MUSHROOMS:** Verticillium brown spot and dry bubble - Apply 4 to 8 fl. oz. of Bravo® 500 per 1,000 sq. ft. of mushroom bed. Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (8 fl.oz.) of Bravo 500 in the first application and the low rate (4 fl.oz.) of Bravo 500 in the second application. The first application should be made within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning. Do not apply within 5 days of first harvest. Make no more than two applications per cropping cycle. Do not apply more than 12 fl. oz. of Bravo 500 per cropping cycle.

**NOTE:** Follow all applicable directions, restrictions, and precautions on the EPA-registered label.

**NOTE:** This labeling must be in the possession of the user at the time of pesticide application.

3/15/96



ISK Biosciences Corporation  
5966 Heisley Road, P.O. Box 8000, Mentor, Ohio 44061-8000, U.S.A.  
216/357-4100 FAX: 216/354-9506