 <p align="center"><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b> Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE:  <input type="checkbox"/> Registration  <input checked="" type="checkbox"/> Reregistration                  (under FIFRA, as amended)</p>	EPA Reg. Number: 50534-8	Date of Issuance: NOV 10 2005
	Terms of Issuance: Unconditional	
	Name of Pesticide Product: Bravo 500	
Name and Address of Registrant (include ZIP Code): GB Biosciences Corporation 410 Swing Road P.O. Box 18300 Greensboro, NC 27409 Attn.: Ruhi Rezaaiyan		
<p><b>Note:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>Based on your response to the Reregistration Eligibility Document(s), EPA has reregistered the product listed above. Enclosed is a copy of your label stamped "Accepted." This action is taken under the authority of section 4(g)(2)(c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.</p>		
Signature of Approving Official: Tony Kish, Product Manager (22) Fungicide Branch Registration Division (7505C)	Date: NOV 10 2005	

2  
26

Pg. 2 50534-8

A copy of your label stamped Accepted is enclosed for your records.

Sincerely,

*TS/CG*  
*1/2/74*

Tony Kish  
Product Manager (22)  
Fungicide Branch  
Registration Division (7505C)

Enclosure

3  
26

**Bravo® 500**

**Agricultural Fungicide**

Active Ingredient:	
Chlorothalonil (tetrachloroisophthalonitrile) .....	40.4%
Other Ingredients: .....	59.6%
Total:	100.0%

Contains 4.17 Pounds Chlorothalonil Per Gallon (500 grams per liter)

EPA Reg. No. 50534-8

EPA Est. 50534-TX-001

**KEEP OUT OF REACH OF CHILDREN.**


**WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

SCP 50534-8X XXX XXXX

XX gallons  
Net Contents



4  
26

<b>FIRST AID</b>	
<b>If inhaled</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>If swallowed</b>	<ul style="list-style-type: none"><li>• Call poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<p style="text-align: center;"><b>Note to Physician</b></p> <p>Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p style="text-align: center;"><b>HOT LINE NUMBER</b></p> <p style="text-align: center;">For 24 Hour Medical Emergency Assistance for Incidents Involving Human or Animal Exposure Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call <b>1-800-888-8372</b></p>	

5  
26

---

## PRECAUTIONARY STATEMENTS

---

### Hazards to Humans and Domestic Animals

#### WARNING/AVISO

May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not breathe spray mist. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

#### Mixers, Loaders, Applicators and all other handlers must wear:

- long-sleeved shirt and long pants
- chemical resistant gloves made of any waterproof material
- shoes plus socks
- a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N, R, P or HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

6  
26

### User Safety Recommendations

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of GB Biosciences Corporation or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold GB Biosciences and Seller harmless for any claims relating to such factors.

2  
26

GB Biosciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or GB Biosciences, and Buyer and User assume the risk of any such use. **GB BIOSCIENCES MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

In no event shall GB Biosciences or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF GB BIOSCIENCES AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF GB BIOSCIENCES OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

GB Biosciences and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of GB Biosciences.

**A Broad Spectrum Agricultural Fungicide**  
**Read entire label carefully and use only as directed.**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Bravo 500 should be used only in accordance with recommendations on this label or in separately published GB Biosciences supplemental labeling recommendations for this product.

DO NOT apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow workers to enter treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical resistant gloves made of any waterproof material
- shoes plus socks
- protective eyewear

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted-entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

- (1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS-required decontamination site intended for workers entering the treated area.
- (2) Workers must be informed, in a manner they can understand:
  - that residues in the treated area may be highly irritating to their eyes
  - that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes
  - that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water
  - how to operate the eyeflush container



9  
26

---

## **STORAGE AND DISPOSAL**

---

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

### **Pesticide Storage**

Store in a cool place. Protect from excessive heat.

### **Pesticide Disposal**

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### **Container Disposal**

#### **Plastic Containers**

DO NOT reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by alternative methods allowed by state and local authorities.

#### **Returnable Refillable Containers**

If Bravo 500 is packaged in a returnable refillable container, then, after use, do not rinse container. The contents of this container cannot be completely removed by cleaning. Return container intact to point of purchase.

This container must only be refilled with Bravo 500. Refilling with materials other than Bravo 500 will result in contamination and may weaken container. **DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE.** Before refilling, inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn threads on closure devices. Check for leaks after refilling and before transport. Do not refill or transport a damaged or leaking container.

#### **Bulk and Minibulk Containers**

Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

**CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

---

## **GENERAL INFORMATION**

---

Bravo<sup>®</sup> 500 is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases. Bravo 500 is recommended for use in programs which are compatible with the principles of Integrated Pest Management (IPM), which

include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

Bravo 500 is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk of disease resistance exhibit a single-site mode of fungicidal action. Bravo 500, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your federal or state Cooperative Extension Service representatives for guidance on the proper use of Bravo 500 in programs which seek to minimize the occurrence of disease resistance to other fungicides.

Bravo 500 can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

### **GENERAL PRECAUTIONS AND RESTRICTIONS**

DO NOT use on greenhouse-grown crops.

DO NOT combine Bravo 500 with with Dipel<sup>®</sup>, Latron B-1956<sup>®</sup> or Latron AG-98<sup>®</sup> as phytotoxicity may result from the combination when applied to the crops on this label.

This product must not be applied within 150 feet (for aerial and airblast applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

#### **Spray Drift Precautions**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information**.

## **Aerial Drift Reduction Advisory Information**

[This section is advisory in nature and does not supercede the mandatory label requirements.]

### **Information on Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable conditions (See Wind, Temperature).

### **Controlling Droplet Size**

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting the nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

### **Boom Length**

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

### **Application Height**

Applications should not be made at a height greater than 10 ft. above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### **Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

### **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### **Temperature Inversions**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

---

## **APPLICATION**

---

**Note:** Slowly invert container several times to assure uniform mixture.

Dosage rates on this label indicate pints of Bravo 500 per acre, unless otherwise stated. Under conditions favoring disease development the high rate specified and shortest application interval should be used.

The required amount of Bravo 500 should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of Bravo 500 in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

### Field and Row Crops:

Apply Bravo 500 in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See application and calibration instruction below.

### Application and Calibration Techniques for Sprinkler Irrigation - Chemigation

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). DO NOT apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

**Always inject Bravo 500 into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.**

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this

valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. DO NOT apply when wind speed favors drift beyond the area intended for treatment.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Bravo 500 may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

#### **A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment**

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix recommended amount of Bravo 500 for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until Bravo 500 has been cleared from last sprinkler head.

#### **B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment**

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

15  
26

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of Bravo 500 for acreage to be covered with water so that the total mixture of Bravo 500 plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. No agitation should be required. Bravo 500 can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until Bravo 500 has been cleared from last sprinkler head.

**Metric Conversion Guide**

2.125 pints equals 1.0 liter

4.25 pints equals 2.0 liters

16  
26

**Directions for Application**

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Bean (Snap)	Rust	2 to 4¼ pts. (1.0 to 2.25)	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat as necessary (the minimum re-treatment interval is 7 days) to maintain control. DO NOT apply more than 17 pints Bravo 500 (9 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.
	Botrytis blight (gray mold)	4¼ pts. (2.25)	
Beans (Dry) (except soybeans) bean, adzuki bean, broad bean, dry bean, lablab bean, navy bean, kidney bean, lima bean, moth bean, mung bean, pink bean, pinto bean, tepary bean, urd bean, yardlong catjang chickpea (garbanzo) cowpea lupin, grain lupin bean, rice bean, runner bean, jackbean pea, blackeyed pea, southern	Rust Anthracnose Downy mildew Cercospora leaf spot (blackeye only) Ascochyta blight	2 to 2¾ pts. (1.0 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days). For use only on beans to be harvested dry with pods removed. DO NOT apply more than four times per growing season. DO NOT apply more than 11.5 pints Bravo 500 (6 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days before harvest.  Bravo 500 may be applied through sprinkler irrigation equipment. See calibration directions preceding this section.



17  
26

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Cabbage Chinese Cabbage (tight-headed varieties only) Cauliflower Broccoli Chinese Broccoli Brussels Sprouts	Alternaria leaf spot Downy mildew	2¼ pts. (1.125)	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control. DO NOT apply more than 23 pints Bravo 500 (12 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest to Chinese cabbage or Chinese broccoli.
	Ring spot (California only)	2¼ pts. (1.5)	For field-seeded Brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.
Carrot	Cercospora (early) blight Alternaria (late) blight	2¼ to 2¾ pts. (1.125 to 1.5)	Use in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control. DO NOT apply more than 29 pints Bravo 500 (15 lbs. a.i.) per acre during each growing season.  Bravo 500 may be applied through sprinkler irrigation equipment (solid set, portable wheel move or center pivot systems only). See calibration directions preceding this section.
Celery	Cercospora (early) blight Septoria (late) blight Basal stalk rot ( <i>Rhizoctonia solani</i> )	2¼ to 4¼ pts. (1.5 to 2.25)	Use in sufficient water to obtain adequate coverage. Start applications when transplants are set in field and repeat at a 7 day interval as needed to maintain control (the minimum re-treatment interval is 7 days). DO NOT apply more than 34.5 pints Bravo 500 (18 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.
	Pink rot (suppression – 7 day schedule)	4¼ pts. (2.25)	Bravo 500 may be applied through sprinkler irrigation equipment (solid set, portable wheel move or center pivot systems only). See calibration directions preceding this section.
	Early blight Late blight	2¼ to 2¾ pts. (1.125 to 1.5) per 100 gals.	For celery seedbeds, apply in a spray volume of 125 gallons per acre twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.
Corn (Sweet) Corn Grown for Seed	Helminthosporium leaf blights Rust	1 1/8 to 2¼ pts. (0.6 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at 7 day intervals as required to maintain control (the minimum re-treatment interval is 7 days). Under severe disease conditions, use 2¼ to 2¾ pts. Bravo 500 per acre. DO NOT apply more than 17 pints Bravo 500 (9 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT apply to sweet corn to be processed. DO NOT allow livestock to graze in treated fields. DO NOT ensile treated corn or use as livestock forage.

18  
26

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Cranberry	Fruit rots Lophodermium leaf/twig blight	5¼ to 9¼ pts. (3.0 to 5.0)	<p>Apply at early bloom and repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days). Under severe disease conditions use the 9¼ pts./acre rate on a 10 day schedule. DO NOT apply Bravo 500 more than 3 times per season. DO NOT apply more than 29 pints Bravo 500 (15 lbs. a.i.) per acre during each growing season. DO NOT apply within 50 days of harvest. DO NOT apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application.</p> <p>Bravo 500 may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.</p>
<b>Cucurbits</b> Cucumber Cantaloupe Muskmelon Honeydew melon Watermelon Squash Pumpkin	Target spot Anthracnose Downy mildew	2¼ to 2¾ pts. (1.125 to 1.5)	<p>Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum re-treatment interval is 7 days). DO NOT apply more than 30 pints Bravo 500 (15.75 lbs. a.i.) per acre during each growing season.</p> <p>Bravo 500 may be applied through sprinkler irrigation equipment (solid set, portable wheel move or center pivot systems only). See calibration directions preceding this section.</p> <p>NOTE: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply Bravo 500 to watermelons when any of the following conditions are present:</p> <ol style="list-style-type: none"> <li>1. Intense heat and sunlight</li> <li>2. Drought conditions</li> <li>3. Poor vine canopy</li> <li>4. Other crop or environmental conditions which may be conducive to increased natural sunburn</li> </ol> <p>Do not combine Bravo 500 with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions for use.</p>
<b>Grasses</b> Grown for Seed	Stem rust Leaf rust Stripe rust Septoria leaf spot Glume blotch Bipolaris & Drechslera leaf spots	1½ to 2¼ pts. (0.75 to 1.125)	<p>Use in sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Reapply at flag (top) leaf emergence and repeat applications at 14 day intervals (the minimum re-treatment interval is 14 days). DO NOT apply more than 8.5 pints Bravo 500 (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze in treated areas or feed treated plant parts to livestock.</p>
	Selenophoma (eyespot)	1½ to 2¼ pts. (0.75 to 1.5)	

19  
26

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS												
Mint (Indiana, Michigan and Wisconsin only)	Rust Septoria leaf spot	2 pts. (1.0)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground and aircraft applications. Begin applications when emerging plants are 4-8 inches high. Repeat applications at 7 to 10 day intervals to maintain control (the minimum re-treatment interval is 7 days). DO NOT apply more than 3 times per season. DO NOT apply more than 5.75 pints Bravo 500 (3 lbs. a.i.) per acre during each growing season. DO NOT apply within 80 days of harvest. DO NOT feed fresh or extracted mint hay from treated fields to livestock.												
Onion (dry bulb) and Garlic	Botrytis leaf blight/blast Botrytis neck rot (suppression) Purple blotch Downy mildew (Suppression)	1½ to 4¼ pts. (0.75 to 2.25)	<p>Apply in sufficient water to obtain thorough coverage of tops. Bravo 500 is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows:</p> <table border="1" data-bbox="834 827 1398 1152"> <thead> <tr> <th data-bbox="834 827 997 1003"></th> <th data-bbox="997 827 1105 1003">Low Disease Hazard &amp; Prior to Infection</th> <th data-bbox="1105 827 1243 1003">Low Disease Hazard &amp; Some Disease Present</th> <th data-bbox="1243 827 1398 1003">High Disease Hazard</th> </tr> </thead> <tbody> <tr> <td data-bbox="834 1003 997 1079">Rate per Acre</td> <td data-bbox="997 1003 1105 1079">1½ pts</td> <td data-bbox="1105 1003 1243 1079">2 pts</td> <td data-bbox="1243 1003 1398 1079">4¼ pts</td> </tr> <tr> <td data-bbox="834 1079 997 1152">Frequency</td> <td data-bbox="997 1079 1105 1152">10 days</td> <td data-bbox="1105 1079 1243 1152">7 to 10 days</td> <td data-bbox="1243 1079 1398 1152">7 days</td> </tr> </tbody> </table> <p>For suppression of neck rot (<i>Botrytis</i> spp.) during storage, a minimum of three weekly applications prior to lifting, using 2 to 4¼ pts. of Bravo 500 per acre, is recommended.</p> <p>The minimum re-treatment interval is 7 days. DO NOT apply more than 29 pints of Bravo 500 (15 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.</p>		Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard	Rate per Acre	1½ pts	2 pts	4¼ pts	Frequency	10 days	7 to 10 days	7 days
	Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard												
Rate per Acre	1½ pts	2 pts	4¼ pts												
Frequency	10 days	7 to 10 days	7 days												
Onion (green bunching) Leek Shallot Onion Grown for Seed	Botrytis leaf blight/blast Purple blotch Downy mildew (suppression)	2¼ to 4¼ pts. (1.125 to 2.25)	<p>Use in sufficient water to obtain thorough coverage to tops. Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals as long as conditions favor disease (the minimum re-treatment interval is 7 days). Use the high rate and a 7 day schedule of applications when heavy dew or rain persist.</p> <p>DO NOT apply more than 13 pints of Bravo 500 (6.75 lbs. a.i.) per acre during each growing season. DO NOT apply more than 3 times per season or within 14 days of harvest on green bunching onions, leeks or shallots. If additional disease control is needed before harvest, use another registered fungicide.</p>												

20  
26

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Papaya	Alternaria fruit spot Anthracnose Stem end rot	2¼ to 4¼ pts. (1.125 to 2.25)	Apply with ground equipment only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatment at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days).  DO NOT apply more than 13 pints of Bravo 500 (6.75 lbs. a.i.) per acre during each growing season.
Parsnip	Alternaria leaf spot Downy mildew Anthracnose Botrytis blight (gray mold) Bottom rot ( <i>Rhizoctonia</i> )	2¼ to 2¾ pts. (1.125 to 1.5)	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule (the minimum re-treatment interval is 7 days).  DO NOT apply more than 4 times per season or within 10 days of harvest. DO NOT apply more than 11.5 pints of Bravo 500 (6 lbs. a.i.) per acre during each growing season.
Passion Fruit (Hawaii only)	Alternaria fruit and leaf spot (passion fruit brown spot)	2¾ pts. (1.5)	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when fruit spots appear (April to July) and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days).  DO NOT apply more than 14.5 pints of Bravo 500 (7.5 lbs. a.i.) per acre during each growing season.
Peanut	Cercospora (early) leafspot Cercosporidium (late) leafspot	1½ to 2¼ pts. (0.75 to 1.125)	Apply in sufficient water for adequate coverage when leaf wetness first occurs or 30 to 40 days after planting; repeat at 14 day intervals (the minimum re-treatment interval is 14 days). When conditions favor late leafspot or when rust or web blotch occur, apply 2¼ pints Bravo 500 per acre at 14 day intervals for the remainder of the season. DO NOT apply more than 17 pints Bravo 500 (9 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze in treated areas. DO NOT feed hay or threshings from treated fields to livestock.  Bravo 500 may be applied through sprinkler irrigation equipment. Use 2¼ pints Bravo 500 per acre in solid set, portable wheel move, center pivot or traveling gun sprinkler irrigation equipment. See calibration directions preceding this section.
	Rust Web blotch	2¼ pts. (1.125)	

21  
26

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Potato	Early blight Late blight Botrytis vine rot	1 1/8 pt. (0.6) — then — 1½ to 2¼ pts. (0.75 to 1.125)	<p>Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).</p> <p>Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:</p> <ul style="list-style-type: none"> <li>• Vines close between rows</li> <li>• Late blight forecasting measures 18 disease severity values (DSV)</li> <li>• The crop reaches 300 P-days</li> </ul> <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. DO NOT apply more than 21.5 pints of Bravo 500 (11.25 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.</p> <p>Bravo 500 may be applied through sprinkler irrigation equipment (solid set, portable wheel move, center pivot or motorized lateral move systems only). DO NOT exceed a 10 day interval between applications when using this technique. Follow calibration and application directions preceding this section.</p>
Soybean Determinante (Southern) Varieties	Anthracnose Diaporthe pod and stem blight Frogeye leaf spot ( <i>Cercospora sojina</i> ) Purple seed stain Cercospora leaf Blight ( <i>Cercospora kikuchii</i> ) Septoria brown spot		Apply in sufficient water to obtain complete coverage, using at least five gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. Bravo 500 may be applied through sprinkler irrigation equipment. Follow application and calibration directions preceding this section. The minimum re-treatment interval is 14 days. DO NOT exceed total of 3 applications per season. DO NOT apply more than 8.5 pints of Bravo 500 (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 6 weeks of harvest. DO NOT feed treated parts to livestock or allow grazing in treated fields.
		2¼ to 3¼ pts. (1.125 to 1.7)	<b>Two application program</b> — Make the first application at early pod set (R3 stage, when majority of pods are 1/8 to 3/8 inch in length) and the second at beginning seed formation (R5) which occurs about 14 days later.
		1½ to 2¼ pts. (0.75 to 1.5)	<b>Three application program</b> — Make the first application at the beginning of flowering (R1), the second at early pod set (R3) and the third at beginning of seed formation (R5).
	Stem canker ( <i>Diaporthe phaseolorum</i> var. <i>caulivora</i> )	1½ pts. (0.75)	Apply in 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make the first application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease make a second and a third application. Make all applications at 14 day intervals.

22  
—  
26

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Soybean Indeterminate (Northern) Varieties	Anthracnose Diaporthe pod and stem blight Frogeye leaf spot ( <i>Cercospora sojina</i> ) Purple seed stain <i>Cercospora</i> leaf blight ( <i>Cercospora kikuchii</i> ) Septoria brown spot		Apply in sufficient water to obtain complete coverage, using at least five gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. Bravo 500 may be applied through sprinkler irrigation equipment. Follow application and calibration directions preceding this section. The minimum re-treatment interval is 14 days. DO NOT exceed total of 3 applications per season. DO NOT apply more than 8.5 pints of Bravo 500 (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 6 weeks of harvest. DO NOT feed soybean hay or threshings from treated fields to livestock.
		2¼ to 3¼ pts. (1.125 to 1.7)	<b>Two application program</b> — Make the first application when largest pods are 1 to 1-1/2 inches in length and make the second application 14 days later.
		1½ to 2¾ pts. (0.75 to 1.5)	<b>Three application program</b> — Make the first application one week after first flowering and continue applications at 14 day intervals.
Tomato	FOLIAGE (apply every 7-10 days) Early blight Late blight Gray leaf spot Gray leaf mold Septoria leaf spot Target spot	2 to 2¾ pts. (1.0 to 1.5)	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Use the highest rate and shortest interval specified when disease conditions are severe. The minimum re-treatment interval is 7 days).  DO NOT apply more than 28.5 pints Bravo 500 (15 lbs. a.i.) per acre during each growing season.
	FRUIT (apply every 7-14 days beginning at fruit set) Anthracnose Alternaria fruit rot (black mold) Rhizoctonia fruit rot Botrytis gray mold Late blight fruit rot	2¾ to 4 pts. (1.5 to 2.1)	Bravo 500 may be combined in the spray tank with EPA-registered pesticide products that claim copper as the active ingredient and are labeled for control of bacterial diseases on tomatoes. Check the copper manufacturer's label for specific instructions, precautions and limitations prior to mixing with Bravo 500. DO NOT use with Copper Count® N in concentrated spray suspensions.  Bravo 500 may be applied through sprinkler irrigation equipment (solid set or portable wheel move systems only). See calibration directions preceding this section.

### Tree and Orchard Crops

Apply Bravo 500 in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, Bravo 500 may be applied with aircraft using at least 20 gallons per acre. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of Bravo 500 listed may be used. DO NOT allow livestock to graze in treated areas. The following spray volumes are recommended as gallons of spray per acre:

23  
26

CROP	SPRAY VOLUME (Gallons Per Acre)
Peach, Nectarine, Apricot, Tart Cherry, Plum, Prune	20 (concentrate) to 300 (full dilute)
Sweet Cherry	20 (concentrate) to 300 (full dilute)
Conifers Forest stands Christmas trees Nursery beds	Dilute Concentrate Not used      10 to 20 (aircraft) 100            10 to 50 (aircraft or ground equipment) 100            5 to 10 (ground equipment only)

24  
26

CROP	DISEASES	BRAVO 500 RATE PER		APPLICATION DIRECTIONS
		ACRE	100 GALS.*	
Peach Nectarine Apricot Cherry Plum Prune	Leaf curl Coryneum blight (shothole)	4½ to 6 pts. (2.3 to 3.1)	1½ to 2 pts. (0.75 to 1.0)	For best control of both diseases apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of Bravo 500 for control of leaf curl may be made at any time prior to budswell the following spring. Where Coryneum blight (shothole) occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections.
	Lacy (russet) scab (plum/prune)	4½ to 6 pts. (2.3 to 3.1)	1½ to 2 pts. (0.75 to 1.0)	Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry leaf spot Peach, Nectarine, Apricot scab	4½ to 6 pts. (2.3 to 3.1)	1½ to 2 pts. (0.75 to 1.0)	In addition to the bloom application listed above, make one application at shuck split. DO NOT apply Bravo 500 after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide.  For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10 to 14 days later.
DO NOT apply more than 29.5 pints of Bravo 500 (15.5 lbs. a.i.) per acre during each growing season. The minimum re-treatment interval is 10 days.				



25  
26

CROP	DISEASES	BRAVO 500 RATE PER		APPLICATION DIRECTIONS
		ACRE	100 GALS.*	
Conifers (pines, spruces)	Swiss needlecast	4 to 8 pts. (2.1 to 4.125)	4 to 8 pts. (2.1 to 4.125)	<b>Single application technique:</b> In Christmas tree plantations or forest stands, make one application in the spring when new shoot growth is ½ to 2 inches in length.
	Scleroderris canker (pines) Swiss needlecast	2¼ to 4 pts. (1.125 to 2.1)	2¼ to 4 pts. (1.125 to 2.1)	Make the first application in spring when new shoot growth is ½ to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3-week schedule.
	Sirococcus tip blight	2¾ to 5 pts. (1.5 to 2.6)	2¾ to 5 pts. (1.5 to 2.6)	
	Rhizosphaera needlecast (spruces) Scirrhia brown spot (pines)	8 pts. (4.125)	8 pts. (4.125)	
	Cylaneusma and Lophodermium needlecasts (pines)	4 to 8 pts. (2.1 to 4.125)	4 to 8 pts. (2.1 to 4.125)	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
	Rhabdocline needlecast (Douglas fir)	2¼ to 4 pts. (1.125 to 2.1)	2¼ to 4 pts. (1.125 to 2.1)	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3-week schedule.
	Botrytis seedling blight Phoma twig blight	2¼ to 4 pts. (1.125 to 2.1)	2¼ to 4 pts. (1.125 to 2.1)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
	DO NOT apply more than 31.5 pints of Bravo 500 (16.5 lbs. a.i.) per acre during each growing season. The minimum re-treatment interval for established trees is 21 days. The minimum re-treatment interval in nursery beds is 7 days.			

\*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

26  
26

**MUSHROOMS:** Verticillium brown spot and dry bubble - Apply 4 to 8 fl. oz. of Bravo 500 per 1,000 sq. ft. of mushroom bed. Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (8 fl. oz.) of Bravo 500 in the first application and the low rate (4 fl. oz.) of Bravo 500 in the second application. The first application should be made within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning. DO NOT apply within five days of first harvest. Make no more than two applications per cropping cycle. DO NOT apply more than 12 fl. oz. of Bravo 500 per cropping cycle.

---

Bravo® and the Syngenta logo are trademarks of a Syngenta Group Company

Copper-Count® is a registered trademark of Mineral Research and Development Corporation

Dipel® is a registered trademark of Valent BioSciences Corporation.

Latron B-1956® and Latron AG-98® are trademarks of Dow AgroSciences LLC.

© 2004 Syngenta

For non-emergency (e.g., current product information) call  
Syngenta Crop Protection at 1-800-334-9481.

GB Biosciences Corporation  
Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

SCP 50534-8X XXX XXXX

050534-00008.20050923.Bravo 500.8 Mo RED Resp.doc  
050534-00008.20050923.Bravo 500.8 Mo RED Resp.pdf