

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning

Causes eye irritation. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using.

Do not take internally.

Avoid breathing spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild to moderate irritation and redness or rash on exposed skin areas. Persons having the type of reaction should contact a physician.

First Aid:

Note to Physician: Persons having an allergic reaction to sprays should be treated with antihistamines or oral steroids and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.**

Apply only to areas specified on label.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. **Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty.** In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

REGISTERED

JUL 2 1985

Registered with EPA Reg. No. 40534-9

Net Contents 2 Gallons/7.87 Liters



Daconil 2787[®]

Flowable Fungicide Turf Care[™]

Turf and Ornamental Fungicide

Contains 4.17 Pounds Chlorothalonil Per Gallon (500 grams per liter)

Active Ingredient: Chlorothalonil (methylchloro-sophthalonitrile) 40.4%

Inert Ingredients 59.6%

Total 100.0%

40.4%

59.6%

100.0%

Patented in U.S. Patent No. 3,770,000
No. 1,111,111, 1,222,222, 1,333,333

Keep Out of Reach of Children

WARNING

See side panel for additional precautionary statements.

SDS Biotech Corporation

Agricultural Chemicals Business

Parisville, Ohio 44077

EPA Reg. No. 40534-9

EPA Est. No. 40534-TX-01

3316 4007 9004 8503

3316 4007 9004 8503

Registered with EPA Reg. No. 40534-9

2 CF4

Daconil 2787™ Flowable Fungicide

A Broad-Spectrum Fungicide

For Use on Golf Course Tees, Greens and Fairways, Ornamental Turfgrass and Ornamentals.

Read entire label carefully and use only as directed.

General Information

Daconil 2787 Flowable Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

NOTE: Slowly invert container several times to assure uniform mixture.

Add the required amount of Daconil 2787 to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations.

Do not combine Daconil 2787 to the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

See attached folder for directions for use.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a cool place. Protect from excessive heat.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SEE ENCLOSED FOLDER FOR DIRECTIONS FOR USE.

PLACE INSERT POUCH HERE

U.S. Patent No. 3,298,353,
and No. 3,948,628

1-800-447-8503



Daconil 2787

Flowable Fungicide

Turf Care

Contains 4.17 Pounds Chlorothalonil Per Gallon (500 grams per liter)

Active Ingredient: Chlorothalonil (tetrachlorosophthalonitrile)*	40.4%
Inert Ingredients:	59.6%
Total:	100.0%

*Covered under U.S. Patent No. 3,290,353 No. 3,948,636 and No. 3,331,735

A Broad-Spectrum Fungicide For Use on Golf Course Tees, Greens and Fairways, Ornamental Turfgrass and Ornamentals.

Read entire label carefully and use only as directed.

3316-4007-8001 8503

General Information

Daconil 2787 Flowable Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

Note: Slowly invert container several times to assure uniform mixture.

Add the required amount of Daconil 2787 to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations.

Do not combine Daconil 2787 in the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use.

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning

Causes eye irritation. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and water after handling or using.

Do not take internally.

Avoid breathing spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

First Aid:

Note to Physician: Persons having allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

Turf: Do not mow or wait after treatment until spray deposit on turfgrass is thoroughly dry. Daconil 2787 should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply Daconil 2787 Flowable Fungicide in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

Disease	Application Interval	Application Rate Per Acre
Sclerotinia dollar spot	7-10 Days	4 to 8 pts
	14-21 Days	8 to 14 pts
Helminthosporium leafspot	7-10 Days	8 pts
	14-21 Days	8 to 14 pts
Rhizoctonia brown patch	7-14 Days	8 to 10 pts
Anthraco-nose	7-14 Days	8 to 12 pts

Golf Course Tees and Greens and Ornamental Turfgrass

Apply Daconil 2787 in an adequate amount of water to provide complete coverage. This amount may vary from 2 to 10 gallons (approximately 8 to 38 liters) per 1,000 square feet. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

Disease	Interval of Application	RATE Liquid Ounces* Per 1,000 sq ft	
		Preventive**	Curative***
Anthraco-nose	7-14 Days	5.9	
Copper spot	7-10 Days	6.9	9.11
Curvularia leaf spot	7-10	3.6	6.11
Dollar spot	7-14	3.6	6.11
Gray leaf spot	7-10	3.6	6.11
Helminthosporium leaf spot and melting out	7-10	3.6	6.11
Large brown patch	7-10	3.6	6.11
Red thread	7-10	3.9	9.11
Stem rust of bluegrass	7-14	6.9	9.11
UNIDENTIFIED Alternaria leaf spot (California only)	7-14	6.9	9.11

* One liquid ounce is approximately equal to 29 milliliters.

** Recommended rates for preventing disease establishment. Use lower rates when disease conditions are light to moderate; higher indicated rates when conditions are severe.

*** Rates for use on a 7 day schedule when disease is present. Higher indic. rate should be applied under severe conditions.

Turfgrasses: Gray Snow Mold caused by *Typhula* spp. - Apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1,000 square feet). Apply 8 to 16 fluid ounces of Daconil 2787 Flowable Fungicide per 1,000 square feet of turf area. Application must be made before snow cover in autumn. Use the higher rate if turf layer remains frozen prior to snow cover. If snow cover is intermittent or lacking during the winter, re-apply Daconil 2787 at 8 fluid ounces per 1,000 square feet at monthly intervals until gray snow mold conditions no longer prevail. In areas where pink snow mold (*Gaelectria* or *Fusarium* patch) is likely to occur, apply Daconil 2787 at 8 fluid ounces per 1,000 square feet in combination with either Tersan® 1991 50WP at 2 ounces per 1,000 square feet or Chippo® 26019 50WP at 4 ounces per 1,000 square feet of turf area. *Fusarium* (*Gaelectria*) Patch: For control of *Fusarium* patch only in areas where snow cover is intermittent or lacking during the winter, apply 8 to 14 fluid ounces of Daconil 2787 Flowable Fungicide per 1,000 sq. ft. of turf area. Begin applications in late autumn and reapply at 21 to 28 day intervals until conditions favorable for *Fusarium* patch no longer prevail.

Ornamentals and Coolers: Apply Daconil 2787 Flowable Fungicide at rate of 2 pints per 100 gallons of water unless other directions are given in the tables below. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, use the higher rate specified and the shortest indicated interval between applications.

Daconil 2787 may be used in greenhouses. Applicators and attending personnel should wear protective clothing including long sleeves, gloves, goggles or eye shield plus a face fitting respirator specifically designed to remove organic pesticide vapors and particulates. Do not use mistblowers or high pressure spray equipment when making applications of Daconil 2787 in greenhouses.

Use of Daconil 2787 is recommended for disease control on the ornamentals and coolers listed in the tables below. Use of Daconil 2787 is not recommended for species that are not listed.

*Tersan is a registered trademark of E. I. DuPont de Nemours & Company, Inc.

**Chippo is a registered trademark of Pheno Poulenc, Inc.

2004

Ornamentals:

Species	Diseases Controlled	Suggested First Application
SHRUBBLEAF SHRUBS AND TREES:		
Ash (Fraxinus)	Cercospora, Cymodactylum leafspots	Spring bud break
Azalea "Rhododendron"	Phytophthora die-back, Oudemans flower blight	New leaf emergence, Early bloom
Baccharis, Neracanthus	Leaf blight, Anthracnose	Spring bud break
Cherry - Laurel	Cercospora leafspot	Petal fall
Crapeole	Scab, Cedar-apple rust, Sphaeria leafspot	Spring bud break
Dogwood	Sphaeria leafspot	Early bloom
Eubotrys	Anthracnose	Spring bud break
Fuchsia	Scab	Spring bud break
Flowering Almond, Quince, Sand Cherry	Monilia blossom branch blight	Early bloom
Hawthorn	Rust, Fabraea leafspot	Pre bloom
Holly	Rhizoctonia web blight	Warm, moist conditions
Mountain Laurel	Cercospora leafspot	Spring bud break
Oak (red group only)	Ischyrus blister, Acinospora leafspot, Anthracnose	Dormant leafswell
Oregon - Grape (Mahonia)	Rust	Spring bud break
Prunus	Fabraea (Ectomycorrhizae) leafspot	Spring bud break
Pieris (Andromeda)	Phytophthora die-back	New leaf emergence
Poplar	Marsipora leafspot	Spring bud break
Privet	Cercospora leafspot	Prolonged wet conditions
Sycamore, Plane tree	Anthracnose	Spring bud break
Viburnum	Powdery mildew	Mid summer

Species	Diseases Controlled	Suggested First Application
BULBS AND FLOWERING PLANTS:		
Caration	Alternaria leafspot, branch rot, Botrytis flower blight	Transplant of cuttings, Cool, moist conditions
Chrysanthemum/Daisy	Mycosphaerella ray blight, Septoria leafspot, Botrytis flower blight (gray mold)	Transplant of cuttings, Pre-bloom
Geranium	Botrytis blight, rust	Cool, moist conditions
Glaucolus	Carotolium leaf-spot, Botrytis leaf-spot, flower spot	Early propagation
Hollyhock	Rust	Early seeding stage
Hydrangea (foliage only)	Leptomyces and Septoria leafspots, Rust	Early propagation
Iris	Botrytis blight, Botrytis blight, Botrytis blight	Cool, moist conditions
Lily	Botrytis gray mold	Pre bloom
Petunia*	Phytophthora blight (leaf phase), Botrytis blight	Pre-bloom
Rose (Use 1 1/2 pt per 100 gallons)	Black spot, Botrytis blight	Spring bud break
Statice	Anthracnose, Cercospora, Alternaria, Botrytis leaf blights	Spring bud break
Zinnia	Powdery mildew	First sign of disease

* Discoloration of blooms has been noted on certain varieties when applications are made during flowering.

Species	Diseases Controlled	Suggested First Application
FOLIAGE PLANTS:		
Dracaena	Fusicladium leafspot	Pre transplant
Pachysandra (Use 4 pt per 100 gallons)	Verticillium leaf blight	Spring bud break
Leatherleaf fern	Ascochyta blight, Cercospora leafspot, Cylindrocapsa leafspot, Rhizoctonia blight	Spring bud break
Parlor palm (Chamaedorea)	Bipolaris (Helminthosporium) leafspot	Cool, moist conditions
Prayer plant (Maranta)	Helminthosporium leafspot	Early propagation
Rusty plant (Pilea)	Leaf leafspot	Early propagation
Synonymum	Cephalosporium leafspot	Warm, moist conditions
Philodendron	Phytophthora blight, Oospora leafspot	Moist conditions

Conifers:

Diseases Controlled	Resin/Flammable Ratio/Rate	Application Schedule
Sclerotinia Canker (pinus), Swiss Needlecast and Rhododendrum Needlecast (Douglas fir)	2 to 4 pts	Make first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
Sirococcus Tip Blight	3 to 5 pts	
Phytophthora Needlecast (spruces), Scirpus brown spot (pines)	8 pts	
Lophodermium Needlecast	2 to 4 pts	North Central and Northeastern States: Begin applications in mid-July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 4 pts per acre on a 3 week schedule.
Botrytis seedling blight, Phoma twig blight	2 to 4 pts	Begin applications in nursery beds when seedlings are 4 inches tall and warm, cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a cool place. Protect from excessive heat.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

SDS Biotech Corporation
Agricultural Chemicals Business
Painesville, Ohio 44077

EPA Reg. No. 50534-9
EPA Est. No. 50534 TX 01

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning

Causes eye irritation. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin. Avoiding. Wash exposed areas of skin with soap and warm water after handling.

Do not breathe vapors.

Avoid breathing dusts.

Note to User: This product may produce temporary allergic reactions, such as redness, itching, and swelling of the skin. If you have a known sensitivity to any of the ingredients, do not use this product.

First Aid

Note to Physician: In case of eye contact, flush eyes with plenty of water. In case of skin contact, wash with soap and water. In case of inhalation, move to fresh air.

Environmental Hazards

This product is toxic to aquatic organisms. Do not apply to streams, rivers, or other bodies of water. Do not apply to lawns or other areas where it may be washed into water.

Avoid contact with aquatic life.

Warranty and Limitation of Damages

This warranty is limited to the product as sold. It does not cover damage caused by misuse or application of the product. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. This warranty is void where prohibited by law.

BEST AVAILABLE COPY

Handwritten notes and stamps, including "EPA REG. NO. 505-M-9" and "EPA EST. NO. 3316-4007-9004-8503".

Net Contents 2 Gallons/7.87 Liters



Daconil 2787[®]

Flowable Fungicide

Turf Care[™]

Turf and Ornamental Fungicide

Contains 4.17 Pounds of Chlorothalonil Per Gallon (500 grams per liter)

Active Ingredient Chlorothalonil (65% orisophthalate) 40.4%

Inert Ingredients 59.6%

Total 100.0%

EPA Reg. No. 505-M-9 EPA Est. No. 3316-4007-9004-8503

Keep Out of Reach of Children

WARNING

See label and label for other precautionary statements.

SDS Biotech Corporation




Agricultural Chemical Business

Parsonsville, Ohio 44130

EPA Reg. No. 505-M-9

EPA Est. No. 505-M-EX-01

3316 4007 9004 8503



Daconil 2787[®] Flowable Fungicide
A Broad-Spectrum Fungicide
For Use on Golf Course Tees, Greens and
Fairways, Ornamental Turfgrass and
Ornamentals.

Read entire label carefully and use only as directed.

General Information

Daconil 2787 Flowable Fungicide is formulated for use on golf course tees, greens, and fairways; on ornamental plants and ornamental shrubs, shrubs, and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

NOTE: Slowly invert container several times before use to obtain uniform slurry. Add the required amount of Daconil 2787 to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operation.

Do not combine Daconil 2787 in the spray tank with pesticides, surfactants, or fertilizers unless your product label shows the combination physically compatible, effective and nonphytotoxic under your conditions of use.

Directions For Use

It is a violation of Federal Law to use this pesticide in a manner inconsistent with its labeling.

See attached folder for directions for use.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Store in original container. Do not use if container is damaged.

Storage: Store in a cool, dry place. Do not store near food.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Pesticide sprays and other treatments containing pesticides should be applied to the soil. If the pesticide cannot be disposed of by a private citizen, contact your local or state health or environmental agency for disposal instructions. Do not use if container is damaged or leaking.

Container Disposal: Do not reuse or refill this container. Do not dispose of this container in any way that could result in the container being reused or refilled. Do not burn. Do not puncture. Do not reuse for any other purpose. Do not reuse for any other purpose. Do not reuse for any other purpose.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SEE ENCLOSED FOLDER FOR DIRECTIONS FOR USE.

PLACE INSERT POUCH HERE

BEST AVAILABLE COPY



Turf Care

Contains 4.17 Pounds Chlorothalonil Per Gallon (500 grams per liter)
Active Ingredient: Chlorothalonil (tetrachloroisophthalonitrile) 40.4%
Inert Ingredients 59.6%
Total 100.0%

*Covered under U.S. Patent No. 3,240,213 No. 3,948,636 and No. 3,341,735

A Broad-Spectrum Fungicide For Use on Golf Course Tees, Greens and Fairways, Ornamental Turfgrass and Ornamentals.

Read entire label carefully and use only as directed

3116 4007 82X11 8/81

General Information

Daconil 2787 Flowable Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

Note: Slowly invert container several times to assure uniform mixture.
 Add the required amount of Daconil 2787 to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations.

Do not combine Daconil 2787 in the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use.

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning

Causes eye irritation. May be a potential skin sensitizer.
Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and water after handling or using.

Do not take internally.

Avoid breathing spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reactions should contact a physician.

First Aid

Note to Physician: Persons having allergic reactions respond to treatment with antihistamines, corticosteroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **Do not** apply directly to water. **Do not** apply when weather conditions favor drift from treated areas. Apply only to areas specified on label.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reactions should contact a physician.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

Turf: **Do not** mow or water after treatment until spray deposit on turfgrass is thoroughly dry. Daconil 2787 should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply Daconil 2787 Flowable Fungicide in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

Disease	Application Interval	Application Rate Per Acre
Sclerotinia dollar spot	7-10 Days	4 to 6 pts
Helminthosporium leaf spot	14-21 Days	8 to 14 pts
Rhizoctonia brown patch	7-10 Days	8 pts
Anthraxnose	14-21 Days	8 to 14 pts

Golf Course Tees and Greens and Ornamental Turfgrass: Apply Daconil 2787 in an adequate amount of water to provide complete coverage. This amount may vary from 2 to 10 gallons (approximately 8 to 38 liters) per 1,000 square feet. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

Disease	Interval of Application	RATE Liquid Ounces* Per 1,000 sq. ft. Preventive**	Curative***
Anthraxnose	7-14 Days	5-9	
Copper spot	7-10 Days	6-9	9-11
Curculatula leaf spot	7-10	3-6	6-11
Dollar spot	7-14	3-6	6-11
Gray leaf spot	7-10	3-6	6-11
Helminthosporium leaf spot and wetting rot	7-10	3-6	6-11
Large brown patch	7-10	3-6	6-11
Red thread	7-10	3-9	9-11
Stem rust of bluegrass	7-14	6-9	9-11
THELETRICHA Acremonia leaf spot (California only)	7-14	6-9	9-11

*One liquid ounce is approximately equal to 29 milliliters.
 **Recommended rates for preventing disease establishment; use lower rates when disease conditions are light to moderate; higher indicated rates when conditions are severe.
 ***Rates for use on a 7 day schedule when disease is present. Higher indicated rate should be applied under severe conditions.

Tortricidae, Gray Snow Mold caused by Typhula spp. - Apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1,000 square feet). Apply 8 to 16 fluid ounces of Daconil 2787 Flowable Fungicide per 1,000 square feet of turf area. Application must be made before snow cover in autumn. Use the higher rate if turf layer remains frozen prior to snow cover. If snow cover is intermittent or lacking during the winter, re-apply Daconil 2787 at 8 fluid ounces per 1,000 square feet at monthly intervals until gray snow mold conditions no longer prevail. In areas where pink snow mold (Gaelectria or Fusarium patch) is likely to occur, apply Daconil 2787 at 8 fluid ounces per 1,000 square feet in combination with either Fersan® 1991 SOWP at 2 ounces per 1,000 square feet or Chipco® 26019 SOWP at 4 ounces per 1,000 square feet of turf area.
Fusarium (Gaelectria) Patch: For control of Fusarium patch only in areas where snow cover is intermittent or lacking during the winter, apply 8 to 14 fluid ounces of Daconil 2787 Flowable Fungicide per 1,000 sq. ft. of turf area. Begin applications in late autumn and reapply at 21 to 28 day intervals until conditions favorable for Fusarium patch no longer prevail.

Ornamentals and Conifers: Apply Daconil 2787 Flowable Fungicide at rate of 2 pints per 100 gallons of water unless other directions are given in the tables below. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, use the higher rate specified and the shortest indicated interval between applications.

Daconil 2787 may be used in greenhouse applications and attending personnel should wear protective clothing including long sleeves, gloves, goggles or eye shield plus a face fitting respirator specifically designed to remove organic pesticide vapors and particulates. **Do not** use mistblowers or high pressure spray equipment when making applications of Daconil 2787 in greenhouses.

Use of Daconil 2787 is recommended for disease control on the ornamentals and conifers listed in the tables below. Use of Daconil 2787 is not recommended for species that are not listed.

*Fersan is a registered trademark of E. I. DuPont de Nemours & Company, Inc.
 **Chipco is a registered trademark of The Dow Chemical Company.

BEST AVAILABLE COPY

4064

Ornamentals:

Species	Diseases Controlled	Suggested First Application
BROADLEAF SHRUBS AND TREES		
Ash (Fraxinus)	Cercospora Cercosporidium Eylendosporium leafspots	Spring bud break
Azalea Rhododendron	Phytophthora die back Oudemania flower blight	New leaf emergence Early bloom
Buckeye Horsechestnut	Leaf blotch Anthracnose	Spring bud break
Cherry Laurel	Cercospora leafspot	Petal fall
Crabapple	Scab Cedar apple rust Sphaeropsis leafspot	Spring bud break
Dogwood	Septoria leafspot	Early bloom
Eucornus	Anthracnose	Spring bud break
Fernthorn	Scab	Spring bud break
Flowering Almond Dunce Sand Cherry	Wolfsia blossom branch blight	Early bloom
Hawthorn	Rust Fabraea leafspot	Pre bloom
Holly	Rhizoctonia web blight	Warm, moist conditions
Mountain Laurel	Cercospora leafspot	Spring bud break
Oak (red group only)	Lophna bleser Actinopelte leafspot Anthracnose	Dormant budswell
Oregon Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea Echinochloa leafspot	Spring bud break
Pieris (Andromeda)	Phytophthora die back	New leaf emergence
Poplar	Marschneria leafspot	Spring bud break
Privet	Leucospiza leafspot	Pruned and transplants
Spalsham Flowering	Anthracnose	Spring bud break
Viburnum	Fireblight mildew	Mid summer

Species	Diseases Controlled	Suggested First Application
BULBS AND FLOWERING PLANTS		
Carnation	Alternaria leafspot branch rot Botrytis flower blight	Transplant of cuttings. Cool moist conditions
Chrysanthemum Daisy	Mycosphaerella ray blight Septoria leafspot Botrytis flower blight (gray mold)	Transplant of cuttings Early bloom
Geranium	Botrytis blight-rust	Cool, moist conditions
Gladiolus	Convolvulium leaf flower spot Botrytis leaf flower spot	Early propagation
Hollyhock	Rust	Early seeding stage
Hydrangea (large only)	Leucospiza and Septoria leafspots Rust	Early propagation
Lily	Botrytis budswell blight Elatymelina leafspot	Cool, moist conditions
Lily	Botrytis gray mold	Pre bloom
Petunia	Phytophthora blight (trunk phase) Botrytis blight	Pre bloom
Rose (tree type per 100 gallons)	Black spot Botrytis blight	Spring bud break
Statice	Anthracnose Echinochloa Alternaria Botrytis leaf blights	Spring bud break
Zinnia	Fireblight mildew	First sign of disease

Transplanting of bulbs has been noted in certain varieties when applications are made during flowering.

Species	Diseases Controlled	Suggested First Application
FOLIAGE PLANTS		
Dracaena	Fusarium leafspot	Pre transplant
Pachysandra (Use 4 pt per 100 gallons)	Volvetia leaf blight	Spring bud break
Leatherleaf fern	Ascochyta blight Cercospora leafspot Cylindrocium leafspot Rhizoctonia blight	Spring bud break
Parlor palm (Chamaedorea)	Bipolaris (Helmintho- sporium) leafspot	Cool, moist conditions
Prayer plant (Maranta)	Helminthosporium leafspot	Early propagation
Oyster plant (Rhoeo)	Tan leafspot	Early propagation
Synonymum	Cephalosporium leafspot	Warm, moist conditions
Philodendron	Phytophthora blight Dactylaria leafspot	Moist conditions

Conifers:

Diseases Controlled	Resinell Flowable Rate / Rate	Application Directions
Sclerotinia Canker (pines) Swiss Needlecast and Phaeocline Needlecast (Douglas fir)	2 to 4 pts	Make first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
Sirococcus Tip Blight	3 to 5 pts	
Rhizosphaera Needlecast (spruces) Scirrhia brown spot (pines)	8 pts	
Lophodermium Needlecast	2 to 4 pts	North Central and Northeastern States. Begin applications in mid July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 4 pts per acre on a 3 week schedule.
Botrytis seedling blight (Thuja) twig blight	2 to 4 pts	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a cool place. Protect from excessive heat.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinse is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

SDS Biotech Corporation
Agricultural Chemicals Business
Painesville, Ohio 44077

EPA Reg. No. 505-34-9
EPA Est. No. 505-34-1X-01

BEST AVAILABLE COPY