

10F
K1001

Net Contents

REF: 100-100
In Accordance with FR Notice 92-2,
Based on Draft Labeling Dated

5-12-88

SPECIMEN LABEL PA

FERMENTA
PLANT PROTECTION

Daconil 2787

Flowable Fungicide

Turf and Ornamental Fungicide

Contains 4.17 Pounds Chlorothalonil Per Gallon (500 grams per liter)

Active Ingredient: Chlorothalonil (tetrachloroisophthalonitrile)*	40.4%
Inert Ingredients:	59.6%
Total:	100.0%

WATER

Fermenta Plant Protection Company
5066 Haisley Road
P.O. Box 8000
Mentor, Ohio 44061-8000

EPA Est. No. 50534-TX-01

EPA Reg. No. 50534-9

3316-4060-9069-8708

General Information

Daconil 2787 Flowable Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

Note: Slowly invert container several times to assure uniform mixture.

Add the required amount of **Daconil 2787** to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations.

Do not combine Daconil 2787 in the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use.

Precautionary Statements

Hazards to Humans and Domestic Animals

Warning

Causes eye irritation. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or eye shield when handling this product. In case of contact with eyes, flush with plenty of water immediately for 15 minutes. Seek medical attention for eyes immediately.

Avoid contact with skin or clothing. Wash exposed areas of skin with soap and water after handling or using.

Do not take internally.

Avoid breathing spray mist.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

First Aid:

Note to Physician: Persons having allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **Do not apply directly to water. Do not apply** when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

Turf: **Do not mow or water** after treatment until spray deposit on turfgrass is thoroughly dry. **Daconil 2787** should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply **Daconil 2787 Flowable Fungicide** in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

Disease	Application Interval	Application Rate Per Acre
Sclerotinia dollar spot	7 - 10 Days	4 to 8 pts
	14 - 21 Days	8 to 14 pts
Helminthosporium leafspot	7 - 10 Days	8 pts
	14 - 21 Days	8 to 14 pts
Rhizoctonia brown patch	7 - 14 Days	8 to 14 pts
Anthraxnose	7 - 14 Days	6 to 12 qts

Golf Course Tees and Greens and Ornamental Turfgrass:

Apply **Daconil 2787** in an adequate amount of water to provide complete coverage. This amount may vary from 2 to 10 gallons (approximately 8 to 38 liters) per 1,000 square feet. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

Disease	Interval of Application	RATE	
		Liquid Ounces* Per 1,000 sq. ft.	Curative***
Anthraxnose	7-14 Days	5-9	-
Copper spot	7-10 Days	6-9	9-11
Curvularia leaf spot	7-10	3-6	6-11
Dollar spot	7-14	3-6	6-11
Gray leaf spot	7-10	3-6	6-11
Helminthosporium leaf spot and melting out	7-10	3-6	6-11
Large brown patch	7-10	3-6	6-11
Red thread	7-10	5-9	9-11
Stem rust of bluegrass	7-14	6-9	9-11
DICHONDRA Alternaria leaf spot (California only)	7-14	6-9	9-11

* One liquid ounce is approximately equal to 29 milliliters.

** Recommended rates for preventing disease establishment use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe.

*** Rates for use on a 7 day schedule when disease is present. Higher indicated rate should be applied under severe conditions.



Turfgrasses: Gray Snow Mold caused by *Typhula* spp. - Apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1,000 square feet). Apply 8 to 16 fluid ounces of **Daconil 2787 Flowable Fungicide** per 1,000 square feet of turf area. Application must be made before snow cover in autumn. Use the higher rate if turf layer remains frozen prior to snow cover. If snow cover is intermittent or lacking during the winter, re-apply **Daconil 2787** at 8 fluid ounces per 1,000 square feet at monthly intervals until gray snow mold conditions no longer prevail. In areas where pink snow mold (*Gerlachia* or *Fusarium* patch) is likely to occur, apply **Daconil 2787** at 8 fluid ounces per 1,000 square feet in combination with either Tersan® 1991 50WP at 2 ounces per 1,000 square feet or Chipco® 26019 50WP at 4 ounces per 1,000 square feet of turf area.

Fusarium (Gerlachia) Patch: For control of *Fusarium* patch only in areas where snow cover is intermittent or lacking during the winter, apply 8 to 14 fluid ounces of **Daconil 2787 Flowable Fungicide** per 1,000 sq. ft. of turf area. Begin applications in late autumn and reapply at 21 to 28 day intervals until conditions favorable for *Fusarium* patch no longer prevail.

Ornamentals and Conifers: Apply **Daconil 2787 Flowable Fungicide** at rate of 2 pints per 100 gallons of water unless other directions are given in the tables below. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, use the higher rate specified and the shortest indicated interval between applications.

Daconil 2787 may be used in greenhouses. Applicators and attending personnel should wear protective clothing including long sleeves, gloves, goggles or eye shield plus a face-fitting respirator specifically designed to remove organic pesticide vapors and particulates. **Do not** use mistblowers or high pressure spray equipment when making applications of **Daconil 2787** in greenhouses.

Use of **Daconil 2787** is recommended for disease control on the ornamentals and conifers listed in the tables below. Use of **Daconil 2787** is not recommended for species that are not listed.

* Tersan is a registered trademark of E. I. DuPont de Nemours & Company, Inc.

** Chipco is a registered trademark of Phone-Pouenc, Inc.

Ornamentals:

Species	Diseases Controlled	Suggested First Application
BROADLEAF SHRUBS AND TREES:		
Ash (Fraxinus)	Cercospora, Cercosporidium, Cylindrosporium leafspots	Spring bud break
Azalea,* Rhododendron*	Phytophthora die-back, Jvulinia flower blight	New leaf emergence, Early bloom
Buckeye, Horsechestnut	Leaf blotch, Anthracnose	Spring bud break
Cherry - Laurel	Cercospora leafspot	Petal fall
Crabapple	Scab, Cedar-apple rust, Sphaeropsis leafspot	Spring bud break
Dogwood	Septoria leafspot	Early bloom
Euonymus	Anthracnose	Spring bud break
Fuchsia	Scab	Spring bud break
Flowering Almond, Quince, Sand Cherry	Monilia blossom branch blight	Early bloom
Hawthorn	Rust, Fabraea leafspot	Pre-bloom
Holly	Rhizoctonia web blight	Warm, moist conditions
Mountain Laurel	Cercospora leafspot	Spring bud break
Oak (red group only)	Taphrina blister, Actinopeltis leafspot, Anthracnose	Dormant budswell
Oregon - Grape (Mahonia)	Rust	Spring bud break
Photinia	Fabraea (Entomosporium) leafspot	Spring bud break
Pine (Andromeda)	Phytophthora die-back	New leaf emergence
Poplar	Marssonina leafspot	Spring bud break
Privet	Cercospora leafspot	Prolonged wet conditions
Sycamore, Planetree	Anthracnose	Spring bud break
Viburnum	Powdery mildew	Mid-summer


Species	Diseases Controlled	Suggested First Application
BULBS AND FLOWERING PLANTS		
Carnation	Alternaria leafspot, branch rot, Botrytis flower blight	Transplant of cuttings, Cool, moist conditions
Chrysanthemum Daisy	Mycosphaerella ray blight, Septoria leafspot, Botrytis flower blight (gray mold)	Transplant of cuttings, Pre-bloom
Geranium	Botrytis blight, rust	Cool, moist conditions
Gladolus	Curvularia leaf flower spot, Botrytis leaf flower spot	Early propagation
Hollyhock	Rust	Early seeding stage
Hydrangea* (Hokage only)	Cercospora and Septoria leafspots, Rust	Early propagation
Iris	Botrytis blossom blight, Didymella leafspot	Cool, moist conditions
Lily	Botrytis gray mold	Pre-bloom
Petunia*	Phytophthora blight (folar phase), Botrytis blight	Pre-bloom
Rose (Use 1 1/2 pt per 100 gallons)	Black spot, Botrytis blight	Spring bud break
Statice	Anthracnose, Cercospora, Alternaria, Botrytis leaf blights	Spring bud break
Zinnia	Powdery mildew	First sign of disease

* Discoloration of blooms has been noted on certain varieties when applications are made during flowering.

Species	Diseases Controlled	Suggested First Application
FOLIAGE PLANTS.		
Dracaena	Fusarium leafspot	Pre-transplant
Pachysandra (Use 4 pt per 100 gallons)	Volvetia leaf blight	Spring bud break
Leatherleaf fern	Ascochyta blight, Cercospora leafspot, Cylindrocladium leafspot, Rhizoctonia blight	Spring bud break
Parlor palm (Chamaedorea)	Bipolaris (Helminthosporium) leafspot	Cool, moist conditions
Prayer plant (Maranta)	Helminthosporium leafspot	Early propagation
Oyster plant (Rhoeo)	Tan leafspot	Early propagation
Syngonium	Cephalosporium leafspot	Warm, moist conditions
Philodendron	Phytophthora blight, Dactyaria leafspot	Moist conditions

Conifers:

Diseases Controlled	Dose/Flower/Rate/Acre	Application Directions
Scieroderms Canker (pines), Swiss Needlecast and Rhadocline Needlecast (Douglas-fir)	2 to 4 pts	Make first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
Sirococcus Tip Blight	3 to 5 pts	
Rhizosphaera Needlecast (spruces), Scirrhous brown spot (pines)	8 pts	
Lophodermium Needlecast	2 to 4 pts	North Central and Northeastern States. Begin applications in mid-July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 4 pts per acre on a 3 week schedule.
Botrytis seedling blight, Phoma twig blight	2 to 4 pts	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.



Application and Calibration Techniques for Sprinkler Irrigation on Sod Farms, Ornamental Herbs, Shrubs and Trees:

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system. Do not use Dacoonl 2787 Flowable Fungicide through sprinkler irrigation equipment on golf courses.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not apply this product through irrigation systems connected to a public water system. "Public water system" means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls on both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject DACOONL 2787 FLOWABLE FUNGICIDE into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Dacoonl 2787 Flowable Fungicide may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix recommended amount of Dacoonl 2787 Flowable Fungicide for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until Dacoonl 2787 Flowable Fungicide has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides, however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of Dacoonl 2787 Flowable Fungicide for acreage to be covered with water so that the total mixture of Dacoonl 2787 Flowable Fungicide plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. No agitation should be required. Dacoonl 2787 Flowable Fungicide can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until Dacoonl 2787 Flowable Fungicide has been cleared from last sprinkler head.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a dry place.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Fermenta Plant Protection Company
5966 Hershey Road, P.O. Box 8000
Mentor, Ohio 44063-8000
EPA Reg. No. 505B4-9
EPA Est. No. 50534-T-01

Printed in U.S.A.