

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

6/27/89

Registration Support Branch
Registration Division
Environmental Protection Agency
Washington, DC 20460

Registration Support Branch
Registration Division
Environmental Protection Agency
Washington, DC 20460

Registration Support Branch
Registration Division
Environmental Protection Agency
Washington, DC 20460

- 1. Submit production information (gross and net use, reduced) for this product for the fiscal year in which the use on the crop group, brassica leafy vegetable, is conditionally registered, in accordance with FIFRA section 24. The fiscal year begins October 1 and ends September 30. Production information will be submitted to the Agency no later than November 15, following the end of the preceding fiscal year.

This information should be submitted to:

Registration Support Branch
Registration Division (R2100-1)
Environmental Protection Agency
Washington, DC 20460

BEST AVAILABLE COPY

52584: I: CR-45: Ikeda: MP-6: KENCO: 9/12/89: 9/22/89: AS: VO: EK: AS

CONCURRENCES							
MBOL							
RNAME							
E							

- 3. Submit one (1) copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
 Acting Product Manager (23)
 Fungicide-Herbicide Branch
 Registration Division (87505C)

cc: [unclear]

50534-17

DACTHAL^R G-2.5 HERBICIDE
EPA Reg. No. 50534-17

**APPLICATION TO BRASSICA LEAFY VEGETABLES CROP GROUP
FOR CONTROL OF ANNUAL GRASSES AND CERTAIN BROADLEAVED WEEDS**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ANY APPLICATION OF DACTHAL SHOULD BE MADE PRIOR TO WEED SEED GERMINATION

Crops	When to Apply	Rate/Acre	Remarks
Broccoli, brussels sprouts, cauliflower, cabbage and all other <u>Brassica</u> (cole) leafy vegetables in this crop group.*	At seeding or transplanting	320 to 420 lbs DACTHAL G-2.5 (See Rates of Application)	Apply uniformly to the soil. DACTHAL can be applied directly over transplants with no injury. If weeds have emerged, soil should be clean cultivated or weeded prior to application. Can be preplant incorporated.

*Brassica leafy vegetables crop group includes broccoli, Chinese broccoli, broccoli raab (rapini), Brussels sprouts, cabbage, Choy (bok choy, napa), Chinese mustard cabbage (gai choy), cauliflower, collards, kale, kohlrabi, mustard greens and rape greens.

NOTE: It is important to read and observe the precautionary statements and all other information appearing on the DACTHAL G-2.5 HERBICIDE label. Use only as directed.

NOTE: This labeling must be in the possession of the user at the time of pesticide application.

17

SUSCEPTIBLE WEEDS

(Controlled at Lowest Recommended Rates for Each Soil Type)

carpetweed	lovegrass
chickweed, common	nettle, burning
crabgrass, large	pansy, field
crabgrass, smooth	purslane
foxtail, green	pusley, Florida
foxtail, yellow	witchgrass
lambquarters, common	

MODERATELY SUSCEPTIBLE WEEDS

(Highest Recommended Rates for Each Soil Type Required for Effective Control)

barnyardgrass	johnsongrass (from seed)
bluegrass, annual	knotweed, prostrate
copperleaf, rhombic	nightshade, black
copperleaf, Virginia (three-seeded mercury)	panicum, brown top
dodder	pigweed, redroot
deadnettle, purple	sandbur
deadnettle, spotted	spurge, nodding
goosegrass	spurge, prostrate
groundcherry	spurge, spotted

DIFFICULT TO CONTROL WEEDS

(Commercially acceptable control of the following weeds may not always be obtained but weed populations will be suppressed when applied at the maximum labeled rate, thus resulting in less competition with the emerging and growing crop.)

buckwheat, wild	panicum, fall
canarygrass, littleseed	panicum, Texas
dock, curly (from seed)	polypogon, rabbitfoot
foxtail, giant	rocket, London
goosefoot, nettleleaf	shepherdspurse
henbit	sowthistle, annual
knotweed, silversheath	vaseygrass
ladysthumb	witchweed

WEEDS NOT CONTROLLED WITH DACTHAL

galinsoga	nutsedge
gimsonweed	ragweed, common
johnsongrass (from rhizomes)	smartweed
mustards	velvetleaf