

PM 22 50534-23

Net Contents 3 Pounds/1.4 Kilograms

ACCEPTED  
SEP 26 1995  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 50534-23

ISK BIOSCIENCES™

# Bravo W-75

**Agricultural Fungicide (75% Wettable Powder)**

Active Ingredient: Chlorothalonil (tetrachloroisophthalonitrile) .....	75.0%
Inert Ingredients: .....	25.0%
Total: .....	100.0%

**Keep Out of Reach of Children  
DANGER—PELIGRO**

See side panel for additional precautionary statements

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

ISK Biosciences Corporation  
5966 Hensley Road  
P.O. Box 8000  
Mentor, Ohio 44061-8000

BRAVO is a registered trademark  
of ISK Biosciences Corporation.

See enclosed folder for directions for use.

EPA Reg. No. 50534-23  
EPA Est. No. 37429-GA-2

3305-4018-1006-9405

**BEST COPY AVAILABLE**

OCT-13-95 09:33 FROM:

ID: 215 357 792

PAGE 2/2  
1572

## Directions For Use:

**BRAVO W-75** is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. **BRAVO W-75** is recommended for use in programs which are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

**BRAVO W-75** is effective for strategic use in programs which attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk from disease resistance exhibit a single-site mode of fungicidal action. **BRAVO W-75**, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your federal or state Cooperative Extension Service representatives for guidance on the proper use of **BRAVO W-75** in programs which seek to minimize the occurrence of disease resistance to other fungicides.