

11/17/77 50534-46 11/12/76

Ansar® 8100

Net Contents:

Herbicide

High Concentrate DSMA Powder
For Selective Post-Emergent Weed Control

| | |
|--|--------------|
| Active Ingredient: Disodium Methanearsonate* | 81.0% |
| Inert Ingredients: | <u>19.0%</u> |
| Total: | 100.0% |

Total Arsenic all in water-soluble form, expressed as elemental 33.0%

*Product equivalent to 128.5% Disodium Methanearsonate Hexahydrate

Keep Out of Reach of Children


CAUTION:

See back panel for additional precautionary statements

ANSAR is a registered trademark of ISK Biosciences Corporation.

ISK Biosciences Corporation
5966 Heisley Road
P.O. Box 8000
Mentor, Ohio 44061-8000

EPA Reg. No. 50534-46
EPA Est. No.39578-TX-1
3170-3007-1030-9509



Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through the skin.

Avoid contact with eyes, skin or clothing.

Avoid breathing dust or spray mist.

If swallowed, induce vomiting immediately by giving two glasses of water and sticking finger down throat. NEVER give anything by mouth to an unconscious person. Seek medical attention.

In case of contact, flush eyes with plenty of water for at least 15 minutes. Seek medical attention.

Wash exposed skin gently with plenty of soap and water.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- * Long-sleeved shirt and long pants
- * Waterproof gloves
- * Shoes plus socks
- * Protective eyewear
- * Chemical-resistant apron when cleaning equipment, mixing or loading

For exposures in enclosed areas:

- * A respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)

For exposures outdoors:

- * Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Follow manufacturer's instructions for cleaning maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- * Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT apply directly to water. DO NOT feed treated foliage to livestock or graze treated areas. DO NOT spray or allow drift onto vegetables, ornamentals or other desirable plants. DO NOT apply when weather conditions favor drift from target area. DO NOT contaminate water by cleaning of equipment or disposal of wastes.

Ansar® 8100 Herbicide

Read entire label before using this product.

General Information

ANSAR 8100 Herbicide is useful for selective post-emergent weed control, particularly for grassy weeds. Its activity is enhanced by addition of a suitable surfactant to the spray solution. **It is most effective on young, actively growing weeds at air temperatures above 70°F.** Its phytotoxic properties are quickly inactivated on contact with the soil.

A list of weeds controlled with ANSAR 8100 includes:

| | |
|---|---|
| Bahiagrass (<i>Paspalum sp.</i>) | Goosegrass (<i>Eleusine indica</i>) |
| Barnyardgrass (<i>Echinochloa crus-galli</i>) | Guineagrass (<i>Panicum sp.</i>) |
| Beggartick, hairy (<i>Biden pilosa</i>) | Jimsonweed (<i>Datura stramonium</i>) |
| Brachiaria spp. | Johnsongrass (<i>Sorghum halipense</i>) |
| Broomsedge | Morningglory (<i>Ipomoea spp.</i>) |
| Chickweed (<i>Stellaria media</i>) | Mustard, wild |
| Cocklebur (<i>Xanthium spinosum</i>) | Nutsedge (<i>Cyperus sp.</i>) |
| Crabgrass, large (<i>Digitaria sp.</i>) | Oats, wild (<i>Avena fatua</i>) |
| Crabgrass, smooth (<i>Digitaria sp.</i>) | Pigweed (<i>Amaranthus sp.</i>) |
| Dallisgrass (<i>Paspalum dilatatum</i>) | Puncturevine (<i>Tribulus terrestris</i>) |
| Dandelion (<i>Taraxacum officinale</i>) | Purslane, common |
| Dayflower (<i>Commelina sp.</i>) | Ragweed (<i>Ambrosia spp.</i>) |
| Fall Panicum (<i>Panicum dichotomiflorum</i>) | Sandbur (<i>Cenchrus sp.</i>) |
| Fiddleneck (<i>Amsinckia sp.</i>) | Sicklepod (<i>Cassia obtusifolia</i>) |
| Florida Beggarweed (<i>Desmodium tortuosum</i>) | Sorrel, Wood (<i>Oxalis sp.</i>) |
| Foxtail (<i>Setaria sp.</i>) | Spurge (<i>Euphorbia sp.</i>) |
| | Witchgrass (<i>Panicum capillare</i>) |

Mixing Instructions

ANSAR 8100 Herbicide is completely water soluble. However, the need for proper mixing is essential to prevent lumping of the powder. Fill the spray equipment tank with at least 80% of the required amount of water and add the required amount of ANSAR 8100 slowly, with agitation. With continued agitation add the required amount of a mild approved agricultural surfactant and the remainder of the water. Do not store spray solution in tank for a prolonged period and clean application equipment thoroughly after using by flushing with water. Although ANSAR 8100 Herbicide is only moderately corrosive, do not use in galvanized or aluminum equipment.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

176

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- * Coveralls
- * Waterproof gloves
- * Shoes plus socks
- * Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and domestic animals off treated areas until this material has been washed into the soil.

DO NOT apply this product through any type of irrigation equipment.

COTTON: ANSAR 8100 Herbicide is useful as a topical or directed spray application in cotton for post-emergent control of weeds and grasses ~~including johnsongrass, nutsedge, dallisgrass, cocklebur, ragweed, sandbur and puncture vine listed above.~~ ANSAR 8100 Herbicide should be applied to weeds in cotton at the rate of 28 pounds per acre.

If ANSAR 8100 Herbicide is to be applied topically (by either ground or aerial application), then a second directed application, one to three weeks after the first, may be made if regrowth occurs.

If ANSAR 8100 Herbicide is not applied topically, then one directed application of ANSAR 8100 Herbicide may be made, followed one to three weeks after the first, with a second directed application of either ANSAR 8100 or ANSAR 529 H.C.

26

MIXING INSTRUCTIONS: Topical by ground equipment - Mix 7.0 pounds of ANSAR 8100 in 100 gallons of water and add one to two quarts of a mild agricultural surfactant, cleared for application to growing crops, suitable for use on cotton. Apply 40 gallons of spray solution per treated acre. For topical band application in 40 inch row cotton, apply one gallon of spray for each inch of band width per acre.

Apply only when cotton has 1-2 true leaves to first square. Do not feed treated foliage to livestock, or graze treated areas.

AERIAL APPLICATION: Apply at the rate of 2.8 pounds of ANSAR 8100 Herbicide per acre. Mix 28 pounds of ANSAR 8100 and one quart of a mild agricultural surfactant, cleared for application to growing crops, for use on cotton in 50 gallons of water and apply at 5 gallons per acre. When humidity is low, mix 28 pounds of ANSAR 8100 and one quart of surfactant in 100 gallons of water and apply at 10 gallons per acre.

Do not apply by air on windy days or in such a manner as will allow spray drift to come in contact with adjacent crops on non-target areas. Do not make topical applications to cotton when drought condition or cool temperatures prevail. Apply topically only when cotton has 1-2 true leaves to first square formation.

Do not make topical applications after first square.

DIRECTED APPLICATIONS: Mix at the rate of 7.0 pounds in 100 gallons of water and add one to two quarts of a mild surfactant, cleared for application to growing crops. Apply at a broadcast rate of 40 gallons per treated acre. For band applications in 40 inch rows, apply one gallon of spray solution per inch of band width per acre of cotton.

Apply only when cotton is 3 inches high to first bloom. Do not apply after first bloom. Do not feed treated foliage to livestock, or graze treated areas.

Slight burning and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally. ANSAR 8100 Herbicide should be applied with a low volume sprayer, with satisfactory pumping and by-pass action.

CITRUS-BEARING AND NON-BEARING (Except Florida): ANSAR 8100 Herbicide is useful as a directed application in citrus orchards, such as orange, grapefruit, tangerine, lemon and lime orchards. It should be applied at the rate of 3 to 6 pounds per acre. Mix ANSAR 8100 Herbicide at the rate of 3 to 6 pounds, plus 1 to 2 quarts of a suitable surfactant, in 100 gallons of water. Apply as a directed spray in interspaces and around base of trees. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required, however, do not exceed 3 applications per year.

Do not allow spray solution to contact fruit, leaves, stems or bark. Use a shield, if necessary, for nursery plantings or young trees.

NON-CROP: ANSAR 8100 Herbicide is useful for controlling johnsongrass, ~~and similar~~ *the above listed* weeds on drainage ditch banks, rights-of-way, storage yards, and similar non-crop areas. Mix ANSAR 8100 Herbicide at the rate of 3 to 6 pounds, plus 1 to 2 quarts of a suitable surfactant in 100 gallons of water. Spray the unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes.

LAWN AND ORNAMENTAL TURF: ANSAR 8100 Herbicide is designed for the selective kill of ~~dallisgrass, sandbur, bahiagrass, nutsedge, crabgrass, chickweed, and wood sorrel~~ with *the above listed weeds* with little or no injury to tolerant lawn grasses. Tolerant grasses may be temporarily affected, especially in hot weather. Rain or waterings will revive them. Zoysia, bluegrasses, and bermuda are tolerant. DO NOT treat new lawns until after three mowings. DO NOT apply to St. Augustinegrass except for commercial sod production. Bent, St. Augustinegrass and fescue grasses are generally more sensitive, and may be temporarily discolored. DO NOT use on Dichondra or centipede. Mow lawns 1 to 1-1/2 inches high before treatment. DO NOT apply to freshly mowed St. Augustinegrass sod. Mix 1.5 ounces of ANSAR 8100 Herbicide in 1 to 2-1/2 gallons of water, containing 2 to 3 fluid ounces of a suitable surfactant, and apply to an area of 1000 sq. ft. Spray thoroughly to wet all undesirable plants. Repeat applications, 10 to 14 days apart, may be needed for good control. For applications to St. Augustinegrass, applications can be made to sod on sod farms, with the last application approximately 4-5 weeks prior to lifting the sod for harvest to allow full recovery.

DO NOT apply with hose-end applicators.

Storage and Disposal

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: DO NOT reuse empty container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants to those persons lawfully acquiring title to this product that at the time of the first sale of this product by seller that this product conformed to its chemical description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions under normal conditions of use and Buyers and users of this product assume the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO EXCEPT IN WRITING WITH A SPECIFIC REFERENCE TO THIS WARRANTY. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in the Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practices.