



SDS Biotech Corporation
Agricultural Chemicals Business
7528 Auburn Road, P.O. Box 348
Painesville, Ohio 44077

Bravo 90DG Fungicide

Supplemental Labeling

SUPPLEMENTAL DIRECTIONS FOR THE USE OF

BRAVO 90DG
(EPA Reg. No. 50534-157)

For Control of Diseases of Cranberry

ACCEPTED

NOV 13 1988

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 50534-157

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Dosage rates on this label indicate pounds of BRAVO 90DG per acre, unless otherwise stated. Applications should be made in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons (approximately 80 to 600 liters) per acre for dilute sprays and 5 to 10 gallons (approximately 20 to 40 liters) per acre for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions for ground application only are given for a crop. Application through sprinkler irrigation systems is recommended for some crops which are specified on the label below. Follow application and calibration instructions.

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Cranberry	Fruit rots, Lophodermium leaf/twig blight	3 1/2 to 5 3/4 lb.	Apply at late bloom and repeat at 10 to 14 day intervals. Under severe disease conditions use the 5 3/4 lbs/acre rate on a 10 day schedule. DO NOT apply BRAVO 90DG more than 3 times per season, or within 50 days before harvest. DO NOT apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application.

BEST AVAILABLE COPY

Warranty and Limitation of Damages: Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
=====			
			BRAVO 90DG may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions following this section.
=====			

APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

SOLID SET IRRIGATION EQUIPMENT:

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of BRAVO 90DG for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. BRAVO 90DG can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until BRAVO 90DG has been cleared from the last sprinkler head.

DO NOT operate irrigation system without safety valves or other devices to prevent back-siphoning of BRAVO 90DG into water source. Irrigation water treated with BRAVO 90DG should be maintained on the treated area until the water is absorbed by the soil. Tank containing BRAVO 90DG should be connected to the discharge side of irrigation pump or other pressurized equipment attached to irrigation line.

NOTE: It is important to read and observe the precautionary statements and all other information appearing on the BRAVO 90DG label. Use only as directed.

NOTE: This labeling must be in the possession of the user at the time of pesticide application.