


PROPOSED LABEL

BRAVO 90DG

Agricultural Fungicide
(90% Water Dispersible Granules)



Active Ingredient:

Chlorothalonil (tetrachloroisophthalonitrile)* . . . 90.0%

Inert Ingredients: 10.0%

Total: 100.0%

*Covered under U.S. Patent No. 3,290,353 and No. 3,331,735

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

See side panel for additional precautionary statements

Precaucion al Usuario: Si usted no lee ingles, no use
este producto hasta que la etiqueta haya sido
explicado ampliamente.

Fermenta Plant Protection Company
7528 Auburn Road
P.O. Box 348
Concord Township
Painesville, Ohio 44077

EPA Reg. No. 50534-157

EPA Est. No. 50534-TX-01

A BROAD SPECTRUM AGRICULTURAL FUNGICIDE

Read entire label carefully and use only as directed.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER

Corrosive, causes severe eye damage. May cause skin irritation. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or face shield when handling this product.

Avoid prolonged contact with skin. Wash exposed areas of skin with soap and warm water after handling or using.

Avoid breathing spray mist.

Do not take internally.

Note to User: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

FIRST AID

In case of contact with eyes, immediately flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes immediately.

Wash exposed areas of skin with soap and warm water after handling or using.

Note to Physician: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

BRAVO 90DG is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases.

BRAVO 90DG can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

Do not use on greenhouse-grown crops.

Do not combine **BRAVO 90DG** in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. Do not combine **BRAVO 90DG** with Dipel 4L, Triton AG-98, or Triton B-1956 as phytotoxicity may result from the combination when applied to some crops on this label.

Dipel is a registered trademark of Abbott Laboratories. Triton and Triton B-1956 are trademarks of Rohm and Haas Company.

Dosage rates on this label indicate pounds of **BRAVO 90DG** per acre, unless otherwise stated. Under conditions favoring disease development the high rate specified and shortest application interval should be used.

The required amount of **BRAVO 90DG** should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of **BRAVO 90DG** in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

Note to User: Do not rotate to crops other than those listed on labels within 12 months of the last treatment. After 12 months from the last application, leafy vegetables (i.e. spinach, lettuce, kale, etc.) may also be rotated.

Reentry: **Note to User:** Do not enter treated area to perform hand labor within 24 hours of application unless protective clothing is worn.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Oral and written warnings must include the following information: "DANGER. Area treated with **BRAVO 90DG** on (date of application). Do not enter without appropriate protective clothing for 24 hours. In case of accidental exposure, wash exposed area with plenty of water and get medical attention. For further information see 'Precautionary Statements' on the label."

BEST AVAILABLE COPY

FIELD AND ROW CROPS

Apply Bravo 90DG in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft application. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See application and calibration instructions below.

BRAVO 90DG:

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Potato	Late blight, Early blight, Botrytis vine rot	5/8 lb -- then -- 7/8 to 1 1/4 lb/A	<p>Apply as a banded treatment directed over the rows beginning when vines are first exposed to blight. Repeat applications at 7 to 10 days when vines close, then increase rate as disease becomes severe.</p> <p>In addition to the early-season applications, when vines close between rows increase water carrying capacity, denser canopy and begin broadcast applications. Continue applications at 7 to 10 day intervals at the highest registered rate weekly when disease is severe.</p> <p>BRAVO 90DG may be applied through any type of equipment (solid set, portable wheel, or motorized lateral move systems only). Follow the recommended interval between applications when used in a broadcast section. Follow calibration and application directions in the product section.</p>

TREE AND ORCHARD CROPS:

Apply Bravo 90DG in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, Bravo 90DG may be applied with aircraft using at least 20 gallons per acre. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of Bravo 90DG listed may be used. Do not allow livestock to graze in treated areas. The following spray volumes are recommended as gallons of spray per acre:

CROP	SPRAY VOLUME (Gallons per Acre)
Peach, Nectarine, Apricot, Tart Cherry, Plum, Prune	20 (concentrate) to 300 (full dilute)
Sweet Cherry	20 (concentrate) to 400 (full dilute)

CROP	DISEASES	BRAVO 90DG RATE PER ACRE	RATE PER 100 GAL*	APPLICATION DIRECTIONS
Peach, Nectarine, Apricot, Cherry, Plum, Prune	Brown rot blossom blight, Lacy (russet) scab (plum/prune)	2 1/2 to 4 1/2 lb	7/8 to 1 1/8 lb	Use 3 1/2 to 4 1/2 lb per acre on trees taller than 20 feet and 2 1/2 to 3 1/2 lb per acre on smaller trees. Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.

* Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.