

PM - 22

Reg # 50534-158

146
7**DSMA 4****Net Contents: 55 Gallons/208 Liters****DSMA For Postemergence Weed Control
in Cotton, Turf, and Non-Crop Areas****Contains No Surfactant****Active Ingredient:** Disodium Methanearsonate

24.07%

Inert Ingredients:

75.93%

Total:

100.00%

Total Arsenic, all in water-soluble form, expressed as elemental

9.84%

Product contains 4 pounds Disodium Methanearsonate Hexahydrate equivalent per gallon.

AntidoteDrink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger.
DO NOT induce vomiting or give anything by mouth to an unconscious person. Get medical attention.**Read entire label before using this product.****Notice:** Use of this material in a manner, or at any time other than in accordance with the directions, may produce plant injury, excessive residues, or other undesirable results.**Keep Out of Reach of Children**
CAUTION:

See side panel for additional precautionary statements

ISK Biosciences Corporation

5966 Heisley Road


P.O. Box 8000

Mentor, Ohio 44061-8000

EPA Reg. No. 50534-158

EPA Est. No. 50534-TX-01

2410-0000-9120-9304



146

Precautionary Statements**Hazards to Humans and Domestic Animals****Caution**

Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing spray mist. Wash hands after using.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- * Long-sleeved shirt and long pants
- * Waterproof gloves
- * Shoes plus socks
- * Protective eyewear
- * Chemical-resistant apron when cleaning equipment, mixing or loading

For exposures in enclosed areas:

- * A respirator with either an organic vapor-removing cartridge with a pre-filter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C) or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)

For exposure outdoors:

- * Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Follow manufacturer's instructions for cleaning maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- * Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

DO NOT feed treated foliage to livestock or graze treated areas.

DO NOT spray or allow drift onto vegetables, ornamentals or other desirable plants.

DO NOT apply when weather conditions favor drift from target area.

DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. DO NOT contaminate water by disposal of equipment wash waters.

DO NOT contaminate water by cleaning of equipment or disposal of wastes.

Avoid storage near food and feed products.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- * Coveralls
- * Waterproof gloves
- * Shoes plus socks
- * Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and domestic animals off treated areas until this material has been washed into the soil.

DO NOT apply this product through any type of irrigation equipment.

Cotton: DSMA 4 is useful as a directed postemergence spray for controlling weeds in cotton such as barnyardgrass, cocklebur, smooth and large crabgrass, dallisgrass, goosegrass, johnsongrass and nutsedge.

DO NOT feed treated foliage to livestock or graze treated areas.

Preplant or postplant up to cracking application on cotton: A single ground or aircraft application of DSMA 4 can be made to prepared cotton seedbeds when planting has been delayed and weeds have emerged, or as a post-plant treatment, but no later than initial cracking of soil in field before emergence of cotton.

Planting of cotton may immediately follow the preplant application. For ground application mix at a rate of 7 pints of DSMA 4 and 1 quart of a nonionic surfactant, cleared for application to growing crops, in 40 gallons of water for broadcast ground application or in 5 to 10 gallons of water for aircraft application and apply to one acre using a properly calibrated sprayer.

Postemergent, over-the-top application on cotton:

A single ground or aircraft application may be made as an over-the-top broadcast spray to actively growing cotton as a salvage operation when cotton is 3 to 6 inches high or up to early first square stage, whichever occurs first. For ground application, mix DSMA 4 at a rate of 3 to 4 pints in 40 gallons of water plus 1 to 2 pints of a nonionic surfactant, cleared for application to growing crops, and apply to one acre using a properly calibrated sprayer. For aerial application, mix DSMA 4 at a rate of 3 to 4 pints in 5 to 10 gallons of water plus 1/2 pint of a nonionic surfactant, cleared for application to growing crops, and apply to one acre using a properly calibrated sprayer. Apply as an over-the-top broadcast spray only when cotton is 3 to 6 inches high or up to early first square stage, whichever occurs first.

DO NOT apply as an over-the-top broadcast spray after cotton is more than 6 inches high or after early first square stage.

Slight burning or reddish discoloration of cotton foliage may occasionally be seen following recommended treatment.

Directed application: Mix 9 quarts of DSMA 4 and 1 to 2 quarts of a nonionic surfactant, cleared for application to growing crops, in 100 gallons of water. Apply the solution as a directed spray at rate of 40 gallons per acre when weeds are small. For band applications, apply 1 gallon of above diluted spray per acre for each 1 inch band width to be treated of cotton grown on 40 inch row spacing. A second or repeat application, if needed, should be timed about 1 to 3 weeks after the first application. Keep spray off cotton foliage.


Apply Only When Cotton is 3 Inches High to First Bloom. Do Not Apply After First Bloom.

Slight burning or reddish discoloration of cotton foliage may occasionally be seen following recommended treatment; however cotton plants will develop normally and yield will not be affected.

Ornamental Turfgrass: DSMA 4 can be used for selective control of bahiagrass, barnyardgrass, chickweed, smooth and large crabgrass, dallisgrass, nutsedge, sandbur and wood sorrel with little or no injury to well-established, actively growing turfgrasses. Mow turfgrass to height of 1 to 1-1/2 inches before treatment. Mix 3 fluid ounces (6 tablespoons) of DSMA 4 plus 1 ounce of a nonionic surfactant in 1 to 2-1/2 gallons of water for application to an area of 1,000 square feet. Application should be uniform and thorough to adequately wet all undesirable plants. Two or more repeat treatments at 14 day intervals may be necessary. Make applications during warm weather when temperature is between 80 and 90°F. DO NOT water turf for at least 24 hours after application. Turfgrasses may be temporarily discolored. Bermuda-grass, bluegrass and zoysiagrass have shown tolerance to properly applied DSMA 4. Injury may result if applied to bentgrass and fescues. DO NOT apply to St. Augustine-grass, carpetgrass, centipedegrass or to Dichondra. DO NOT apply with hose-end applicators.

Agricultural Plantings: DSMA 4 is useful as a directed application in non-bearing vineyards and in non-bearing deciduous fruit, nut and citrus orchards such as almond, apple, apricot, cherry, grapefruit, lemon, lime, orange, peach, pear, plum, prune and tangerine orchards. It should be applied at the rate of 3-1/2 to 6-1/2 quarts per acre. Mix DSMA 4 at the rate of 3-1/2 quarts plus 1 quart of a nonionic surfactant in 50 gallons of water. Apply as a directed spray in interspaces and around bases of trees or vines. Spray unwanted vegetation to just short of run-off. If regrowth occurs, reapply as required. However, do not exceed 3 applications per year.

DO NOT allow spray solution to contact leaves, stem or bark. Use a shield, if necessary, for nursery plantings or young trees. DO NOT apply around trees or vines from which fruit will be harvested within one year of treatment.



Non-Crop: DSMA 4 is effective in controlling barnyardgrass, cocklebur, smooth and large crabgrass, dallisgrass, goosegrass, johnsongrass, nutsedge and many similar weeds on drainage ditchbanks, fence rows, rights-of-way, storage yards and similar non-crop areas. Application should be made when weeds are small and conditions are favorable for good weed growth. Mix at a rate of 1 gallon of DSMA 4 plus 1 quart of a nonionic surfactant in 40 gallons of water. Spray undesirable vegetation thoroughly to point of run-off. Adequate coverage and complete wetting of foliage is important for effective control. Repeat applications may be necessary if regrowth occurs. DO NOT contaminate waters used for domestic purposes, or by animals including wildlife and aquatic life, or water being used for irrigation purposes.

Storage and Disposal

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of at an approved waste disposal facility.

Container Refilling and Cleaning: This is a returnable dedicated container to be refilled only with ISK Biotech's Methanearsonate products by an authorized Repackager. Return this container to the Repackager from which it was purchased for refilling. If the seals are intact, do not rinse this container between refills. If the seals are not intact, or, other evidence exists that tampering may have occurred, triple rinse (or equivalent) before returning container to authorized Repackager. Add rinsate to the spray tank as make-up water for normal spray dilution of concentrate product.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless expressly provided in the Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practices.

