

50534-188


08/02/2001

1/28

Page 1 of 28

Bravo 720

BRAVO188RSE - 5/10/01



BRAVO® 720
Agricultural Fungicide

ACTIVE INGREDIENT:

Chlorothalonil (tetrachloroisophthalonitrile) 54.0%

INERT INGREDIENTS: 46.0%

TOTAL 100.0%

Contains 6.0 Pounds Chlorothalonil Per Gallon
(720 grams per liter)

EPA Reg. No. 50534-188

EPA Est. No. 50534-TX-1

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

GB Biosciences Corporation

1800 Concord Pike

Wilmington, DE 19850-5458

**CONDITIONS OF SALE
AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of GB Biosciences or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold GB Biosciences and Seller harmless for any claims relating to such factors.

GB Biosciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or GB Biosciences, and Buyer and User assume the risk of any such use. GB BIOSCIENCES MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall GB Biosciences or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF GB BIOSCIENCES AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF GB BIOSCIENCES OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

GB Biosciences and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of GB Biosciences.

A Broad Spectrum Agricultural Fungicide
Read entire label carefully and use only as directed

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes eye irritation. May cause skin irritation. May be a potential skin sensitizer.

DO NOT get in eyes. Avoid prolonged contact with skin. Avoid breathing spray mist. DO NOT take internally.

NOTE TO USER: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact x lenses, if present, after the first 15 minutes, then continue rinsing eye. Call a poison control center for doctor for treatment advice.

Wash exposed areas of skin with soap and warm water after handling or using.

NOTE TO PHYSICIAN: Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL 1-800-F-A-S-T-M-E-D (327-8633).

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

Personal Protective Equipment (PPE)

Mixers, Loaders, Applicators and all other handlers must wear:

- long-sleeved shirt and long pants,
- chemical resistant gloves made of any waterproof material - Category A (e.g. barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton),
- shoes plus socks,
- protective eyewear,
- and a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with

this product's concentrate. DO NOT reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to aquatic invertebrates, and wildlife. DO NOT apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment wash water or rinsate. This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

See attached booklet for Directions for Use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow workers to enter treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

(1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.

(2) Workers must be informed, in a manner they can understand:

- that residues in the treated area may be highly irritating to their eyes,
- that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes,
- that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water, and
- how to operate the eyeflush container.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in a cool place. Protect from excessive heat.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: DO NOT reuse empty container. Triple rinse (or equivalent), and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

RETURNABLE REFILLABLE CONTAINER: If Bravo 720 is packaged in a returnable refillable container, then, after use, do not rinse container. Return container intact to point of purchase. This container must only be refilled with Bravo 720. DO NOT REUSE THE CONTAINER FOR ANY OTHER PURPOSE. Before refilling, inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn threads on closure devices. Check for leaks after refilling and before transport.

GENERAL INFORMATION

BRAVO[®] 720 is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. BRAVO 720 is recommended for use in programs which are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

BRAVO 720 is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk from disease resistance exhibit a single-site mode of fungicidal action. BRAVO 720, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your federal or state Cooperative Extension Service representatives for guidance on the proper use of BRAVO 720 in programs which seek to minimize the occurrence of disease resistance to other fungicides.

BRAVO 720 can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

GENERAL PRECAUTIONS AND RESTRICTIONS

DO NOT use on greenhouse-grown crops.

DO NOT combine BRAVO 720 in spray tank with pesticides, surfactants or fertilizers, unless your prior

use has shown the combination physically compatible, effective and noninjurious under your conditions of use. DO NOT combine BRAVO 720 with Dipel 4L, Foil, Triton AG-98, Triton B-1956 or Latron B-1956 as phytotoxicity may result from the combination when applied to the crops on this label.

This product must not be applied within 150 feet (for aerial applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Spray Drift Precautions:

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Aerial Drift Reduction Advisory Information.

([This section is advisory in nature and does not supercede the mandatory label requirements.]

Information on Droplet Size:

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential but will not prevent drift if applications are made improperly, or under unfavorable conditions (See Wind, Temperature).

Controlling Droplet Size:

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.

- **Nozzle Orientation** - Orienting the nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

Boom Length:

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Wind:

Drift potential is lowest between wind speeds of 2 - 10mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity:

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions:

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

APPLICATION:

Dosage rates on this label indicate pints of BRAVO 720 per acre, unless otherwise stated. Under conditions favoring disease development the high rate specified and shortest application interval should be used.

Note: Slowly invert container several times to assure uniform mixture.

The required amount of BRAVO 720 should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of BRAVO 720 in a clean container and add to the

spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

Field and Row Crops

Apply BRAVO 720 in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See application and calibration instruction below.

Application and Calibration Techniques for Sprinkler Irrigation

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). DO NOT apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject BRAVO 720 into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either

automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. DO NOT apply when wind speed favors drift beyond the area intended for treatment.

BRAVO 720 may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix recommended amount of BRAVO 720 for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until BRAVO 720 has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of BRAVO 720 for acreage to be covered with water so that the total mixture of BRAVO 720 plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. No agitation should be required. BRAVO 720 can be injected at the

beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until BRAVO 720 has been cleared from last sprinkler head.

DIRECTIONS FOR APPLICATION:

(

12/28

Page 11 of 28

Bravo 720

BRAVO188R9E - 5/10/01

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Asparagus	Rust Purple Spot Cercospora leaf blight	2 to 4 pints (1.5 to 3.0)	Use water volumes of 25-50 gallons per acre. Begin applications following final harvest of spears. Repeat applications at 14-28 day intervals (the minimum re-treatment interval is 14 days), depending on disease pressure. Use the higher rate and shorter interval if disease severity begins to increase during the season or weather conditions are conducive for severe epidemics. Do not apply within 190 days of the harvest of spears in the following season. Do not apply more than 12 pints BRAVO 720 (9.0 lb ai) per acre during each growing season.
Bean (Snap)	Rust	1-3/8 to 3 pts (1.0 to 2.25)	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat as necessary (the minimum re-treatment interval is 7 days) to maintain control. DO NOT apply more than 12 pints BRAVO 720 (9 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.
	Botrytis blight (gray mold)	3 pts (2.25)	
Beans (Dry) Navy, Pinto, Kidney, Lima, Blackeye	Rust, Anthracnose, Downy mildew, Cercospora leaf spot (blackeye only), Ascochyta blight	1-3/8 to 2 pts (1.0 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days). For use only on beans to be harvested dry with pods removed. DO NOT apply more than 4 times per growing season. DO NOT apply more than 8 pints BRAVO 720 (8 lbs a.i.) per acre during each growing season. DO NOT apply within 14 days before harvest. BRAVO 720 may be applied through sprinkler irrigation equipment. See calibration directions which appear on the product label.

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Blueberries	Anthracnose Mummy Berry	3 to 4 pints (2.25 to 3.0)	<p>Use 3 to 4 pints in sufficient water to obtain adequate coverage, normally 20-100 gallons per acre. Begin applications at budbreak (green tip). Repeat applications through the bloom period and repeat at 10-day intervals (the minimum re-treatment interval is 10 days). Under heavy disease pressure, use the higher rate.</p> <p>Do not apply after full bloom or within 42 days of harvest. Do not apply more than 12 pints BRAVO 720 (9.0 lb ai) per acre during each growing season.</p>
Cabbage, Chinese Cabbage (tight-headed varieties only) Cauliflower, Broccoli, Chinese Broccoli, Brussels Sprouts	Alternaria leaf spot, Downy mildew	1-1/2 pts (1.125)	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary (the minimum re-treatment interval is 7 days) to maintain control. DO NOT apply more than 16 pints BRAVO 720 (12 lbs a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest to Chinese cabbage or Chinese broccoli.
	Ring spot (California only)	2 pts (1.5)	For field-seeded brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary (the minimum re-treatment interval is 7 days) to maintain control.

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Celery	Cercospora (Early) blight, Septoria (Late) blight, Basal stalk rot (<i>Rhizoctonia solani</i>)	2 to 3 pts (1.5 to 2.25)	<p>Use in sufficient water to obtain adequate coverage. Start applications when transplants are set in the field and repeat at a 7 day interval as needed to maintain control (the minimum re-treatment interval is 7 days). DO NOT apply more than 24 pints BRAVO 720 (18 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.</p> <p>BRAVO 720 may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). See calibration directions preceding this section.</p>
	Pink rot (Suppression - 7 day schedule)	3 pts (2.25)	
	Early blight, Late blight	1-1/2 to 2 pts (1.125 to 1.5) per 100 gal.	For celery seedbeds, apply in a spray volume of 125 gallons per acre twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.
Corn (Sweet), Corn grown for seed	Helminthosporium leaf blights, Rust	3/4 to 2 pts (0.6 to 1.5)	<p>Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at 7 day intervals as required to maintain control (the minimum re-treatment interval is 7 days). Under severe disease conditions, use 1-1/2 to 2 pints BRAVO 720 per acre. DO NOT apply more than 12 pints BRAVO 720 (9 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT apply to sweet corn to be processed. DO NOT allow livestock to graze in treated fields. DO NOT ensile treated corn or use as livestock forage.</p>

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Cranberry	Fruit rots, Lophodermium leaf/twig blight	4 to 6-1/2 pts (3.0 to 4.9)	<p>Apply at early bloom and repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days). Under severe disease conditions, use the 6-1/2 pint per acre rate on a 10 day schedule. DO NOT apply BRAVO 720 more than 3 times per season. DO NOT apply more than 20 pints BRAVO 720 (15 lbs. a.i.) per acre during each growing season. DO NOT apply within 50 days of harvest. DO NOT apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application.</p> <p>BRAVO 720 may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.</p>
Cucurbits: Cucumber, Cantaloupe, Muskmelon, Honeydew melon, Watermelon, Squash, Pumpkin	Anthracnose, Downy mildew, Target spot	1-1/2 to 2 pts (1.125 to 1.5)	<p>Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum re-treatment interval is 7 days). DO NOT apply more than 21 pints BRAVO 720 (15.75 lbs. a.i.) per acre during each growing season.</p> <p>BRAVO 720 may be applied through sprinkler irrigation equipment (solid set, portable wheel move or center pivot systems only). See calibration directions preceding this section.</p> <p>Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply BRAVO 720 to watermelons when any of the following conditions are present:</p> <ol style="list-style-type: none"> 1. Intense heat and sunlight; 2. Drought conditions; 3. Poor vine canopy; 4. Other crop and environmental conditions which may be conducive to increased natural sunburn. <p>DO NOT combine BRAVO 720 with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use.</p>

16/28

Page 16 of 28
Bravo 720
BRAVO188RSE - 5/10/01

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
	Cercospora leaf spot, Gummy stem blight (black rot), Alternaria leaf blight, Scab, Powdery mildew (Sphaerotheca only)	2 to 3 pts (1.5 to 2.25)	

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Grasses Grown for Seed	Stem rust, Leaf rust, Stripe rust, Septoria leaf spot, Glume blotch, Bipolaris and Drechslera leaf spots	1 to 1-1/2 pts (0.75 to 1.125)	Use in sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat applications at 14 day intervals (the minimum re-treatment interval is 14 days). DO NOT apply more than 6 pints BRAVO 720 (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze in treated areas or feed treated plant parts to livestock.
	Selenophoma (eyespot)	1 to 2 pts (0.75 to 1.5)	
Mango	Anthracnose	2 to 3.5 pints (1.5 to 2.6)	Use a water volume of 20 to 300 gallons per acre. Begin applications at early bloom and repeat on a 7-14 day interval until early fruit development. Begin the season with the 2 pint rate on a 14-day interval. If disease pressure is severe, use the higher rate and shorter interval. The minimum re-treatment interval is 7 days. Do not apply within 21 days of harvest. Do not apply more than 32 pints BRAVO 720 (24 lb ai) per acre during each growing season.
Mint (Indiana, Michigan and Wisconsin only)	Rust, Septoria leaf spot	1-3/8 pts (1.0)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground and aircraft applications. Begin applications when emerging plants are 4 - 8 inches high. Repeat applications at 7 to 10 day intervals or as necessary to maintain control (the minimum re-treatment interval is 7 days). DO NOT apply more than 3 times per season. DO NOT apply more than 4 pints BRAVO 720 (3 lbs. a.i.) per acre during each growing season. DO NOT apply within 80 days of harvest. DO NOT feed fresh or extracted mint hay from treated fields to livestock.

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS												
Onion (Dry bulb) and Garlic	Botrytis leaf blight/blast. Botrytis neck rot (suppression), Purple blotch, Downey mildew (suppression)	1 to 3 pts (0.75 to 2.25)	<p>Apply in sufficient water to obtain thorough coverage of tops. BRAVO 720 is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows:</p> <table border="1"> <thead> <tr> <th></th><th data-bbox="987 636 1133 751">Low Disease Hazard & Prior to Infection</th><th data-bbox="1170 636 1344 751">Low Disease Hazard & Some Disease Present</th><th data-bbox="1382 667 1471 751">High Disease Hazard</th></tr> </thead> <tbody> <tr> <td data-bbox="902 783 1000 842">Rate per Acre:</td><td data-bbox="1024 783 1073 814">1 pt</td><td data-bbox="1170 783 1268 814">1-3/8 pts</td><td data-bbox="1382 783 1430 814">3 pts</td></tr> <tr> <td data-bbox="902 846 1122 877">Frequency:</td><td data-bbox="1024 846 1122 877">10 days</td><td data-bbox="1170 846 1308 877">7 to 10 days</td><td data-bbox="1382 846 1430 877">7 days</td></tr> </tbody> </table> <p>For suppression of neck rot (Botrytis spp.) during storage, a minimum of three weekly applications prior to lifting, using 1-3/8 to 3 pints of BRAVO 720 per acre, is recommended.</p> <p>The minimum re-treatment interval is 7 days. DO NOT apply more than 20 pints BRAVO 720 (15 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.</p>		Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard	Rate per Acre:	1 pt	1-3/8 pts	3 pts	Frequency:	10 days	7 to 10 days	7 days
	Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard												
Rate per Acre:	1 pt	1-3/8 pts	3 pts												
Frequency:	10 days	7 to 10 days	7 days												
Onion (Green bunching), Leek, Shallot, Onion grown for seed	Botrytis leaf blight/ (blast), Purple blotch, Downy mildew (suppression)	1-1/2 to 3 pts (1.125 to 2.25)	<p>Use in sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease (the minimum re-treatment interval is 7 days). Use the high rate and a 7 day schedule of applications when heavy dew or rain persist. DO NOT apply more than 9 pints BRAVO 720 (6.75 lbs. a.i.) per acre during each growing season. DO NOT apply more than 3 times per season or within 14 days of harvest on green bunching onions, leeks or shallots. If additional disease control is needed before harvest, use another registered fungicide.</p>												

19/28

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Papaya	Alternaria fruit spot, Anthracnose, Stem end rot	2 to 3 pts (1.5 to 2.25)	Apply with ground equipment only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days). DO NOT apply more than 9 pints BRAVO 720 (6.75 lbs a.i.) per acre during each growing season.
Parsnip	Alternaria leaf spot, Downy mildew, Anthracnose, Botrytis blight (gray mold), Bottom rot (Rhizoctonia)	1-1/2 to 2 pts (1.125 to 1.5)	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule (the minimum re-treatment interval is 7 days). DO NOT apply more than 4 times per season or within 10 days of harvest. DO NOT apply more than 8 pints BRAVO 720 (6 lbs. a.i.) per acre during each growing season.
Passion Fruit (Hawaii only)	Alternaria fruit and leaf spot, (passion fruit brown spot)	2 pts (1.5)	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when fruit spots appear (April to July) and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days). DO NOT apply more than 10 pints BRAVO 720 (7.5 lbs a.i.) per acre during each growing season.

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Peanut	Early leafspot (Cercospora), Late leafspot (Cercosporidium)	1 to 1-1/2 pts (0.75 to 1.125)	<p>Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting; repeat at 14 day intervals (the minimum re-treatment interval is 14 days). When conditions favor late leafspot or when rust or web blotch occur, apply 1-1/2 pints BRAVO 720 per acre at 14 day intervals for the remainder of the season. DO NOT apply more than 12 pints BRAVO 720 (9 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze in treated areas. DO NOT feed hay or threshings from treated fields to livestock.</p> <p>BRAVO 720 may be applied through sprinkler irrigation equipment. Use 1-1/2 pints BRAVO 720 per acre in solid set, portable wheel move, center pivot, motorized lateral move or traveling gun sprinkler irrigation equipment. See calibration directions preceding this section.</p>
	Rust, Web blotch	1-1/2 pts (1.125)	
Potato	Late blight, Early blight, Botrytis vine rot	3/4 pt (0.6) - then - 1 to 1-1/2 pts (0.75 to 1.125)	<p>Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).</p> <p>Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur.</p> <ul style="list-style-type: none"> • Vines close within the rows; • Late blight forecasting measures 18 disease severity values (DSV); • The crop reaches 300 P-days <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. DO NOT apply more than 15 pints of BRAVO 720 (11.25 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.</p> <p>BRAVO 720 may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). DO NOT exceed a 10 day interval between applications when using this technique. See calibration directions preceding this section.</p>

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Soybean Determinate (Southern) Varieties	Anthracnose, Diaporthe pod and stem blight, Frogeye leaf spot (<i>Cercospora sojina</i>), Purple seed stain, Cercospora leaf blight (<i>Cercospora kikuchii</i>), Septoria brown spot		Apply in sufficient water to obtain complete coverage, using at least five gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. BRAVO 720 may be applied through sprinkler irrigation equipment. Follow application and calibration directions preceding this section. The minimum re-treatment interval is 14 days. DO NOT exceed total of 3 applications per season. DO NOT apply more than 6 pints BRAVO 720 (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 6 weeks of harvest. DO NOT feed treated parts to livestock or allow grazing in treated fields.
		1-1/2 to 2-1/4 pts (1.125 to 1.7)	Two application program - Make the first application at early pod set (R3 stage, when majority of pods are 1/8 to 3/8 inch in length) and the second at beginning of seed formation (R5) which occurs about 14 days later.
		1 to 2 pts (0.75 to 1.5)	Three application program - Make the first application at the beginning of flowering (R1), the second at early pod set (R3) and the third at beginning of seed formation (R5).
	Stem canker (<i>Diaporthe phaseolorum</i> var. <i>caulivora</i>)	1 pt (0.75)	Apply in 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make the application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease make a second and third application. Make all applications at 10 to 14 day intervals.
Soybean Indeterminate (Northern) Varieties	Anthracnose, Diaporthe pod and stem blight, Frogeye leaf spot (<i>Cercospora sojina</i>), Purple seed stain, Cercospora leaf blight (<i>Cercospora kikuchii</i>), Septoria brown spot		Apply in sufficient water to obtain complete coverage, using at least five gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. BRAVO 720 may be applied through sprinkler irrigation equipment. Follow application and calibration directions preceding this section. The minimum re-treatment interval is 14 days. DO NOT exceed total of 3 applications per season. DO NOT apply more than 6 pints BRAVO 720 (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 6 weeks of harvest. DO NOT feed soybean hay or threshings from treated fields to livestock.

CROP	DISEASES	RATE PER ACRE (lbs. a.i./A)	APPLICATION DIRECTIONS
Soybean Indeterminate (Northern) Varieties		1-1/2 to 2-1/4 pts (1.125 to 1.7)	Two application program - Make the first application when largest pods are 1 to 1-1/2 inches in length and make the second application 14 days later. BRAVO 720 may be co-applied with BENLATE® SP as a tank mix for disease control in indeterminate (northern) soybeans. Use 1 pint of BRAVO 720 plus 8 ounces of BENLATE SP per acre. Make the first application when pods near the top of plants are 1/2 to 1 inch in length and a second application 14 days later.
		1 to 2 pts (0.75 to 1.5)	Three application program - Make the first application one week after first flowering and continue applications at 14 day intervals.
Tomato	FOLIAGE (apply every 7-10 days): Early blight, Late blight, Gray leaf spot, Gray leaf mold, Septoria leaf spot, Target spot	1-3/8 to 2 pts (1.0 to 1.5)	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Use the highest rate and shortest interval specified when disease conditions are severe. The minimum re-treatment interval is 7 days. DO NOT apply more than 20 pints BRAVO 720 (15 lbs. a.i.) per acre during each growing season. BRAVO 720 may be combined in the spray tank with EPA-registered pesticide products that claim copper as the active ingredient and are labeled for control of bacterial diseases of tomatoes. Check the copper manufacturer's label for specific instructions, precautions and limitations prior to mixing with BRAVO 720. DO NOT use with Copper-Count® N in concentrated spray suspensions. BRAVO 720 may be applied through sprinkler irrigation equipment (solid set or portable wheel move systems only). See calibration directions preceding this section.
	FRUIT (apply every 7-14 days beginning at fruit set): Anthracnose, Alternaria fruit rot (black mold), Botrytis gray mold, Late blight fruit rot Rhizoctonia fruit rot	2 to 2-3/4 pts (1.5 to 2.1)	

Tree and Orchard Crops

Apply BRAVO 720 in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy.

If application with ground equipment is not feasible, BRAVO 720 may be applied with aircraft using at least 20 gallons of spray per acre. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of BRAVO 720 listed may be used. DO NOT allow livestock to graze in treated areas. The following spray volumes are recommended as gallons of spray per acre:

CROP		SPRAY VOLUME (Gallons per Acre)	
Pistachios		20 (concentrate) to 200 (full dilute)	
Almonds, Peach, Nectarine		20 (concentrate) to 300 (full dilute)	
Apricot, Tart Cherry, Plum, Prune			
Filberts, Sweet Cherry		20 (concentrate) to 400 (full dilute)	
Conifers:	Dilute	Concentrate	
Forest stands	Not used	10 to 20 (aircraft)	
Christmas trees	100	10 to 50 (aircraft or ground equipment)	
Nursery beds	100	5 to 10 (ground equipment only)	

24/28

24
Page 22 of 28
Bravo 720

BRAVO188RSE - 5/10/01

CROP	DISEASES	BRAVO 720 RATE PER (lbs. a.i. per)		APPLICATION DIRECTIONS
		ACRE	100 GAL*	
Almonds	Blossom blight/brown rot	4 pints (3.0)	1.33 pints (1.0)	<p>Use water volumes of 20-300 gallons per acre. For blossom blight, begin application at popcorn (pink bud) and follow with an application at full bloom. If weather is still conducive for disease development, another application may be made at petal fall.</p> <p>For control of shothole, make an application in the autumn at leaf fall. In the spring, make the first application at budbreak, followed by an application at shuck split to control nut infections and to control scab.</p> <p>Do not apply within 150 days of harvest.</p> <p>Do not apply more than 25 pints BRAVO 720 (18.75 lb ai) per acre during each growing season (leaf fall through shuck split).</p>
	Shothole			
	Scab			
Filberts (Hazelnuts)	Eastern filbert blight	4 pints (3.0)	1.33 pints (1.0)	<p>Use a water volume of 20 to 300 gallons per acre. Begin applications at the onset of disease or when weather conditions favor disease development. Make applications on a 14-28 day schedule, using the shorter interval under heavy disease pressure (the minimum re-treatment interval is 14 days).</p> <p>Do not apply with oils, other pesticides, surfactants or fertilizers. Do not apply within one week of an oil-based pesticide application.</p> <p>Do not apply through irrigation.</p> <p>Do not apply within 120 days of harvest.</p> <p>Do not apply more than 12 pints BRAVO 720 (9 lb ai) per acre during each growing season.</p>

25/28

CROP	DISEASES	BRAVO 720 RATE PER		APPLICATION DIRECTIONS
		(lbs. a.i. per)		
		ACRE	100 GAL*	
Peach, Nectarine, Apricot, Cherry, Plum, Prune	Leaf curl, Coryneum blight (shothole)	3-1/8 to 1 to 4-1/8 pts (2.3 to 3.1)	1-3/8 pts (0.75 to 1.0)	For best control of both diseases apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of BRAVO 720 for control of leaf curl may be made at any time prior to budswell the following spring. Where Coryneum blight (shothole) occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections.
DO NOT apply more than 20-1/2 pints BRAVO 720 (15.4 lbs. a.i.) per acre during each growing season. The minimum re-treatment interval is 10 days.				
	Lacy (russet) scab (plum/prune)	3-1/8 to 1 to 4-1/8 pts (2.3 to 3.1)	1-3/8 pts (0.75 to 1.0)	
	Cherry leaf spot, Peach, Nectarine, Apricot scab; Black knot (cherry, plum)	3-1/8 to 1 to 4-1/8 pts (2.3 to 3.1)	1-3/8 pts (0.75 to 1.0)	
				Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
				In addition to the bloom application listed above, make one application at shuck split. DO NOT apply BRAVO 720 after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide.
				For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10-14 days later.

CROP	DISEASES	BRAVO 720 RATE PER		APPLICATION DIRECTIONS
		ACRE	(lbs. a.i. per) 100 GAL*	
Platanillo	Alcornaria late blight	6 pints (4.5)	3 pints (2.25)	Use a water volume of 20 to 200 gallons per acre. Make the first application at the beginning of the blossom period followed by an application at full bloom. Make additional applications as required on a 28-day schedule. (The minimum re-treatment interval is 28 days).
	Botryosphaeria blight			
	Septoria leaf spot	4 to 6 pints (3.0 to 4.5)	2 to 3 pints (1.50 to 2.25)	Do not apply within 14 days of harvest. Do not apply more than 30 pints BRAVO 720 (22.5 lb ai) per acre during each growing season. NOTE: Use of this product may result in speckling or reddening of the fruit hull (epicarp). This affect is superficial and has not resulted in any change in nut quality.
	Botrytis blight			
Conifers (Pine, Spruce) DO NOT apply more than 22 pints BRAVO 720 (16.5 lbs. a.i.) per acre during each growing season. The minimum re-treatment interval for established trees is 21 days. The minimum re-treatment interval in nursery beds is 7 days.	Swiss needlecast	2-3/4 to 5-1/2 pts (2.1 to 4.125)	2-3/4 to 5-1/2 pts (2.1 to 4.125)	Single application technique: In Christmas tree plantations or forest stands make one application in the spring when new shoot growth is 1/2 to 2 inches in length.
	Sclerodermis canker (pines), Swiss needlecast	1-1/2 to 2-3/4 pts (1.125 to 2.1)	1-1/2 to 2-3/4 pts (1.125 to 2.1)	Make the first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.

CROP	DISEASES	<u>BRAVO 720 RATE PER</u> (lbs. a.i. per)		APPLICATION DIRECTIONS
		ACRE	100 GAL*	
	Sirococcus tip blight	2 to 3-1/2 pts (1.5 to 2.6)	2 to 3-1/2 pts (1.5 to 2.6)	
	Rhizosphaera needlecast (spruces), Scirrhia brown spot (pines)	5-1/2 pts (4.125)	5-1/2 pts (4.125)	
Conifers (Pine, Spruce)	Cyclaneusma and Lophodermium needlecasts (pines)	2-3/4 to 5-1/2 pts (2.1 to 4.125)	2-3/4 to 5-1/2 pts (2.1 to 4.125)	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
	Rhabdocline needlecast (Douglas-fir)	1-1/2 to 2-3/4 pts (1.125 to 2.1)	1-1/2 to 2-3/4 pts (1.125 to 2.1)	
	Botrytis seedling blight, Phoma twig blight	1-1/2 to 2-3/4 pts (1.125 to 2.1)	1-1/2 to 2-3/4 pts (1.125 to 2.1)	

DO NOT apply more than 22 pints BRAVO 720 (16.5 lbs. a.i.) per acre during each growing season. The minimum re-treatment interval for established trees is 21 days. The minimum re-treatment interval in nursery beds is 7 days.

Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.

Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

CROP	DISEASES	<u>BRAVO 720 RATE PER</u> (lbs. a.i. per)		APPLICATION DIRECTIONS
		ACRE	100 GAL*	
	Autoecious needle rust (Weir's cushion) (spruce)	5-1/2 pts (4.125)	5-1/2 pts (4.125)	Begin applications when 10% of buds have broken and twice thereafter at 7-10 day intervals.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

MUSHROOMS: Verticillium brown spot and dry bubble - Apply 2.75 to 5.5 fl. oz. of Bravo 720 per 1,000 sq. ft. of mushroom bed. Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (5.5 fl.oz.) of Bravo 720 in the first application and the low rate (2.75 fl.oz.) of Bravo 720 in the second application. The first application should be made within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning. Do not apply within 5 days of first harvest. Make no more than two applications per cropping cycle. Do not apply more than 8.25 fl. oz. of Bravo 720 per cropping cycle.

BRAVO®, SuperWeatherStik™ and WEATHER STIK® are trademarks of a ZENECA company.

Benlate is a registered trademark of E.I. DuPont de Nemours and Company.

Copper-Count® is a registered trademark of Mineral Research and Development Corporation.

Dipel is a registered trademark of Abbott Laboratories.

Foil is a registered trademark of Ecogen, Inc.

Latron B-1956 and B-1956 are trademarks of Rohm and Haas Company.