

PROPOSED LABEL

Net Contents

Logo

BRAVO^R C/M

Agricultural Fungicide/ Bactericide

Active Ingredients:

Chlorothalonil (tetrachloroisophthalonitrile) . . .	27.0%
Copper Oxychloride*	45.8%
Maneb (manganese ethylene bisdithiocarbamate) . .	5.4%

Inert Ingredients:	<u>21.8%</u>
------------------------------	--------------

Total	100.0%
-----------------	--------

*Metallic copper equivalent 27.0%

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO


See side panel for additional precautionary statements

Precaucion al Usuario:

Si usted no lee ingles, no use este producto
hasta que la etiqueta haya sido explicado ampliamente.

FERMENTA PLANT PROTECTION COMPANY
7528 Auburn Road
Concord Township
P.O. Box 348
Painesville, OH 44077

(EPA Reg. No. 50534-191)



EPA Est. No. 50534-TX-01

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

WARNING

Causes substantial but temporary eye injury.

Do not get in eyes. Wear goggles or face shield when handling this product.

Avoid prolonged contact with skin.

Do not take internally.

Avoid breathing spray mist.

FIRST AID

Note to User: Wash exposed areas of skin with soap and warm water after handling or using.

In case of contact with eyes, flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes immediately.

Note to User: BRAVO C/M may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note to physician: Persons having a temporary allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARD

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Apply only to crops specified on label.

A BROAD SPECTRUM AGRICULTURAL FUNGICIDE/ BACTERICIDE

Read entire label carefully and use only as directed.

GENERAL INFORMATION

BRAVO C/M contains chlorothalonil which is an excellent fungicide when used according to label directions for control of a broad spectrum of diseases of tomatoes, peaches and nectarines. In addition, **BRAVO C/M** contains copper oxychloride plus maneb to provide control of bacterial spot and bacterial speck of tomatoes and bacterial canker, bud blast and bacterial spot of peaches and nectarines when applied as indicated in the following use directions.

BRAVO C/M is a broad spectrum protectant fungicide/ bactericide which is most effective when applied in dilute sprays. Applications should be started early before infection and continued as indicated in the Directions for Use. Application should be made at the recommended label rate and timing starting at the first sign of disease, first report of disease in the area, or when environmental conditions become favorable for disease development. Thorough, uniform coverage is essential for effective disease control.

DO NOT combine **BRAVO C/M** in the spray tank with surfactants, fertilizers or other pesticides unless your prior use has shown the combination physically compatible, effective and non-injurious under your conditions of use.

DO NOT combine **BRAVO C/M** with Dipel 4L or Triton AG-98 as phytotoxicity may result from the combination when applied to the crops on this label.

Dipel is a registered trademark of Abbott Laboratories;
Triton AG-98 is a trademark of Rohm and Haas Company.

The required amount of **BRAVO C/M** should be added slowly into the spray tank during filling. Keep agitator running when filling spray tank and during spray operations. DO NOT use on greenhouse-grown tomatoes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

Note to User: This product may produce temporary allergic side effects characterized by redness of eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician.

Note to User: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

TOMATO: BRAVO C/M is a specially designed product which controls bacterial and fungal diseases of tomato. The following diseases can be controlled through proper use of BRAVO C/M:

Bacterial Spot (*Xanthomonas*)
Bacterial Speck (*Pseudomonas*)
Early Blight
Late Blight (*Phytophthora infestans*)
Gray Leaf Mold (*Cladosporium*)
Gray Leaf Spot (*Stemphylium*)
Septoria Leafspot
Target Spot (*Corynespora*)
Alternaria Fruit Rot (Black Mold)
Gray Mold (*Botrytis*)
Rhizoctonia Fruit Rot
Anthracnose

Apply in sufficient water to obtain complete coverage of canopy, generally 50 to 125 gallons per acre. On upright staked tomatoes use application equipment capable of thoroughly covering the crop canopy, generally over-row sprayers with vertical boom orientation. On bush tomatoes, use nozzle placement that will insure thorough coverage of the canopy. Begin applications when dew or rain occur and disease threatens. Repeat applications on a weekly or semi-weekly (every 3-4 days) schedule, using the rates listed below:

<u>SCHEDULE</u>	<u>APPLICATION RATE</u> <u>(Per A·re)</u>	<u>RECOMMENDED SPRAY</u> <u>VOLUME (GAL/A)</u>	
		<u>Pre-Bloom</u>	<u>Bloom-Harvest</u>
Weekly	4 - 6 lbs	50 - 75	75 - 125
Semi-weekly	2 - 3 lbs	50 - 75	75 - 125

When conditions favor severe disease, use the highest rate listed for the application schedule selected. DO NOT apply more than 6 lbs/A of BRAVO C/M per application. DO NOT apply BRAVO C/M within 5 days of harvest.

PEACH, NECTARINE: The following diseases can be controlled through proper use of **BRAVO C/M**:

Leaf curl
Coryneum blight (shothole)
Bacterial canker, Bud blast (dead bud)
Bacterial spot

Apply **BRAVO C/M** in sufficient water and with proper calibration to obtain uniform coverage of the tree canopy, generally 50-200 gallons/A. Application with ground equipment is preferable because ground applications generally give better coverage. When treating non-bearing or immature trees, the lower rate of **BRAVO C/M** may be used.

For best control of Leaf curl and Coryneum blight and bacterial diseases, apply at leaf fall in late autumn. When conditions favor high disease levels, and for best control of bacterial diseases, use the high rate of application and apply once more in mid to late winter before budswell. If the late fall application is not practical, applications of **BRAVO C/M** for control of Leaf curl may be made prior to budswell the following spring. When Coryneum blight (shothole) and Bacterial spot occur, also apply at budbreak.

<u>DISEASES</u>	<u>RECOMMENDED APPLICATION RATE, / ACRE</u>	<u>SPRAY VOLUME (GAL/A)</u>
=====	=====	=====
Leaf curl, Coryneum blight, Bacterial canker, Bud blast (dead bud), Bacterial spot	6 - 10 lbs	50 - 200

Note to User: Do not rotate to crops other than those listed on labels within 12 months of the last treatment. After 12 months from the last application, leafy vegetables (i.e. spinach, lettuce, kale, etc.) may also be rotated.

REENTRY

Note to User: Do not enter treated area to perform hand labor within 24 hours of application unless protective clothing is worn.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information:

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Oral and written warnings must include the following information: **"WARNING.** Area treated with **BRAVO C/M** on (date of application). **Do not enter** without appropriate protective clothing until the sprays have dried. In case of accidental exposure, wash exposed area with plenty of water and get medical attention. For further information see 'Precautionary Statements' on the label".

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a cool place. Protect from excessive heat.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. **Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchability.** In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

FERMENTA PLANT PROTECTION COMPANY
7528 Auburn Road
Concord Township
P.O. Box 348
Painesville, Ohio 44077

EPA Reg. No. 50534-191

EPA Est. No. 50534-TX-01

12/86