

PM 22

50534-201

10/13

JUL 22 1993

Mr. Ralph P. Burton
ISK Biotech Corporation
P.O. Box 8000
Mentor, OH 44061-8000

Dear Mr. Burton:

Subject: Revised Warranty and Cucurbit Directions for Use
Bravo 500
EPA Registration No. 50534-8
Bravo W-75
EPA Registration No. 50534-23
Bravo 720
EPA Registration No. 50534-188
Bravo 825
EPA Registration No. 50534-201
Your Submissions Dated June 10, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

This acceptance of your label does not relieve you of your obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR Part 156, Subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely yours,



Cynthia Giles-Parker
Product Manager (22)
Fungicide Herbicide Branch
Registration Division (H7505C)

Enclosure

CROP	DISEASES	BRAVO 825 RATE PER ACRE	APPLICATION DIRECTIONS
Cucurbits: Cucumber, Cantaloupe, Muskmelon, Honeydew melon, Watermelon, Squash, Pumpkin	Target spot, Anthracnose, Downy mildew ----- Cercospora leafspot, Gummy stem blight (black rot), Alternaria leaf blight, Scab, Powdery mildew (<i>Sphaerotheca</i> only)	1.4 to 1.8 lbs 1.8 to 2.7 lbs	Use in sufficient water to obtain adequate coverage in applications when plants are in first true leaf stage. Under favorable conditions are favorable for disease development, applications at 7 day intervals. Under severe conditions, shorten spray interval. BRAVO 825 may be applied through sprinkler irrigation equipment (solid stream, move, motorized lateral move or center pivot system) in accordance with calibration directions preceding this section. NOTE: Do not combine BRAVO 825 in the spray mixture with any material other than water if applications are made to watermelons. Do not spray watermelons when environmental conditions are conducive to sunburn. Do not apply BRAVO to watermelons within 21 days of fruit maturity.
	Cucumber belly rot (<i>Rhizoctonia solani</i>)	7.6 lbs	Use BRAVO 825 in sufficient water to obtain runoff on soil surface. Make a single application when vines are in fruiting stage. BRAVO 825 may be applied through sprinkler irrigation as directed above.

ACCEPTED
with COMMENTS
by EPA Letter Dated:

JUL 22 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
registered, for the pesticide
classified under EPA Reg. No.
505-34-201

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. **SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO.** In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this company, including, but not limited to, incompatibility with products other than those set forth in the Directions for Use, weather conditions, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural practice.