· , PM 22 50534-201

18/2

MAR 9 1995

Mr. Ralph P. Burton
ISK Biotech Corporation
P.O. Box 8000

Mentor, OH 44061-8000

Dear Mr. Burton:

Subject: Add Cranberries

Bravo 825

EPA Registration No. 50534-201

Your Submission Dated February 6, 1995

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

- 1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) or 4(a)
 when the Agency requires all registrants of similar products to
 submit such data.
- 2. Submit one (1) copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

15)

Cynthia Giles-Parker Product Manager (22) Fungicide Herbicide Branch Registration Division (7505C)

Enclosure

PROPOSED LABELING

ISK BIOSCIENCES"

ACCEPTED
with COMMENTS
In EPA Letter Dated:

DIRECTIONS FOR THE USE OF

MAR 9 1995

BRAVO ULTREXTM (EPA Reg. No. 50534-201)

FOR THE CONTROL OF DISEASES ON CRANBERRY

50534-301

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply BRAVO ULTREX in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop.

Discuses	BRAVO ULTREX Rate per Acre	Application Directions
Fruit rots, Lophodermium leaf/twig blight	3.8 to 6.3 lb	Apply at early bloom and repeat at 10 to 14 day intervals. Under severe disease conditions use the 6.3 lb/acre rate on a 10 day schedule. DO NOT apply BRAVO ULTREX more than 3 times per season, or within 50 days before harvest. DO NOT apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application. BRAVO ULTREX may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.

Follow all applicable directions, restrictions, and precautions on the EPA-registered label.

NOTE: This labeling must be in the possession of the user at the time of pesticide application.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants to those persons lawfully acquiring title to this product that at the time of the first sale of this product by Seller that this product conformed to its chemical description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions under normal conditions of use and Buyers and users of this product assume the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in the Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practice.

2/95