

**Proposed Labeling****ISK BIOSCIENCES™****SUPPLEMENTAL DIRECTIONS FOR THE USE OF****BRAVO ULTREX®**

EPA Reg. No. 50534-201

**TO CONTROL DISEASES OF MUSHROOMS****DIRECTIONS FOR USE:**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**MUSHROOMS:** Verticillium brown spot and dry bubble - Apply 2.5 to 5 oz. of Bravo Ultrex® per 1,000 sq. ft. of mushroom bed. Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (5 oz.) of Bravo Ultrex in the first application and the low rate (2.5 oz.) of Bravo Ultrex in the second application. The first application should be made within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning. Do not apply within 5 days of first harvest. Make no more than two applications per cropping cycle. Do not apply more than 7.5 fl. oz. of Bravo Ultrex per cropping cycle.

**NOTE:** Follow all applicable directions, restrictions, and precautions on the EPA-registered label.

**NOTE:** This labeling must be in the possession of the user at the time of pesticide application.

3/15/96

