

**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Adora Clark
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419

FEB 26 2009

SUBJECT: Label Amendment to Request Application of Accepted Acute Inhalation Study
Bravo ZN
EPA Reg. No. 50534-204
Your Submission Dated October 10, 2008
Decision Number D-401488

Dear Ms. Clark:


The amended labeling referred to above, requesting that the approved acute inhalation study (MRID 45684001) with toxicological category III, also be applied toward the above mentioned submission and to remove the respirator restriction except for enclosed areas as outlined in the Reregistration Eligibility Document (RED; 1999), in connection with registration under the Federal Insecticide Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. The Hazards to Humans and Domestic Animals in both the container label and master label sections must read as follows: "Harmful if absorbed through skin. Harmful if inhaled. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and chemical resistant gloves (such as natural rubber, Selection Category A. Remove and wash contaminated clothing before reuse. Avoid breathing spray mist. Avoid contact with eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Submit one copy of your final printed labeling before you release the product for shipment.

If you have any questions regarding this correspondence, contact Rose Kearns of my staff by phone at 703-305-5611 or via email at kearns.rosemary@epa.gov or myself at 703-308-9443 or via email at kish.tony@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Rose M. Kearns".

Tony Kish
Product Manager Team 22
Fungicide Branch
Registration Division (7505P)

Enclosure

Bravo® Zn

Agricultural Fungicide

[Insert SuperWeatherStik® logo]

Active Ingredient:

Chlorothalonil (tetrachloroisophthalonitrile) 38.5%

Other Ingredients: 61.5%

Total: 100.0%

Contains 4.17 pounds chlorothalonil per gallon (500 grams per liter)

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

*Covered under U.S. Pat. No. 5,667,795

EPA Reg. No. 50534-204

EPA Est.

_____ gallons
Net Contents

**ACCEPTED
with COMMENTS
In EPA Letter Dated
FEB 26 2009**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.**

50534-204

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Note to Physician	
Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER	
For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING/AVISO

May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not breathe spray mist. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Mixers, Loaders, Applicators and all other handlers must wear:

- long-sleeved shirt and long pants
- chemical resistant gloves made of any waterproof material
- shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of GB BIOSCIENCES CORPORATION or Seller. To the extent permitted by applicable law,

Buyer and User agree to hold GB BIOSCIENCES and Seller harmless for any claims relating to such factors.

GB BIOSCIENCES warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or GB BIOSCIENCES, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, GB BIOSCIENCES MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall GB BIOSCIENCES be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF GB BIOSCIENCES AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF GB BIOSCIENCES OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

GB BIOSCIENCES and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of GB BIOSCIENCES.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Bravo Zn should be used only in accordance with recommendations on this label or in separately published GB BIOSCIENCES supplemental labeling recommendations for this product.

Do not apply this product in a way that will contact workers, other persons or pets, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical resistant gloves made of any waterproof material
- shoes plus socks
- protective eyewear

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted-entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

(1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS-required decontamination site intended for workers entering the treated area.

(2) Workers must be informed, in a manner they can understand:

- that residues in the treated area may be highly irritating to their eyes
- that they should take precautions, such as refraining from rubbing their eyes to keep the residues out of their eyes
- that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site, or using other readily available clean water
- how to operate the eyeflush container

GENERAL INFORMATION

Bravo Zn is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases.

Bravo Zn is recommended for use in programs which are compatible with the principles

of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

Bravo Zn is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk from disease resistance exhibit a single-site model of fungicidal action. Bravo Zn, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your Federal or State Cooperative Extension Service representatives for guidance on the proper use of Bravo Zn in programs which seek to minimize the occurrence of disease resistance to other fungicides.

Bravo Zn can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

General Precautions and Restrictions

Do not use on greenhouse-grown crops.

This product must not be applied within 150 feet for aerial applications, or 25 feet for ground applications of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Do not combine Bravo Zn in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. Do not combine Bravo Zn with Dipel[®], Latron B-1956[®] or Latron AG-98[®] as phytotoxicity may result from the combination when applied to some crops on this label.

Spray Drift Precautions

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information**.

Aerial Drift Reduction Advisory Information

[This section is advisory in nature and does not supercede the mandatory label requirements.]

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable conditions (See **Wind, Temperature**).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting the nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

Boom Length

For some use patterns, reducing the effective boom length to less than ¾ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 ft. above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller

drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION

Note: Slowly invert container several times to assure uniform mixture.

Dosage rates on this label indicate pints of Bravo Zn per acre, unless otherwise stated. Under conditions favoring disease development, the high rate specified and shortest application interval should be used.

The required amount of Bravo Zn should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of Bravo Zn in a clean container and add to the spray tank as it is being filled. Keep agitator running when

filling spray tank and during spray operations.

Apply Bravo Zn in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth.

For field and row crops, spray volume usually will range from 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications.

For tree and orchard crops, apply Bravo Zn in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. For conifers, the maximum volume is 100 gallons per acre.

Application and Calibration Techniques for Sprinkler Irrigation - Chemigation

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not apply this product through irrigation systems connected to a public water system. "Public water system" means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject Bravo Zn into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Bravo Zn may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Thoroughly mix recommended amount of Bravo Zn for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until Bravo Zn has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides;

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Beans (Dry) (except soybeans) bean, adzuki bean, broad bean, dry bean, lablab bean, navy bean, kidney bean, lima bean, moth bean, mung bean, pink bean, pinto bean, tepary bean, urd bean, yardlong catjang chickpea (garbanzo) cowpea lupin, grain lupin bean, rice bean, runner bean, jackbean pea, blackeyed pea, southern	Anthracnose (<i>Colletotrichum lindemuthianum</i>) Ascochyta blight (<i>A. phaseolorum</i>) Cercospora leaf blotch (<i>C. cruenta</i>) Downy mildew <i>Phytophthora nicotianae</i> Rust (<i>Uromyces appendiculatus</i>)	2 to 2½ (1.0 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications at first onset of disease, which may occur as early as 2 to 4 weeks before flowering. Repeat applications at 7 to 10 day intervals (the minimum re-treatment interval is 7 days). For use only on beans to be harvested dry with pods removed. Apply by ground, air or chemigation.

Specific Use Restrictions:

- Do not apply more than 11.5 pints of Bravo Zn (6 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days before harvest.

Blueberries	Suppression: Anthracnose (ripe rot) (<i>C. gloeosporoides</i>) Mummy berry (<i>M. vaccinicorymbosi</i>)	4¼ to 5¼ (2.25 to 3.0)	Bravo Zn should be integrated into an overall disease management strategy which includes alternation with a fungicide with a different mode of action. Diseases may only be suppressed and russetting may occur under heavy disease pressure or unfavorable environmental conditions. Apply in sufficient water to obtain adequate coverage, normally 20 to 100 gallons per acre. Begin applications at budbreak (green tip) and repeat at 10-day intervals through early bloom (the minimum re-treatment interval is 10 days). Under heavy disease pressure, use the higher rate. Apply by ground or air.
--------------------	--	---------------------------	--

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
	Rust (<i>Pucciniastrum vaccinii</i>) Septoria leaf spot (<i>Septoria albopunctata</i>)	4¼ to 5¼ (2.25 to 3.0)	Foliar Use After Harvest (after all berries are harvested): To maintain healthy leaves for the following season, apply in sufficient water to obtain adequate coverage (normally 20 to 100 gallons per acre). Repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days). Apply by ground or air.

Specific Use Restrictions:

- Do not apply more than 17 pints of Bravo Zn (9.0 lbs. a.i.) per acre during each growing season.
- Do not apply after full bloom (except for foliar use after harvest) or within 42 days of harvest.

Cabbage Chinese Cabbage (tight-headed varieties only) Cauliflower Broccoli Chinese Broccoli Brussels Sprouts	Alternaria leaf spot (<i>Alternaria</i> spp.) Downy mildew (<i>Peronospora parasitica</i>)	2¼ (1.125)	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control. Apply by ground, air or chemigation.
	Ring spot (California only)	2¼ (1.5)	For field-seeded Brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.

Specific Use Restrictions:

- Do not apply more than 23 pints of Bravo Zn (12 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

Carrot	Alternaria leaf blight (<i>A. dauci</i>) Cercospora leaf spot (<i>C. carotae</i>)	2¼ to 2¼ (1.125 to 1.5)	Use in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control. Apply by ground, air or chemigation.
--------	--	-------------------------	---

Specific Use Restrictions:

- Do not apply more than 29 pints of Bravo Zn (15 lbs. a.i.) per acre during each growing season.
- Bravo Zn may be applied the day of harvest.

Celery	Basal stalk rot (<i>Rhizoctonia solani</i>) Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>)	2¼ to 4¼ (1.5 to 2.25)	Use in sufficient water to obtain adequate coverage. Start applications when transplants are set in the field and repeat at a 7 day interval as needed to maintain control (the minimum re-treatment interval is 7 days). Apply by ground, air or chemigation.
--------	---	------------------------	---

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
	Suppression (7 day schedule): Pink rot (<i>Sclerotinia sclerotiorum</i>)	4¼ (2.25)	
	Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>)	2¼ to 2¾ (1.125 to 1.5) per 100 gal.	For celery seedbeds, apply in a spray volume of 125 gallons per acre twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.

Specific Use Restrictions:

- Do not apply more than 34.5 pints of Bravo Zn (18 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

Corn (Sweet), Corn (Grown for seed)	Helminthosporium leaf blights	1 1/8 to 2¼ (0.6 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at a 7 day interval as required to maintain control (the minimum re-treatment interval is 7 days). Under severe disease conditions, use 2¼ to 2¾ pints Bravo Zn per acre. Apply by ground, air or chemigation.
	Rust (<i>Puccinia</i> spp.)		

Specific Use Restrictions:

- Do not apply more than 17 pints of Bravo Zn (9 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest.
- Do not apply to sweet corn to be processed.
- Do not allow livestock to graze in treated fields.
- Do not ensile treated corn or use as livestock forage.

Cranberry	Fruit rots Lophodermium leaf/twig blight (<i>L. hypophyllum</i>)	5¼ to 9¼ (3.0 to 4.9)	Apply at early bloom and repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days). Under severe disease conditions, use the 9¼ pints per acre rate on a 10 day schedule. Apply by ground, air or chemigation. When applying by chemigation, use 300 gallons of water per acre through solid set systems only.
	Upright Dieback (<i>Phomopsis vaccinii</i>)	5¼ to 9¼ (3.0 to 4.9)	Apply in sufficient water to obtain coverage of uprights and runners. Make the first application before bloom, at the time shoots begin growth in the spring. Make additional applications at 10 to 14 day intervals. Apply by ground, air or chemigation. When applying by chemigation, use 300 gallons of water per acre through solid set systems only.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
------	------------------------	------------------------------------	------------------------

Specific Use Restrictions:

- Do not apply more than 29 pints of Bravo Zn (15 lbs. a.i.) per acre during each growing season.
- Do not apply within 50 days of harvest.
- Do not apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application.

Cucurbits Cantaloupe Cucumber Honeydew melon Muskmelon Pumpkin Squash Watermelon	Anthracnose <i>(Colletotrichum spp.)</i> Downy mildew <i>(Pseudoperonospora cubensis)</i> Target spot <i>(Corynespora cassicola)</i>	2¼ to 2¾ (1.125 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum re-treatment interval is 7 days). Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Bravo Zn to watermelons when any of the following conditions are present:
	Alternaria leaf blight <i>(A. cucumerina)</i> Alternaria leaf spot <i>(A. alternata)</i> Cercospora leaf spot <i>(C. citrullina)</i> Gummy stem blight /vine decline <i>(Didymella bryoniae)</i> Powdery mildew <i>(Sphaerotheca only)</i> Scab <i>(Cladosporium cucumerinum)</i>	2¼ to 4¼ (1.5 to 2.25)	<ol style="list-style-type: none"> 1. Intense heat and sunlight 2. Drought conditions 3. Poor vine canopy 4. Other crop and environmental conditions which may be conducive to increased natural sunburn Do not combine Bravo Zn with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use. Apply by ground, air or chemigation.

Specific Use Restrictions:

- Do not apply more than 30 pints of Bravo Zn (15.75 lbs. a.i.) per acre during each growing season.
- Bravo Zn may be applied the day of harvest.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Grasses Grown for Seed	Bipolaris and Drechslera leaf spots Glume blotch Leaf rust Septoria leaf spot Stem rust Stripe rust	1½ to 2¼ (0.75 to 1.125)	Use in sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat applications at 14 day intervals (the minimum re-treatment interval is 14 days). Apply by ground, air or chemigation.
	Selenophoma (eyespot)	1½ to 2¼ (0.75 to 1.5)	

Specific Use Restrictions:

- Do not apply more than 8.5 pints of Bravo Zn (4.5 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest.
- Do not allow livestock to graze in treated areas or feed hay produced before harvest. Feeding of treated plant parts after harvest of seed is allowed.

Mango	Anthracnose (<i>Colletotrichum</i> spp.)	2¾ to 5 (1.5 to 2.6)	Use a water volume of 20 to 300 gallons per acre. Begin applications at early bloom and repeat on a 7-14 day interval until early fruit development. Begin the season with the 2¾ pint rate on a 14-day interval (the minimum re-treatment interval is 7 days). If disease pressure is severe, use the higher rate and shorter interval. Apply by ground or air.
-------	---	----------------------	---

Specific Use Restrictions:

- Do not apply more than 46 pints of Bravo Zn (24 lbs. a.i.) per acre during each growing season.
- Do not apply within 21 days of harvest.

Mint (Indiana, Michigan and Wisconsin only)	Rust (<i>Puccinia menthae</i>) Septoria leaf spot (<i>S. menthae</i>)	2 (1.0)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground and aircraft applications. Begin applications when emerging plants are 4-8 inches high. Repeat applications at 7 to 10 day intervals to maintain control (the minimum re-treatment interval is 7 days).
---	--	---------	--

Specific Use Restrictions:

- Do not apply more than 5.75 pints of Bravo Zn (3 lbs. a.i.) per acre during each growing season.
- Do not apply within 80 days of harvest. Do not feed fresh or extracted mint hay from treated fields to livestock.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS			
Onion (Dry bulb) and Garlic	Botrytis leaf blight (<i>Botrytis</i> spp.)	1½ to 4¼ (0.75 to 2.25)	Apply in sufficient water to obtain thorough coverage of tops. Bravo Zn is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows:			
	Purple blotch (<i>Alternaria porri</i>)		Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard	
	Suppression: Botrytis neck rot		Rate per Acre	1½ pt.	2 pts.	4¼ pts.
	Downy mildew (<i>Peronospora destructor</i>)		Frequency	10 days	7 to 10 days	7 days
	For suppression of neck rot (<i>Botrytis</i> spp.) during storage, a minimum of three weekly applications prior to lifting, using 2 to 4¼ pints of Bravo Zn per acre, is recommended.					
The minimum re-treatment interval is 7 days.						
Apply by ground, air or chemigation.						

Specific Use Restrictions:

- Do not apply more than 29 pints of Bravo Zn (15 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

Onion (green bunching) Leek Shallots Onion and Garlic (grown for seed)	Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Downy mildew (<i>Peronospora destructor</i>)	2¼ to 4¼ (1.125 to 2.25)	Use in sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease (the minimum re-treatment interval is 7 days). Use the high rate and a 7 day schedule of applications when heavy dew or rain persist. Apply by ground, air or chemigation.			
---	--	--------------------------	---	--	--	--

Specific Use Restrictions:

- Do not apply more than 13 pints of Bravo Zn (6.75 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest on garlic.
- Do not apply within 14 days of harvest on green bunching onions, leeks or shallots.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Papaya	<p>Alternaria fruit spot (<i>A. alternata</i>)</p> <p>Anthrachnose (<i>Colletotrichum</i> spp.)</p> <p>Stem end rot (<i>A. alternata</i>, <i>Colletotrichum</i> spp.)</p>	2¼ to 4¼ (1.125 to 2.25)	Apply with ground equipment only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days).

Specific Use Restrictions:

- Do not apply more than 13 pints of Bravo Zn (6.75 lbs. a.i.) per acre during each growing season.
- Bravo Zn may be applied the day of harvest.

Parsnip	<p>Alternaria leaf spot (<i>Alternaria</i> spp.)</p> <p>Anthrachnose (<i>Colletotrichum</i> spp.)</p> <p>Botrytis blight (gray mold) (<i>B. cinerea</i>)</p> <p>Bottom rot (<i>Rhizoctonia</i>)</p> <p>Downy mildew (<i>Plasmopara crustosa</i>)</p>	2¼ to 2¾ (1.125 to 1.5)	<p>Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule (the minimum re-treatment interval is 7 days).</p> <p>Apply by ground, air or chemigation.</p>
---------	--	-------------------------	--

Specific Use Restrictions:

- Do not apply more than 11.5 pints of Bravo Zn (6 lbs. a.i.) per acre during each growing season.
- Do not apply within 10 days of harvest.

Passion Fruit	<p>Alternaria fruit and leaf spot (<i>Alternaria</i> spp.)</p> <p>Anthrachnose (<i>Colletotrichum</i> spp.)</p> <p>Cercospora fruit spot</p>	2¾ (1.5)	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin applications during late bloom and repeat at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days).
---------------	--	----------	---

Specific Use Restrictions:

- Do not apply more than 14.5 pints of Bravo Zn (7.5 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Peanut	Early leaf spot (<i>Cercospora arachidicola</i>)	1½ to 2¼ (0.75 to 1.125)	Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting; repeat at 14 day intervals (the minimum re-treatment interval is 14 days). When conditions favor late leaf spot or when rust or web blotch occur, apply 2¼ pints Bravo Zn per acre at 14 day intervals for the remainder of the season.
	Late leaf spot (<i>Cercosporidium personatum</i>)		
	Pepper spot (<i>Leptosphaerulina crassiasca</i>)		
	Rust (<i>Puccinia arachidis</i>)	2¼ (1.125)	Apply by ground, air, or chemigation. If applying by chemigation, use 2¼ pints Bravo Zn per acre. It is recommended to alternate chemigation applications with ground or aerial applications.
	Web blotch (<i>Phoma arachidicola</i>)		

Specific Use Restrictions:

- Do not apply more than 17 pints of Bravo Zn (9 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest.
- Do not allow livestock to graze in treated areas.
- Do not feed hay or threshings from treated fields to livestock.

Potato	Black dot (<i>Colletotrichum coccodes</i>)	1 1/8 (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Botrytis vine rot (<i>B. cinerea</i>)	- then -	
	Early blight (<i>Alternaria solani</i>)	1½ to 2¼ (0.75 to 1.125)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:
	Late blight (<i>Phytophthora infestans</i>)		

Specific Use Restrictions:

- Do not apply more than 21.5 pints of Bravo Zn (11.25 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Soybean	Anthracnose (<i>Colletotrichum truncatum</i>)		Apply in sufficient water to obtain complete coverage, using at least five gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. The minimum re-treatment interval is 14 days.
	Cercospora leaf blight (<i>C. kikuchii</i>)		
	Diaporthe pod and stem rot (<i>D. phaseolorum</i>)	2¼ to 3¼ (1.125 to 1.7)	Apply by ground, air, or chemigation.
	Frogeye leaf spot (<i>Cercospora sojina</i>)		Two application program: For determinate varieties, make the first application at R3 stage (early pod set) and the second application at R5 (seed formation). For indeterminate varieties, make the first application when largest pods are 1-1¼ inches in length. Make the second application 14 days later.
	Purple seed stain (<i>C. kikuchii</i>)		
	Septoria brown spot (<i>S. glycines</i>) Suppression: Rust (<i>Phakopsora pachyrhizi</i>)	1½ to 2¾ (0.75 to 1.5)	Three application program: For determinate varieties, make the first application at the beginning of flowering (R1), the second at early pod set (R3), and the third at beginning of seed formation (R5). For indeterminate varieties, make the first application one week after first flowering and continue applications at 14 day intervals.
Stem canker (<i>Diaporthe phaseolorum</i>)	1½ (0.75)	Apply in 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make the first application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease make a second and third application. Make all applications at 14 day intervals.	

Specific Use Restrictions:

- Do not apply more than 8.5 pints of Bravo Zn (4.5 lbs. a.i.) per acre during each growing season.
- Do not apply within 6 weeks of harvest.
- Do not feed hay or threshings from treated fields to livestock.

CROP	DISEASES (Pathogen)	Pts. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Tomato	FOLIAGE Early blight (<i>Alternaria solani</i>) Gray leaf mold (<i>Fluvia fluva</i> ; <i>Cladosporium</i>) Gray leaf spot (<i>Stemphyllium botryosum</i>) Late blight (<i>Phytophthora infestans</i>) Septoria leaf spot (<i>S. lycopersici</i>) Target spot (<i>Corynespora cassicola</i>)	2 to 2½ (1.0 to 1.5)	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Apply on a 7 to 10 day interval for foliage diseases. For fruit diseases, begin at fruit set and apply on a 7 to 14 day interval. Use the highest rate and shortest interval specified when disease conditions are severe. The minimum re-treatment interval is 7 days. Apply by ground, air, or chemigation.
	FRUIT Alternaria fruit rot (black mold) (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum spp.</i>) Botrytis gray mold (<i>B. cinerea</i>) Late blight fruit rot (<i>P. infestans</i>) Rhizoctonia fruit rot (<i>R. solani</i>)	2¾ to 4 (1.5 to 2.1)	

Specific Use Restrictions:

- Do not apply more than 28.5 pints of Bravo Zn (15 lbs. a.i.) per acre during each growing season.
- Bravo Zn may be applied the day of harvest.

Tree and Orchard Crops

Apply Bravo Zn in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. For conifers, the maximum volume is 100 gallons per acre.

Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, Bravo Zn may be applied with aircraft using at least 20 gallons of spray per acre. The minimum volume for application by aircraft to forest stands and Christmas trees is 10 gallons per acre.

When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of Bravo Zn listed may be used. Do not allow livestock to graze in treated areas.

CROP	DISEASES (Pathogen)	Pts. Product PER (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 gal.*	
Almonds	Blossom blight/brown rot (<i>Monilinia</i> spp.)	5¼ (3.0)	2 (1.0)	Use water volumes of 20-300 gallons per acre. For blossom blight, begin application at popcorn (pink bud) and follow with an application at full bloom. If weather is still conducive for disease development, another application may be made at petal fall. For control of shothole, make an application in the autumn at leaf fall. In the spring, make the first application at budbreak, followed by an application at shuck split to control nut infections and to control scab. Apply by ground or air.
	Scab (<i>Venturia carpophila</i>)			
	Shothole (<i>Wilsonomyces carpophilus</i>)			

Specific Use Restrictions:

- Do not apply more than 36 pints of Bravo Zn (18.75 lbs. a.i.) per acre during each growing season (leaf fall through shuck split).
- Do not apply within 150 days of harvest.

Filberts (Hazelnuts)	Eastern filbert blight (<i>Anisogramma anomala</i>)	5¼ (3.0)	2 (1.0)	Use a water volume of 20 to 300 gallons per acre. Begin applications at the onset of disease or when weather conditions favor disease development. Make applications on a 14-28 day schedule, using the shorter interval under heavy disease pressure (the minimum re-treatment interval is 14 days).
-------------------------	--	-------------	------------	---

Specific Use Restrictions:

- Do not apply more than 17 pints of Bravo Zn (9 lbs. a.i.) per acre during each growing season.
- Do not apply within 120 days of harvest.
- Do not apply through irrigation.
- Do not apply with oils, other pesticides, surfactants or fertilizers.
- Do not apply within one week of an oil-based pesticide application.

CROP	DISEASES (Pathogen)	Pts. Product PER (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 gal.*	
Peach Nectarine Apricot Cherry Plum Prune	Leaf curl (<i>Taphrina deformans</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)	4½ to 6 (2.3 to 3.1)	1½ to 2 (0.75 to 1.0)	For best control of both diseases, apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of Bravo Zn for control of leaf curl may be made at any time prior to budswell the following spring. Where shothole occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections. Apply by ground or air.
	Brown rot blossom blight <i>Monilinia</i> spp. Lacy (russet) scab (plum/prune)	4½ to 6 (2.3 to 3.1)	1½ to 2 (0.75 to 1.0)	Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Black knot (cherry, plum) (<i>Apiosporina morbosa</i>) Cherry leaf spot (<i>Blumeriella jaapii</i>) Scab (<i>Cladosporium carpophilum</i>)	4½ to 6 (2.3 to 3.1)	1½ to 2 (0.75 to 1.0)	In addition to the bloom application listed above, make one application at shuck split. Do not apply Bravo Zn after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10-14 days later. Apply by ground or air.

Specific Use Restrictions:

- Do not apply more than 29.5 pints of Bravo Zn (15.5 lbs. a.i.) per acre during each growing season.
- Bravo Zn may be applied the day of harvest.
- The minimum re-treatment interval is 10 days.

Pistachio	Botryosphaeria blight (<i>B. dothidea</i>) Suppression: Alternaria late blight (<i>A. alternata</i>)	8½ (4.5)	4¼ (2.25)	Use a water volume of 20 to 200 gallons per acre. Make the first application at the beginning of the blossom period followed by an application at full bloom. Make additional applications as required on a 28-day schedule. (The minimum re-treatment interval is 28 days). For Septoria and
-----------	---	-------------	--------------	---

CROP	DISEASES (Pathogen)	Pts. Product PER (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 gal.*	
	Botrytis blight (<i>B. cinerea</i>) Septoria leaf spot (<i>S. pistacina</i>)	5¼ to 8½ (3.0 to 4.5)	2¼ to 4¼ (1.50 to 2.25)	Botrytis, use the higher rate if disease pressure is severe. NOTE: Use of this product may result in speckling or reddening of the fruit hull (epicarp). This effect is superficial and has not resulted in any change in nut quality. Apply by ground or air.

Specific Use Restrictions:

- Do not apply more than 43 pints of Bravo Zn (22.5 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest.

Conifers (pines, spruces)	Swiss needlecast (<i>Phaeocryptopus gaeumannii</i>)	4 to 8 (2.1 to 4.125)	4 to 8 (2.1 to 4.125)	Single application technique: In Christmas tree plantations or forest stands, make one application in the spring when new shoot growth is ½ to 2 inches in length.
	Scleroderris canker (pines) (<i>Gremmeniella abietina</i>) Swiss needlecast (<i>P. gaeumannii</i>)	2¼ to 4 (1.125 to 2.1)	2¼ to 4 (1.125 to 2.1)	
	Sirococcus tip blight (<i>S. conigenus</i>)	2¼ to 5 (1.5 to 2.6)	2¼ to 5 (1.5 to 2.6)	
	Rhizosphaera needlecast (spruces) (<i>Rhizosphaera spp.</i>) Scirrhia brown spot (pines) (<i>Mycosphaerella dearnessii</i>)	8 (4.125)	8 (4.125)	
	Cyclaneusma and Lophodermium needlecasts (pines)	4 to 8 (2.1 to 4.125)	4 to 8 (2.1 to 4.125)	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.

CROP	DISEASES (Pathogen)	Pts. Product PER (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 gal.*	
	Rhabdocline needlecast (Douglas-fir)	2¼ to 4 (1.125 to 2.1)	2¼ to 4 (1.125 to 2.1)	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
	Botrytis seedling blight Phoma twig blight	2¼ to 4 (1.125 to 2.1)	2¼ to 4 (1.125 to 2.1)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
	Autoecious needle rust (Weir's cushion) (spruce)	8 (4.125)	8 (4.125)	Begin applications when 10% of buds have broken and twice thereafter at 7-10 day intervals.

Specific Use Restrictions:

- Do not apply more than 31.5 pints of Bravo Zn (16.5 lbs. a.i.) per acre during each growing season.
- The minimum re-treatment interval for established trees is 21 days.
- The minimum re-treatment interval in nursery beds is 7 days.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

MUSHROOMS: Verticillium brown spot and dry bubble - Apply 4 to 8 fl. oz. of Bravo Zn per 1,000 sq. ft. of mushroom bed. Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1,000 sq. ft. of mushroom bed. Make two applications. Apply the high rate (8 fl. oz.) of Bravo Zn in the first application and the low rate (4 fl. oz.) of Bravo Zn in the second application. The first application should be made within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. The second application should be made at pinning. Do not apply within 5 days of first harvest. Make no more than two applications per cropping cycle. Do not apply more than 12 fl. oz. of Bravo Zn per cropping cycle.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool place. Protect from excessive heat.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be

disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by alternative methods allowed by state and local authorities.

Returnable Refillable Containers

If Bravo Zn is packaged in a returnable refillable container, then, after use, do not rinse container. The contents of this container cannot be completely removed by cleaning. Return container intact to point of purchase.

This container must only be refilled with Bravo Zn. Refilling with materials other than Bravo Zn will result in contamination and may weaken container. Do not REUSE THE CONTAINER FOR ANY OTHER PURPOSE. Before refilling, inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn threads on closure devices. Check for leaks after refilling and before transport. Do not refill or transport a damaged or leaking container.

Bulk and Minibulk Containers

Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling or reconditioning, or clean in accordance with manufacturer's instructions.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Bravo[®], SuperWeatherStik[®] and the Syngenta logo are trademarks of a Syngenta Group Company.

Dipel[®] is a registered trademark of Valent BioSciences Corporation

Latron B-1956[®] and Latron AG-98[®] are trademarks of Dow AgroSciences LLC

For non-emergency (e.g., current product information), call
1-800-334-9481.

Manufactured for:
GB Biosciences Corporation
P.O. Box 18300
Greensboro, North Carolina 27419-8300

CONTAINER LABEL

Bravo® Zn

Agricultural Fungicide

[Insert SuperWeatherStik® logo]

Active Ingredient:

Chlorothalonil (tetrachloroisophthalonitrile) 38.5%

Other Ingredients: 61.5%

Total: 100.0%

Contains 4.17 pounds chlorothalonil per gallon (500 grams per liter)

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements in the Directions for Use section for information about this standard.

EPA Reg. No. 50534-204

EPA Est.

_____ gallons
Net Contents

FIRST AID	
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Note to Physician Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals**WARNING/AVISO**

May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Do not breathe spray mist. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool place. Protect from excessive heat.

Container Disposal

Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by alternative methods allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

Chemigation

Refer to supplemental labeling in attached booklet for use directions on chemigation. Do not apply this product through any type of irrigation system, unless the supplemental labeling on chemigation is followed.

Bravo[®], SuperWeatherStik[®] and the Syngenta logo are trademarks of a Syngenta Group Company.

For non-emergency (e.g., current product information), call
1-800-334-9481.

Manufactured for:
GB Biosciences Corporation
P.O. Box 18300
Greensboro, North Carolina 27419-8300

BRA ZN 50534-204 SEPT 08 AMEND CLEAN – df – 9/24/08
050534-00204.20080924.BRA_ZN-SEPT08-AMEND-CLEAN.PDF